

02/25/98

0535 U.S. PTO

UTILITY
PATENT APPLICATION
TRANSMITTAL

Attorney Docket No.

210121.428C3

First Inventor or Application Identifier

Jiangchun Xu

Title

COMPOUNDS FOR IMMUNODIAGNOSIS OF
PROSTATE CANCER AND METHODS FOR
THEIR USE

(Only for nonprovisional applications under 37 CFR § 1.53(b))

Express Mail Label No.

EM175835678US

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Box Patent Application
Assistant Commissioner for Patents
Washington, D.C. 20231

1. ☒ General Authorization Form
(Submit an original and a duplicate for fee processing)
2. ☒ Specification [Total Pages] 134
(preferred arrangement set forth below)
- Descriptive Title of the Invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to Microfiche Appendix
 - Background of the Invention

- Brief Summary of the Invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure

3. ☐ Drawing(s) (35 USC 113) [Total Sheets]

4. Oath or Declaration [Total Pages]

- a. ☐ Newly executed (original or copy)
- b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)
- i. ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting
inventor(s) named in the prior
application,
see 37 CFR 1.63(d)(2) and 1.33(b)

5. Incorporation By Reference (useable if box 4b is checked)
The entire disclosure of the prior application, from which
a copy of the oath or declaration is supplied under Box
4b, is considered as being part of the disclosure of the
accompanying application and is hereby incorporated by
reference therein.

6. ☐ Microfiche Computer Program (Appendix)
7. Nucleotide and Amino Acid Sequence Submission
(if applicable, all necessary)
- a. ☒ Computer-Readable Copy
- b. ☒ Paper Copy (identical to computer copy)
- c. ☒ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & document(s))
9. ☐ 37 CFR 3.73(b) Statement (when there is an assignee) ☐ Power of Attorney
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard
14. ☐ Small Entity Statement(s) ☐ Statement filed in prior application, Status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
16. ☒ Other: Certificate of Express Mail

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information below and in a preliminary amendment

☐ Continuation ☐ Divisional ☒ Continuation-In-Part (CIP) of prior Application No.: Not Available

Prior application information: Examiner _____ Group / Art Unit _____

☐ Claims the benefit of Provisional Application No. _____

CORRESPONDENCE ADDRESS

David J. Maki
Seed and Berry LLP
6300 Columbia Center
701 Fifth Avenue
Seattle, Washington 98104-7092
(206) 622-4900 phone
(206) 682-6031 fax

Respectfully submitted,

TYPED or PRINTED NAME David J. Maki

SIGNATURE

c:\CORIXA\1266.DOC

REGISTRATION NO. 31,392

Date FEBRUARY 25, 1998

COMPOUNDS FOR IMMUNODIAGNOSIS OF PROSTATE CANCER
AND METHODS FOR THEIR USE

REFERENCE TO RELATED APPLICATIONS

5 This application is a continuation-in-part of U.S. Patent Application No. _____, filed February 9, 1998 (Attorney Docket No. 210121.428C2), which is a continuation-in-part of U.S. Patent Application No. 08/904,809, filed August 1, 1997, which is a continuation-in-part of U.S. Patent Application No. 08/806,596, filed February 25, 1997.

10

TECHNICAL FIELD

 The present invention relates generally to the treatment and monitoring of prostate cancer. The invention is more particularly related to polypeptides comprising at least a portion of a prostate protein. Such polypeptides may be used for
15 the production of compounds, such as antibodies, useful for diagnosing and monitoring the progression of prostate cancer, and possibly other tumor types, in a patient.

BACKGROUND OF THE INVENTION

 Prostate cancer is the most common form of cancer among males, with
20 an estimated incidence of 30% in men over the age of 50. Overwhelming clinical evidence shows that human prostate cancer has the propensity to metastasize to bone, and the disease appears to progress inevitably from androgen dependent to androgen refractory status, leading to increased patient mortality. This prevalent disease is currently the second leading cause of cancer death among men in the U.S.

25 In spite of considerable research into diagnosis and therapy of the disease, prostate cancer remains difficult to detect and to treat. Commonly, treatment is based on surgery and/or radiation therapy, but these methods are ineffective in a significant percentage of cases. Two previously identified prostate specific proteins - prostate specific antigen (PSA) and prostatic acid phosphatase (PAP) - have limited
30 diagnostic and therapeutic potential. For example, PSA levels do not always correlate well with the presence of prostate cancer, being positive in a percentage of non-prostate cancer cases, including benign prostatic hyperplasia (BPH). Furthermore, PSA

measurements correlate with prostate volume, and do not indicate the level of metastasis.

Accordingly, there remains a need in the art for improved and diagnostic methods for prostate cancer.

5

SUMMARY OF THE INVENTION

The present invention provides methods for immunodiagnosis of prostate cancer, together with kits for use in such methods. Polypeptides are disclosed which comprise at least an immunogenic portion of a prostate tumor protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein the prostate tumor protein comprises an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-11, 115-171, 173-175, 177, 179-224 and variants thereof. Such polypeptides may be usefully employed in the diagnosis and monitoring of prostate cancer.

In one specific aspect of the present invention, methods are provided for detecting prostate cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the above polypeptides; and (b) detecting in the sample a protein or polypeptide that binds to the binding agent. In preferred embodiments, the binding agent is an antibody, most preferably a monoclonal antibody.

In related aspects, methods are provided for monitoring the progression of prostate cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the above polypeptides; (b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent; (c) repeating steps (a) and (b); and comparing the amounts of polypeptide detected in steps (b) and (c).

Within related aspects, the present invention provides antibodies, preferably monoclonal antibodies, that bind to the inventive polypeptides, as well as diagnostic kits comprising such antibodies, and methods of using such antibodies to inhibit the development of prostate cancer.

The present invention further provides methods for detecting prostate cancer comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with a first and a second oligonucleotide primer in a polymerase chain reaction, at least one of the oligonucleotide primers being specific for a DNA molecule that encodes one of the above polypeptides; and (c) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers. In a preferred embodiment, at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule having a partial sequence selected from the group consisting of SEQ ID Nos: 2-3, 5-107, 109-11, 115-171, 173-175, 177 and 179-224.

In a further aspect, the present invention provides a method for detecting prostate cancer in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a DNA molecule that encodes one of the above polypeptides; and (c) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe. Preferably, the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule having a partial sequence selected from the group consisting of SEQ ID Nos: : 2-3, 5-107, 109-11, 115-171, 173-175, 177 and 179-224.

In related aspects, diagnostic kits comprising the above oligonucleotide probes or primers are provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for the immunodiagnosis and monitoring of prostate cancer. The inventive compositions are generally polypeptides that comprise at least a portion of a prostate tumor protein. Also included within the present invention are molecules (such as an antibody or fragment thereof) that bind to the inventive polypeptides. Such molecules are referred to herein as "binding agents."

In particular, the subject invention discloses polypeptides comprising at least a portion of a human prostate tumor protein, or a variant thereof such a protein, wherein the prostate tumor protein includes an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-11, 115-171, 173-175, 177, 179-224, the complements of said nucleotide sequences and variants thereof. As used herein, the term "polypeptide" encompasses amino acid chains of any length, including full length proteins, wherein the amino acid residues are linked by covalent peptide bonds. Thus, a polypeptide comprising a portion of one of the above prostate proteins may consist entirely of the portion, or the portion may be present within a larger polypeptide that contains additional sequences. The additional sequences may be derived from the native protein or may be heterologous, and such sequences may be immunoreactive and/or antigenic.

As used herein, an "immunogenic portion" of a human prostate tumor protein is a portion that is capable of eliciting an immune response in a patient inflicted with prostate cancer and as such binds to antibodies present within sera from a prostate cancer patient. Immunogenic portions of the proteins described herein may thus be identified in antibody binding assays. Such assays may generally be performed using any of a variety of means known to those of ordinary skill in the art, as described, for example, in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988. For example, a polypeptide may be immobilized on a solid support (as described below) and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A. Alternatively, a polypeptide may be used to generate monoclonal and polyclonal antibodies for use in detection of the polypeptide in blood or other fluids of prostate cancer patients.

The compositions and methods of the present invention also encompass variants of the above polypeptides and DNA molecules. A polypeptide "variant," as used herein, is a polypeptide that differs from the recited polypeptide only in conservative substitutions and/or modifications, such that the therapeutic, antigenic and/or immunogenic properties of the polypeptide are retained. Polypeptide variants

preferably exhibit at least about 70%, more preferably at least about 90% and most preferably at least about 95% identity to the identified polypeptides. For prostate tumor polypeptides with immunoreactive properties, variants may, alternatively, be identified by modifying the amino acid sequence of one of the above polypeptides, and evaluating the immunoreactivity of the modified polypeptide. For prostate tumor polypeptides useful for the generation of diagnostic binding agents, a variant may be identified by evaluating a modified polypeptide for the ability to generate antibodies that detect the presence or absence of prostate cancer. Such modified sequences may be prepared and tested using, for example, the representative procedures described herein.

As used herein, a "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his.

Variants may also, or alternatively, contain other modifications, including the deletion or addition of amino acids that have minimal influence on the antigenic properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (*e.g.*, poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

A nucleotide "variant" is a sequence that differs from the recited nucleotide sequence in having one or more nucleotide deletions, substitutions or additions. Such modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983). Nucleotide variants may be naturally occurring allelic variants, or non-naturally occurring variants. Variant nucleotide sequences preferably exhibit at least about 70%, more preferably at least about 80% and

most preferably at least about 90% identity to the recited sequence. Such variant nucleotide sequences will generally hybridize to the recite nucleotide sequence under stringent conditions. As used herein, "stringent conditions" refers to prewashing in a solution of 6X SSC, 0.2% SDS; hybridizing at 65 °C, 6X SSC, 0.2% SDS overnight; 5 followed by two washes of 30 minutes each in 1X SSC, 0.1% SDS at 65 °C and two washes of 30 minutes each in 0.2X SSC, 0.1% SDS at 65 °C.

"Polypeptides" as used herein also include combination, or fusion, polypeptides. A "combination polypeptide" is a polypeptide comprising at least one of the above immunogenic portions and one or more additional immunogenic prostate 10 tumor-specific sequences, which are joined via a peptide linkage into a single amino acid chain. The sequences may be joined directly (*i.e.*, with no intervening amino acids) or may be joined by way of a linked sequence (*e.g.*, Gly-Cys-Gly) that does not significantly diminish the immunogenic properties of the component polypeptides.

The prostate tumor proteins of the present invention, and DNA 15 molecules encoding such proteins, may be isolated from prostate tumor tissue using any of a variety of methods well known in the art. DNA sequences corresponding to a gene (of a portion thereof) encoding one of the inventive prostate tumor proteins may be isolated from a prostate tumor cDNA library using a subtraction technique as described in detail below. Examples of such DNA sequences are provided in SEQ ID Nos: 1-107, 20 109-111, 115-171, 173-175, 177 and 179-224. Partial DNA sequences thus obtained may be used to design oligonucleotide primers for the amplification of full-length DNA sequences in a polymerase chain reaction (PCR), using techniques well known in the art (see, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989). Once a DNA sequence 25 encoding a polypeptide is obtained, any of the above modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983).

The prostate tumor polypeptides disclosed herein may also be generated by synthetic or recombinant means. Synthetic polypeptides having fewer than about 30 100 amino acids, and generally fewer than about 50 amino acids, may be generated using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase

techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain (see, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963). Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

Alternatively, any of the above polypeptides may be produced recombinantly by inserting a DNA sequence that encodes the polypeptide into an expression vector and expressing the protein in an appropriate host. Any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant polypeptides of this invention. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line, such as CHO cells. The DNA sequences expressed in this manner may encode naturally occurring polypeptides, portions of naturally occurring polypeptides, or other variants thereof.

In general, regardless of the method of preparation, the polypeptides disclosed herein are prepared in substantially pure form (*i.e.*, the polypeptides are homogenous as determined by amino acid composition and primary sequence analysis). Preferably, the polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. In certain embodiments, described in more detail below, the substantially pure polypeptides are incorporated into pharmaceutical compositions or vaccines for use in one or more of the methods disclosed herein.

In a related aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, a polypeptide of the present invention and a known prostate antigen, together with variants of such fusion proteins. The fusion proteins of the present invention may also include a linker peptide between the first and second polypeptides.

A DNA sequence encoding a fusion protein of the present invention is constructed using known recombinant DNA techniques to assemble separate DNA

5 translation of the two DNA sequences into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons require to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

30 Polypeptides and/or fusion proteins of the present invention may be used to generate binding agents, such as antibodies or fragments thereof, that are capable of detecting metastatic human prostate tumors. Binding agents of the present invention

may generally be prepared using methods known to those of ordinary skill in the art, including the representative procedures described herein. Binding agents are capable of differentiating between patients with and without prostate cancer, using the representative assays described herein. In other words, antibodies or other binding agents raised against a prostate tumor protein, or a suitable portion thereof, will generate a signal indicating the presence of primary or metastatic prostate cancer in at least about 20% of patients afflicted with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without primary or metastatic prostate cancer. Suitable portions of such prostate tumor proteins are portions that are able to generate a binding agent that indicates the presence of primary or metastatic prostate cancer in substantially all (*i.e.*, at least about 80%, and preferably at least about 90%) of the patients for which prostate cancer would be indicated using the full length protein, and that indicate the absence of prostate cancer in substantially all of those samples that would be negative when tested with full length protein. The representative assays described below, such as the two-antibody sandwich assay, may generally be employed for evaluating the ability of a binding agent to detect metastatic human prostate tumors.

The ability of a polypeptide and/or fusion protein prepared as described herein to generate antibodies capable of detecting primary or metastatic human prostate tumors may generally be evaluated by raising one or more antibodies against the polypeptide (using, for example, a representative method described herein) and determining the ability of such antibodies to detect such tumors in patients. This determination may be made by assaying biological samples from patients with and without primary or metastatic prostate cancer for the presence of a polypeptide that binds to the generated antibodies. Such test assays may be performed, for example, using a representative procedure described below. Polypeptides that generate antibodies capable of detecting at least 20% of primary or metastatic prostate tumors by such procedures are considered to be useful in assays for detecting primary or metastatic human prostate tumors. Polypeptide specific antibodies may be used alone or in combination to improve sensitivity.

Polypeptides and/or fusion proteins capable of detecting primary or metastatic human prostate tumors may be used as markers for diagnosing prostate

03030606, 022398

cancer or for monitoring disease progression in patients. In one embodiment, prostate cancer in a patient may be diagnosed by evaluating a biological sample obtained from the patient for the level of one or more of the above polypeptides, relative to a predetermined cut-off value. As used herein, suitable "biological samples" include
5 blood, sera, urine and/or prostate secretions.

The level of one or more of the above polypeptides may be evaluated using any binding agent specific for the polypeptide(s). A "binding agent," in the context of this invention, is any agent (such as a compound or a cell) that binds to a polypeptide as described above. As used herein, "binding" refers to a noncovalent
10 association between two separate molecules (each of which may be free (*i.e.*, in solution) or present on the surface of a cell or a solid support), such that a "complex" is formed. Such a complex may be free or immobilized (either covalently or noncovalently) on a support material. The ability to bind may generally be evaluated by determining a binding constant for the formation of the complex. The binding constant
15 is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind" in the context of the present invention when the binding constant for complex formation exceeds about 10^3 L/mol. The binding constant may be determined using methods well known to those of ordinary skill in the art.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome with or without a peptide component, an RNA molecule or a peptide. In a preferred embodiment, the binding partner is an antibody, or a fragment thereof. Such antibodies may be polyclonal, or monoclonal. In addition, the antibodies may be single chain, chimeric, CDR-grafted or humanized.
25 Antibodies may be prepared by the methods described herein and by other methods well known to those of skill in the art.

There are a variety of assay formats known to those of ordinary skill in the art for using a binding partner to detect polypeptide markers in a sample. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory,
30 1988. In a preferred embodiment, the assay involves the use of binding partner immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a second

binding partner that contains a reporter group. Suitable second binding partners include antibodies that bind to the binding partner/polypeptide complex. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding partner after incubation of the binding partner with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding partner is indicative of the reactivity of the sample with the immobilized binding partner.

The solid support may be any material known to those of ordinary skill in the art to which the antigen may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an

aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay.

5 This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a second antibody (containing a reporter group) capable of binding to a
10 different site on the polypeptide is added. The amount of second antibody that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically
15 blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact
20 time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with prostate cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve
25 equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second
30 antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and

biotin. The conjugation of antibody to reporter group may be achieved using standard methods known to those of ordinary skill in the art.

The second antibody is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide.

5 An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound second antibody is then removed and bound second antibody is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally
10 appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction
15 products.

To determine the presence or absence of prostate cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value is the average mean signal obtained when the
20 immobilized antibody is incubated with samples from patients without prostate cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for prostate cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for*
25 *Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the
30 most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or

to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for prostate cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the antibody is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized antibody as the sample passes through the membrane. A second, labeled antibody then binds to the antibody-polypeptide complex as a solution containing the second antibody flows through the membrane. The detection of bound second antibody may then be performed as described above. In the strip test format, one end of the membrane to which antibody is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second antibody and to the area of immobilized antibody. Concentration of second antibody at the area of immobilized antibody indicates the presence of prostate cancer. Typically, the concentration of second antibody at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of antibody immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the antigens or antibodies of the present invention. The above descriptions are intended to be exemplary only.

In another embodiment, the above polypeptides may be used as markers for the progression of prostate cancer. In this embodiment, assays as described above for the diagnosis of prostate cancer may be performed over time, and the change in the level of reactive polypeptide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, prostate cancer is progressing in those patients in whom the level of

polypeptide detected by the binding agent increases over time. In contrast, prostate cancer is not progressing when the level of reactive polypeptide either remains constant or decreases with time.

Antibodies for use in the above methods may be prepared by any of a
 5 variety of techniques known to those of ordinary skill in the art. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the antigenic polypeptide is initially injected into any of a wide variety of mammals (*e.g.*, mice, rats, rabbits, sheep and goats). In this step, the polypeptides of this invention may serve as the immunogen without
 10 modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal
 15 antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for the antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J.*
 20 *Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a
 25 myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine,
 30 aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and tested for binding

activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Monoclonal antibodies of the present invention may also be used as therapeutic reagents, to diminish or eliminate prostate tumors. The antibodies may be used on their own (for instance, to inhibit metastases) or coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in

chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described
5 in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody
10 portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a
15 photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one
20 embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for
25 attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may
30 also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating

compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For
5 example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the
10 precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

Diagnostic reagents of the present invention may also comprise DNA sequences encoding one or more of the above polypeptides, or one or more portions thereof. For example, at least two oligonucleotide primers may be employed in a
15 polymerase chain reaction (PCR) based assay to amplify prostate tumor-specific cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for a DNA molecule encoding a prostate tumor protein of the present invention. The presence of the amplified cDNA is then detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes specific for a
20 DNA molecule encoding a prostate tumor protein of the present invention may be used in a hybridization assay to detect the presence of an inventive polypeptide in a biological sample.

As used herein, the term "oligonucleotide primer/probe specific for a DNA molecule" means an oligonucleotide sequence that has at least about 80%,
25 preferably at least about 90% and more preferably at least about 95%, identity to the DNA molecule in question. Oligonucleotide primers and/or probes which may be usefully employed in the inventive diagnostic methods preferably have at least about 10-40 nucleotides. In a preferred embodiment, the oligonucleotide primers comprise at least about 10 contiguous nucleotides of a DNA molecule having a sequence selected
30 from SEQ ID Nos: 1-107, 109-111, 115-171, 173-175, 177 and 179-224. Preferably, oligonucleotide probes for use in the inventive diagnostic methods comprise at least about 15 contiguous oligonucleotides of a DNA molecule having a sequence provided

in SEQ ID Nos: 1-107, 109-111, 115-171, 173-175, 177 and 179-224. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis et al. *Ibid*; Ehrlich, *Ibid*). Primers or probes may thus be used to detect prostate tumor-specific sequences in biological samples, including blood, semen,
5 prostate tissue and/or prostate tumor tissue.

Polypeptides of the present invention that comprise an immunogenic portion of a prostate tumor protein may also be used for immunotherapy of prostate cancer, wherein the polypeptide stimulates the patient's own immune response to prostate tumor cells. In further aspects, the present invention provides methods for
10 using one or more of the immunoreactive polypeptides encoded by a DNA molecule having a sequence provided in SEQ ID NO: 1-107, 109-111, 115-171, 173-175, 177 and 179-224 (or DNA encoding such polypeptides) for immunotherapy of prostate cancer in a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with a disease, or may be free of detectable disease.
15 Accordingly, the above immunoreactive polypeptides may be used to treat prostate cancer or to inhibit the development of prostate cancer. The polypeptides may be administered either prior to or following surgical removal of primary tumors and/or treatment by administration of radiotherapy and conventional chemotherapeutic drugs.

In these aspects, the polypeptide is generally present within a
20 pharmaceutical composition and/or a vaccine. Pharmaceutical compositions may comprise one or more polypeptides, each of which may contain one or more of the above sequences (or variants thereof), and a physiologically acceptable carrier. The vaccines may comprise one or more of such polypeptides and a non-specific immune response enhancer, such as an adjuvant, biodegradable microsphere (*e.g.*, polylactic
25 galactide) or a liposome (into which the polypeptide is incorporated). Pharmaceutical compositions and vaccines may also contain other epitopes of prostate tumor antigens, either incorporated into a combination polypeptide (*i.e.*, a single polypeptide that contains multiple epitopes) or present within a separate polypeptide.

Alternatively, a pharmaceutical composition or vaccine may contain
30 DNA encoding one or more of the above polypeptides, such that the polypeptide is generated *in situ*. In such pharmaceutical compositions and vaccines, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in

the art, including nucleic acid expression systems, bacteria and viral expression systems. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an epitope of a prostate cell antigen on its cell surface. In a preferred embodiment, the DNA may be introduced using a viral expression system (*e.g.*, vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., *PNAS* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *PNAS* 91:215-219, 1994; Kass-Eisler et al., *PNAS* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in published PCT application WO 90/11092, and Ulmer et al., *Science* 259:1745-1749, 1993, reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

Routes and frequency of administration, as well as dosage, will vary from individual to individual and may parallel those currently being used in immunotherapy of other diseases. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Between 1 and 10 doses may be administered over a 3-24 week period. Preferably, 4 doses are administered, at an interval of 3 months, and booster administrations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of polypeptide or DNA that is effective to raise an immune response (cellular and/or humoral) against prostate tumor cells in a treated patient. A suitable immune response is at least 10-50% above the basal (*i.e.*, untreated) level. In

general, the amount of polypeptide present in a dose (or produced *in situ* by the DNA in a dose) ranges from about 1 pg to about 100 mg per kg of host, typically from about 10 pg to about 1 mg, and preferably from about 100 pg to about 1 μ g. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.01 mL to about 5 mL.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a lipid, a wax and/or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and/or magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactic glycolide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Any of a variety of non-specific immune response enhancers may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a nonspecific stimulator of immune response, such as lipid A, *Bordella pertussis* or *Mycobacterium tuberculosis*. Such adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI) and Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ).

Polypeptides disclosed herein may also be employed in *ex vivo* treatment of prostate cancer. For example, cells of the immune system, such as T cells, may be isolated from the peripheral blood of a patient, using a commercially available cell separation system, such as CellPro Incorporated's (Bothell, WA) CEPRATE™ system (see U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). The separated cells are stimulated with one or more of the immunoreactive polypeptides contained within a delivery vehicle, such as a microsphere, to provide antigen-specific T cells. The population of tumor antigen-

specific T cells is then expanded using standard techniques and the cells are administered back to the patient.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLES

EXAMPLE 1

10 ISOLATION AND CHARACTERIZATION OF PROSTATE TUMOR POLYPEPTIDES

This Example describes the isolation of prostate tumor polypeptides from a prostate tumor cDNA library.

A human prostate tumor cDNA expression library was constructed from prostate tumor poly A⁺ RNA using a Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning kit (BRL Life Technologies, Gaithersburg, MD 20897) following the manufacturer's protocol. Specifically, prostate tumor tissues were homogenized with polytron (Kinematica, Switzerland) and total RNA was extracted using Trizol reagent (BRL Life Technologies) as directed by the manufacturer. The poly A⁺ RNA was then purified using a Qiagen oligotex spin column mRNA purification kit (Qiagen, Santa Clarita, CA 91355) according to the manufacturer's protocol. First-strand cDNA was synthesized using the NotI/Oligo-dT18 primer. Double-stranded cDNA was synthesized, ligated with EcoRI/BAXI adaptors (Invitrogen, San Diego, CA) and digested with NotI. Following size fractionation with Chroma Spin-1000 columns (Clontech, Palo Alto, CA 94303), the cDNA was ligated into the EcoRI/NotI site of pCDNA3.1 (Invitrogen) and transformed into ElectroMax *E. coli* DH10B cells (BRL Life Technologies) by electroporation.

Using the same procedure, a normal human pancreas cDNA expression library was prepared from a pool of six tissue specimens (Clontech). The cDNA libraries were characterized by determining the number of independent colonies, the percentage of clones that carried insert, the average insert size and by sequence analysis. The prostate tumor library contained 1.64×10^7 independent colonies, with 70% of

clones having an insert and the average insert size being 1745 base pairs. The normal pancreas cDNA library contained 3.3×10^6 independent colonies, with 69% of clones having inserts and the average insert size being 1120 base pairs. For both libraries, sequence analysis showed that the majority of clones had a full length cDNA sequence and were synthesized from mRNA, with minimal rRNA and mitochondrial DNA contamination.

cDNA library subtraction was performed using the above prostate tumor and normal pancreas cDNA libraries, as described by Hara et al. (*Blood*, 84:189-199, 1994) with some modifications. Specifically, a prostate tumor-specific subtracted cDNA library was generated as follows. Normal pancreas cDNA library (70 μ g) was digested with EcoRI, NotI, and SfuI, followed by a filling-in reaction with DNA polymerase Klenow fragment. After phenol-chloroform extraction and ethanol precipitation, the DNA was dissolved in 100 μ l of H₂O, heat-denatured and mixed with 100 μ l (100 μ g) of Photoprobe biotin (Vector Laboratories, Burlingame, CA). As recommended by the manufacturer, the resulting mixture was irradiated with a 270 W sunlamp on ice for 20 minutes. Additional Photoprobe biotin (50 μ l) was added and the biotinylation reaction was repeated. After extraction with butanol five times, the DNA was ethanol-precipitated and dissolved in 23 μ l H₂O to form the driver DNA.

To form the tracer DNA, 10 μ g prostate tumor cDNA library was digested with BamHI and XhoI, phenol chloroform extracted and passed through Chroma spin-400 columns (Clontech). Following ethanol precipitation, the tracer DNA was dissolved in 5 μ l H₂O. Tracer DNA was mixed with 15 μ l driver DNA and 20 μ l of 2 x hybridization buffer (1.5 M NaCl/10 mM EDTA/50 mM HEPES pH 7.5/0.2% sodium dodecyl sulfate), overlaid with mineral oil, and heat-denatured completely. The sample was immediately transferred into a 68 °C water bath and incubated for 20 hours (long hybridization [LH]). The reaction mixture was then subjected to a streptavidin treatment followed by phenol/chloroform extraction. This process was repeated three more times. Subtracted DNA was precipitated, dissolved in 12 μ l H₂O, mixed with 8 μ l driver DNA and 20 μ l of 2 x hybridization buffer, and subjected to a hybridization at 68 °C for 2 hours (short hybridization [SH]). After removal of biotinylated double-stranded DNA, subtracted cDNA was ligated into BamHI/XhoI site of chloramphenicol

resistant pBCSK⁺ (Stratagene, La Jolla, CA 92037) and transformed into ElectroMax *E. coli* DH10B cells by electroporation to generate a prostate tumor specific subtracted cDNA library (prostate subtraction 1).

To analyze the subtracted cDNA library, plasmid DNA was prepared from 100 independent clones, randomly picked from the subtracted prostate tumor specific library and grouped based on insert size. Representative cDNA clones were further characterized by DNA sequencing with a Perkin Elmer/Applied Biosystems Division Automated Sequencer Model 373A (Foster City, CA). Six cDNA clones, hereinafter referred to as F1-13, F1-12, F1-16, H1-1, H1-9 and H1-4, were shown to be abundant in the subtracted prostate-specific cDNA library. The determined 3' and 5' cDNA sequences for F1-12 are provided in SEQ ID NO: 2 and 3, respectively, with determined 3' cDNA sequences for F1-13, F1-16, H1-1, H1-9 and H1-4 being provided in SEQ ID No: 1 and 4-7, respectively.

The cDNA sequences for the isolated clones were compared to known sequences in the gene bank using the EMBL and GenBank databases (release 96). Four of the prostate tumor cDNA clones, F1-13, F1-16, H1-1, and H1-4, were determined to encode the following previously identified proteins: prostate specific antigen (PSA), human glandular kallikrein, human tumor expression enhanced gene, and mitochondria cytochrome C oxidase subunit II. H1-9 was found to be identical to a previously identified human autonomously replicating sequence. No significant homologies to the cDNA sequence for F1-12 were found.

Subsequent studies led to the isolation of a full-length cDNA sequence for F1-12. This sequence is provided in SEQ ID NO: 107, with the corresponding predicted amino acid sequence being provided in SEQ ID NO: 108.

To clone less abundant prostate tumor specific genes, cDNA library subtraction was performed by subtracting the prostate tumor cDNA library described above with the normal pancreas cDNA library and with the three most abundant genes in the previously subtracted prostate tumor specific cDNA library: human glandular kallikrein, prostate specific antigen (PSA), and mitochondria cytochrome C oxidase subunit II. Specifically, 1 μ g each of human glandular kallikrein, PSA and mitochondria cytochrome C oxidase subunit II cDNAs in pCDNA3.1 were added to the driver DNA and subtraction was performed as described above to provide a second

subtracted cDNA library hereinafter referred to as the "subtracted prostate tumor specific cDNA library with spike".

Twenty-two cDNA clones were isolated from the subtracted prostate tumor specific cDNA library with spike. The determined 3' and 5' cDNA sequences for the clones referred to as J1-17, L1-12, N1-1862, J1-13, J1-19, J1-25, J1-24, K1-58, K1-63, L1-4 and L1-14 are provided in SEQ ID Nos: 8-9, 10-11, 12-13, 14-15, 16-17, 18-19, 20-21, 22-23, 24-25, 26-27 and 28-29, respectively. The determined 3' cDNA sequences for the clones referred to as J1-12, J1-16, J1-21, K1-48, K1-55, L1-2, L1-6, N1-1858, N1-1860, N1-1861, N1-1864 are provided in SEQ ID Nos: 30-40, respectively. Comparison of these sequences with those in the gene bank as described above, revealed no significant homologies to three of the five most abundant DNA species, (J1-17, L1-12 and N1-1862; SEQ ID Nos: 8-9, 10-11 and 12-13, respectively). Of the remaining two most abundant species, one (J1-12; SEQ ID NO:30) was found to be identical to the previously identified human pulmonary surfactant-associated protein, and the other (K1-48; SEQ ID NO:33) was determined to have some homology to *R. norvegicus* mRNA for 2-arylpropionyl-CoA epimerase. Of the 17 less abundant cDNA clones isolated from the subtracted prostate tumor specific cDNA library with spike, four (J1-16, K1-55, L1-6 and N1-1864; SEQ ID Nos:31, 34, 36 and 40, respectively) were found to be identical to previously identified sequences, two (J1-21 and N1-1860; SEQ ID Nos: 32 and 38, respectively) were found to show some homology to non-human sequences, and two (L1-2 and N1-1861; SEQ ID Nos: 35 and 39, respectively) were found to show some homology to known human sequences. No significant homologies were found to the polypeptides J1-13, J1-19, J1-24, J1-25, K1-58, K1-63, L1-4, L1-14 (SEQ ID Nos: 14-15, 16-17, 20-21, 18-19, 22-23, 24-25, 26-27, 28-29, respectively).

Subsequent studies led to the isolation of full length cDNA sequences for J1-17, L1-12 and N1-1862 (SEQ ID NOS: 109-111, respectively). The corresponding predicted amino acid sequences are provided in SEQ ID NOS: 112-114.

In a further experiment, four additional clones were identified by subtracting a prostate tumor cDNA library with normal prostate cDNA prepared from a pool of three normal prostate poly A+ RNA (prostate subtraction 2). The determined cDNA sequences for these clones, hereinafter referred to as U1-3064, U1-3065, V1-

3692 and 1A-3905, are provided in SEQ ID NO: 69-72, respectively. Comparison of the determined sequences with those in the gene bank revealed no significant homologies to U1-3065.

A second subtraction with spike (prostate subtraction spike 2) was performed by subtracting a prostate tumor specific cDNA library with spike with normal pancreas cDNA library and further spiked with PSA, J1-17, pulmonary surfactant-associated protein, mitochondrial DNA, cytochrome c oxidase subunit II, N1-1862, autonomously replicating sequence, L1-12 and tumor expression enhanced gene. Four additional clones, hereinafter referred to as V1-3686, R1-2330, 1B-3976 and V1-3679, were isolated. The determined cDNA sequences for these clones are provided in SEQ ID NO:73-76, respectively. Comparison of these sequences with those in the gene bank revealed no significant homologies to V1-3686 and R1-2330.

Further analysis of the three prostate subtractions described above (prostate subtraction 2, subtracted prostate tumor specific cDNA library with spike, and prostate subtraction spike 2) resulted in the identification of sixteen additional clones, referred to as 1G-4736, 1G-4738, 1G-4741, 1G-4744, 1G-4734, 1H-4774, 1H-4781, 1H-4785, 1H-4787, 1H-4796, 1I-4810, 1I-4811, 1J-4876, 1K-4884 and 1K-4896. The determined cDNA sequences for these clones are provided in SEQ ID NOS: 77-92, respectively. Comparison of these sequences with those in the gene bank as described above, revealed no significant homologies to 1G-4741, 1G-4734, 1I-4807, 1J-4876 and 1K-4896 (SEQ ID NOS: 79, 81, 87, 90 and 92, respectively). Further analysis of the isolated clones led to the determination of extended cDNA sequences for 1G-4736, 1G-4738, 1G-4741, 1G-4744, 1H-4774, 1H-4781, 1H-4785, 1H-4787, 1H-4796, 1I-4807, 1J-4876, 1K-4884 and 1K-4896, provided in SEQ ID NOS: 179-188 and 191-193, respectively, and to the determination of additional partial cDNA sequences for 1I-4810 and 1I-4811, provided in SEQ ID NOS: 189 and 190, respectively.

An additional subtraction was performed by subtracting a normal prostate cDNA library with normal pancreas cDNA (prostate subtraction 3). This led to the identification of six additional clones referred to as 1G-4761, 1G-4762, 1H-4766, 1H-4770, 1H-4771 and 1H-4772 (SEQ ID NOS: 93-98). Comparison of these sequences with those in the gene bank revealed no significant homologies to 1G-4761 and 1H-4771 (SEQ ID NOS: 93 and 97, respectively). Further analysis of the isolated

clones led to the determination of extended cDNA sequences for 1G-4761, 1G-4762, 1H-4766 and 1H-4772 provided in SEQ ID NOS: 194-196 and 199, respectively, and to the determination of additional partial cDNA sequences for 1H-4770 and 1H-4771, provided in SEQ ID NOS: 197 and 198, respectively.

5 Subtraction of a prostate tumor cDNA library, prepared from a pool of polyA+ RNA from three prostate cancer patients, with a normal pancreas cDNA library (prostate subtraction 4) led to the identification of eight clones, referred to as 1D-4297, 1D-4309, 1D-4278, 1D-4288, 1D-4283, 1D-4304, 1D-4296 and 1D-4280 (SEQ ID NOS: 99-107). These sequences were compared to those in the gene bank as described
10 above. No significant homologies were found to 1D-4283 and 1D-4304 (SEQ ID NOS: 103 and 104, respectively). Further analysis of the isolated clones led to the determination of extended cDNA sequences for 1D-4309, 1D-4278, 1D-4288, 1D-4283, 1D-4304, 1D-4296 and 1D-4280, provided in SEQ ID NOS: 200-206, respectively.

15 cDNA clones isolated in prostate subtraction 1 and prostate subtraction 2, described above, were colony PCR amplified and their mRNA expression levels in prostate tumor, normal prostate and in various other normal tissues were determined using microarray technology (Synteni, Palo Alto, CA). Briefly, the PCR amplification products were dotted onto slides in an array format, with each product occupying a
20 unique location in the array. mRNA was extracted from the tissue sample to be tested, reverse transcribed, and fluorescent-labeled cDNA probes were generated. The microarrays were probed with the labeled cDNA probes, the slides scanned and fluorescence intensity was measured. This intensity correlates with the hybridization intensity. Two novel clones (referred to as P509S and P510S) were found to be over-
25 expressed in prostate tumor and normal prostate and expressed at low levels in all other normal tissues tested (liver, pancreas, skin, bone marrow, brain, breast, adrenal gland, bladder, testes, salivary gland, large intestine, kidney, ovary, lung, spinal cord, skeletal muscle and colon). The determined cDNA sequences for P509S and P510S are provided in SEQ ID NO: 223 and 224, respectively. Comparison of these sequences
30 with those in the gene bank as described above, revealed some homology to previously identified ESTs.

EXAMPLE 2

DETERMINATION OF TISSUE SPECIFICITY OF PROSTATE TUMOR POLYPEPTIDES

5

Using gene specific primers, mRNA expression levels for the representative prostate tumor polypeptides F1-16, H1-1, J1-17, L1-12, F1-12 and N1-1862 were examined in a variety of normal and tumor tissues using RT-PCR.

Briefly, total RNA was extracted from a variety of normal and tumor
 10 tissues using Trizol reagent as described above. First strand synthesis was carried out using 1-2 μ g of total RNA with SuperScript II reverse transcriptase (BRL Life Technologies) at 42 °C for one hour. The cDNA was then amplified by PCR with gene-specific primers. To ensure the semi-quantitative nature of the RT-PCR, β -actin was used as an internal control for each of the tissues examined. First, serial dilutions of the
 15 first strand cDNAs were prepared and RT-PCR assays were performed using β -actin specific primers. A dilution was then chosen that enabled the linear range amplification of the β -actin template and which was sensitive enough to reflect the differences in the initial copy numbers. Using these conditions, the β -actin levels were determined for each reverse transcription reaction from each tissue. DNA contamination was
 20 minimized by DNase treatment and by assuring a negative PCR result when using first strand cDNA that was prepared without adding reverse transcriptase.

mRNA Expression levels were examined in four different types of tumor tissue (prostate tumor from 2 patients, breast tumor from 3 patients, colon tumor, lung tumor), and sixteen different normal tissues, including prostate, colon, kidney, liver,
 25 lung, ovary, pancreas, skeletal muscle, skin, stomach, testes, bone marrow and brain. F1-16 was found to be expressed at high levels in prostate tumor tissue, colon tumor and normal prostate, and at lower levels in normal liver, skin and testes, with expression being undetectable in the other tissues examined. H1-1 was found to be expressed at high levels in prostate tumor, lung tumor, breast tumor, normal prostate, normal colon
 30 and normal brain, at much lower levels in normal lung, pancreas, skeletal muscle, skin, small intestine, bone marrow, and was not detected in the other tissues tested. J1-17

and L1-12 appear to be specifically over-expressed in prostate, with both genes being expressed at high levels in prostate tumor and normal prostate but at low to undetectable levels in all the other tissues examined. N1-1862 was found to be over-expressed in 60% of prostate tumors and detectable in normal colon and kidney. The RT-PCR results thus indicate that F1-16, H1-1, J1-17, N1-1862 and L1-12 are either prostate specific or are expressed at significantly elevated levels in prostate.

Further RT-PCR studies showed that F1-12 is over-expressed in 60% of prostate tumors, detectable in normal kidney but not detectable in all other tissues tested. Similarly, R1-2330 was shown to be over-expressed in 40% of prostate tumors, detectable in normal kidney and liver, but not detectable in all other tissues tested. U1-3064 was found to be over-expressed in 60% of prostate tumors, and also expressed in breast and colon tumors, but was not detectable in normal tissues.

RT-PCR characterization of R1-2330, U1-3064 and 1D-4279 showed that these three antigens are over-expressed in prostate and/or prostate tumors.

Northern analysis with four prostate tumors, two normal prostate samples, two BPH prostates, and normal colon, kidney, liver, lung, pancreas, skeletal muscle, brain, stomach, testes, small intestine and bone marrow, showed that L1-12 is over-expressed in prostate tumors and normal prostate, while being undetectable in other normal tissues tested. J1-17 was detected in two prostate tumors and not in the other tissues tested. N1-1862 was found to be over-expressed in three prostate tumors and to be expressed in normal prostate, colon and kidney, but not in other tissues tested. F1-12 was found to be highly expressed in two prostate tumors and to be undetectable in all other tissues tested.

The micro-array technology described above was used to determine the expression levels of representative antigens described herein in prostate tumor, breast tumor and the following normal tissues: prostate, liver, pancreas, skin, bone marrow, brain, breast, adrenal gland, bladder, testes, salivary gland, large intestine, kidney, ovary, lung, spinal cord, skeletal muscle and colon. L1-12 was found to be over-expressed in normal prostate and prostate tumor, with some expression being detected in normal skeletal muscle. Both J1-12 and F1-12 were found to be over-expressed in prostate tumor, with expression being lower or undetectable in all other tissues tested. N1-1862 was found to be expressed at high levels in prostate tumor and normal

prostate, and at low levels in normal large intestine and normal colon, with expression being undetectable in all other tissues tested. R1-2330 was found to be over-expressed in prostate tumor and normal prostate, and to be expressed at lower levels in all other tissues tested. 1D-4279 was found to be over-expressed in prostate tumor and normal prostate, expressed at lower levels in normal spinal cord, and to be undetectable in all other tissues tested.

Example 3

ISOLATION AND CHARACTERIZATION OF PROSTATE TUMOR POLYPEPTIDES BY PCR-BASED SUBTRACTON

A cDNA subtraction library, containing cDNA from normal prostate subtracted with ten other normal tissue cDNAs (brain, heart, kidney, liver, lung, ovary, placenta, skeletal muscle, spleen and thymus) and then submitted to a first round of PCR amplification, was purchased from Clontech. This library was subjected to a second round of PCR amplification, following the manufacturer's protocol. The resulting cDNA fragments were subcloned into the vector pT7 Blue T-vector (Novagen, Madison, WI) and transformed into XL-1 Blue MRF' *E. coli* (Stratagene). DNA was isolated from independent clones and sequenced using a Perkin Elmer/Applied Biosystems Division Automated Sequencer Model 373A.

Fifty-nine positive clones were sequenced. Comparison of the DNA sequences of these clones with those in the gene bank, as described above, revealed no significant homologies to 25 of these clones, hereinafter referred to as P5, P8, P9, P18, P20, P30, P34, P36, P38, P39, P42, P49, P50, P53, P55, P60, P64, P65, P73, P75, P76, P79, and P84. The determined cDNA sequences for these clones are provided in SEQ ID NO:41-45, 47-52 and 54-65, respectively. P29, P47, P68, P80 and P82 (SEQ ID NO:46, 53 and 66-68, respectively) were found to show some degree of homology to previously identified DNA sequences. To the best of the inventors' knowledge, none of these sequences have been previously shown to be present in prostate.

Further studies using the PCR-based methodology described above resulted in the isolation of more than 180 additional clones, of which 23 clones were

found to show no significant homologies to known sequences. The determined cDNA sequences for these clones are provided in SEQ ID NO: 115-123, 127, 131, 137, 145, 147-151, 153, 156-158 and 160. Twenty-three clones (SEQ ID NO: 124-126, 128-130, 132-136, 138-144, 146, 152, 154, 155 and 159) were found to show some homology to previously identified ESTs. An additional ten clones (SEQ ID NO: 161-170) were found to have some degree of homology to known genes. An additional clone, referred to as P703, was found to have five splice variants. The determined DNA sequence for the variants referred to as DE1, DE13 and DE14 are provided in SEQ ID NOS: 171, 175 and 177, respectively, with the corresponding predicted amino acid sequences being provided in SEQ ID NO: 172, 176 and 178, respectively. The DNA sequences for the splice variants referred to as DE2 and DE6 are provided in SEQ ID NOS: 173 and 174, respectively.

mRNA Expression levels for representative clones in tumor tissues (prostate (n=5), breast (n=2), colon and lung) normal tissues (prostate (n=5), colon, kidney, liver, lung (n=2), ovary (n=2), skeletal muscle, skin, stomach, small intestine and brain), and activated and non-activated PBMC was determined by RT-PCT as described above. Expression was examined in one sample of each tissue type unless otherwise indicated.

P9 was found to be highly expressed in normal prostate and prostate tumor compared to all normal tissues tested except for normal colon which showed comparable expression. P20 was found to be highly expressed in normal prostate and prostate tumor, compared to all twelve normal tissues tested. A modest increase in expression of P20 in breast tumor (n=2), colon tumor and lung tumor was seen compared to all normal tissues except lung (1 of 2). Increased expression of P18 was found in normal prostate, prostate tumor and breast tumor compared to other normal tissues except lung and stomach. A modest increase in expression of P5 was observed in normal prostate compared to most other normal tissues. However, some elevated expression was seen in normal lung and PBMC. Elevated expression of P5 was also observed in prostate tumors (2 of 5), breast tumor and one lung tumor sample. For P30, similar expression levels were seen in normal prostate and prostate tumor, compared to six of twelve other normal tissues tested. Increased expression was seen in breast tumors, one lung tumor sample and one colon tumor sample, and also in normal

PBMC. P29 was found to be over-expressed in prostate tumor (5 of 5) and normal prostate (5 of 5) compared to the majority of normal tissues. However, substantial expression of P29 was observed in normal colon and normal lung (2 of 2). P80 was found to be over-expressed in prostate tumor (5 of 5) and normal prostate (5 of 5) compared to all other normal tissues tested, with increased expression also being seen in colon tumor.

Further studies using the above methodology resulted in the isolation of sixteen additional clones. The determined DNA sequences for these clones are provided in SEQ ID NO: 207-222.

EXAMPLE 4

SYNTHESIS OF POLYPEPTIDES

Polypeptides may be synthesized on an Applied Biosystems 430A peptide synthesizer using Fmoc chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation, binding to an immobilized surface, or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using the following cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0%-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to elute the peptides. Following lyophilization of the pure fractions, the peptides may be characterized using electrospray or other types of mass spectrometry and by amino acid analysis.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANTS: Xu, Jiangchun
Dillon, Davin C.

(ii) TITLE OF INVENTION: COMPOUNDS FOR IMMUNODIAGNOSIS OF PROSTATE CANCER
AND METHODS FOR THEIR USE

(iii) NUMBER OF SEQUENCES: 224

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: SEED and BERRY LLP
(B) STREET: 6300 Columbia Center, 701 Fifth Avenue
(C) CITY: Seattle
(D) STATE: WA
(E) COUNTRY: USA
(F) ZIP: 98104

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE: 23-FEB-1998
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Maki, David J.
(B) REGISTRATION NUMBER: 31,392
(C) REFERENCE/DOCKET NUMBER: 210121.428C3

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (206) 622-4900
(B) TELEFAX: (206) 682-6031

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 814 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TTTTTTTTTT	TTTTTCACAG	TATAACAGCT	CTTTATTTCT	GTGAGTTCTA	CTAGGAAATC	60
ATCAAATCTG	AGGGTTGTCT	GGAGGACTTC	AATACACCTC	CCCCCATAGT	GAATCAGCTT	120
CCAGGGGGTC	CAGTCCCTCT	CCTTACTTCA	TCCCCATCCC	ATGCCAAAGG	AAGACCCTCC	180
CTCCTTGGCT	CACAGCCTTC	TCTAGGCTTC	CCAGTGCCTC	CAGGACAGAG	TGGGTTATGT	240

265220 90902050

TTTCAGCTCC	ATCCTTGCTG	TGAGTGTCTG	GTGCGTTGTG	CCTCCAGCTT	CTGCTCAGTG	300
CTTCATGGAC	AGTGTCCAGC	ACATGTCACT	CTCCACTCTC	TCAGTGTGGA	TCCACTAGTT	360
CTAGAGCGGC	CGCCACCGCG	GTGGAGCTCC	AGCTTTTGTG	CCCTTTAGTG	AGGGTTAATT	420
GCGCGCTTGG	CGTAATCATG	GTCATAACTG	TTTCCTGTGT	GAAATTGTTA	TCCGCTCACA	480
ATTCCACACA	ACATACGAGC	CGGAAGCATA	AAGTGTAAG	CCTGGGGTGC	CTAATGAGTG	540
ANCTAACTCA	CATTAATTGC	GTTGCGCTCA	CTGNCCGCTT	TCCAGTCNGG	AAAACTGTCTG	600
TGCCAGCTGC	ATTAATGAAT	CGGCCAACGC	NCGGGGAAAA	GCGGTTTGGC	TTTTGGGGGC	660
TCTTCCGCTT	CTCGCTCACT	NANTCCTGCG	CTCGGTCNTT	CGGCTGCGGG	GAACGGTATC	720
ACTCCTCAAA	GGNGGTATTA	CGGTTATCCN	NAAATCNGGG	GATACCCNGG	AAAAAANTTT	780
AACAAAAGGG	CANCAAAGGG	CNGAAACGTA	AAAA			814

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 816 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ACAGAAATGT	TGGATGGTGG	AGCACCTTTC	TATACGACTT	ACAGGACAGC	AGATGGGGAA	60
TTCATGGCTG	TTGGAGCAAT	AGAACCCAG	TTCTACGAGC	TGCTGATCAA	AGGACTTGGA	120
CTAAAGTCTG	ATGAACCTCC	CAATCAGATG	AGCATGGATG	ATTGGCCAGA	AATGAAGAAG	180
AAGTTTGCTG	ATGTATTTGC	AAAGAAGACG	AAGGCAGAGT	GGTGTCAAAT	CTTTGACGGC	240
ACAGATGCCT	GTGTGACTCC	GGTTCTGACT	TTTGAGGAGG	TTGTTTCATCA	TGATCACAAC	300
AAGGAACGGG	GCTCGTTTAT	CACCACTGAG	GAGCAGGACG	TGAGCCCCCG	CCCTGCACCT	360
CTGCTGTTAA	ACACCCACAG	CATCCCTTCT	TTCAAAAGGG	ATCCACTAGT	TCTAGAAGCG	420
GCCGCCACCG	CGGTGGAGCT	CCAGCTTTTG	TTCCCTTTAG	TGAGGGTTAA	TTGCGCGCTT	480
GGCGTAATCA	TGGTCATAGC	TGTTTCCTGT	GTGAAATTGT	TATCCGCTCA	CAATTCCCCC	540
AACATACGAG	CCGGAACATA	AAGTGTTAAG	CCTGGGGTGC	CTAATGANTG	AGCTAACTCN	600
CATTAATTGC	GTTGCGCTCA	CTGCCCCTT	TCCAGTCGGG	AAAACTGTCTG	TGCCACTGCN	660
TTANTGAATC	NGCCACCCCC	CGGGAAAAGG	CGGTTGCNTT	TTGGGCCTCT	TCCGCTTTCC	720
TCGCTCATTG	ATCCTNGCNC	CCGGTCTTCG	GCTGCGGNGA	ACGGTTCACT	CCTCAAAGGC	780
GGTNTNCCCG	TTATCCCCAA	ACNGGGGATA	CCNGA			816

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 773 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CTTTTGAAAG	AAGGGATGGC	TGGGGTGTTT	AACAGCAGAG	GTGCAGGGCG	GGGGCTCACG	60
TCCTGCTCCT	CACTGGTGAT	AAACGAGCCC	CGTTCCTTGT	TGTGATCATG	ATGAACAACC	120
TCCTCAAAAG	TCAGAACCGG	AGTCACACAG	GCATCTGTGC	CGTCAAAGAT	TTGACACCAC	180
TCTGCCTTCG	TCTTCTTTGC	AAATACATCT	GCAAACCTTCT	TCTTCATTTT	TGGCCAATCA	240
TCCATGCTCA	TCTGATTGGG	AAGTTCATCA	GACTTTAGTC	CANNTCCTTT	GATCAGCAGC	300
TCGTAGAACT	GGGGTTCTAT	TGCTCCAACA	GCCATGAATT	CCCCATCTGC	TGTCCTGTAA	360
GTCGTATAGA	AAGGTGCTCC	ACCATCCAAC	ATGTTCTGTC	CTCGAGGGGG	GGCCCGGTAC	420
CCAATTCGCC	CTATANTGAG	TCGTATTACG	CGCGCTCACT	GGCCGTCGTT	TTACAACGTC	480

GTGACTGGGA	AAACCCTGGG	CGTTACCAAC	TTAATCGCCT	TGCAGCACAT	CCCCCTTTCG	540
CCAGCTGGGC	GTAATANCGA	AAAGGCCCGC	ACCGATCGCC	CTTCCAACAG	TTGCGCACCT	600
GAATGGGNAA	ATGGGACCCC	CCTGTTACCG	CGCATTNAAC	CCCCGCNGGG	TTTNGTTGTT	660
ACCCCCACNT	NNACCGCTTA	CACTTTGCCA	GCGCCTTANC	GCCCGCTCCC	TTTCNCCTTT	720
CTTCCCTTCC	TTTCNCNCCN	CTTTCCCCCG	GGGTTTCCCC	CNTCAAACCC	CNA	773

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 828 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCTCCTGAGT	CCTACTGACC	TGTGCTTTCT	GGTGTGGAGT	CCAGGGCTGC	TAGGAAAAGG	60
AATGGGCAGA	CACAGGTGTA	TGCCAATGTT	TCTGAAATGG	GTATAATTTT	GTCCTCTCCT	120
TCGGAACACT	GGCTGTCTCT	GAAGACTTCT	CGCTCAGTTT	CAGTGAGGAC	ACACACAAAG	180
ACGTGGGTGA	CCATGTTGTT	TGTGGGGTGC	AGAGATGGGA	GGGGTGGGGC	CCACCCTGGA	240
AGAGTGGACA	GTGACACAAG	GTGGACACTC	TCTACAGATC	ACTGAGGATA	AGCTGGAGCC	300
ACAATGCATG	AGGCACACAC	ACAGCAAGGA	TGACNCTGTA	AACATAGCCC	ACGCTGTCCT	360
GNGGGCACTG	GGAAGCCTAN	ATNAGGCCGT	GAGCANAAAG	AAGGGGAGGA	TCCACTAGTT	420
CTANAGCGGC	CGCCACCGCG	GTGGANCTCC	ANCTTTTGTT	CCCTTTAGTG	AGGGTTAATT	480
GCGCGCTTGG	CNTAATCATG	GTCATANCTN	TTTCCTGTGT	GAAATTGTTA	TCCGCTCACA	540
ATTCCACACA	ACATACGANC	CGGAAACATA	AANTGTAAAC	CTGGGGTGCC	TAATGANTGA	600
CTAACTCACA	TTAATTGCGT	TGCGCTCACT	GCCCCTTTT	CAATCNGGAA	ACCTGTCTTG	660
CCNCTTGCA	TNATGAATCN	GCCAACCCCC	GGGGAAAAGC	GTTTGCGTTT	TGGGCGCTCT	720
TCCGCTTCCT	CNCTCANTTA	NTCCCTNCNC	TCGGTCATTC	CGGCTGCNGC	AAACCGGTTT	780
ACCNCCTCCA	AAGGGGGTAT	TCCGGTTTCC	CCNAATCCGG	GGANANCC		828

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 834 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TTTTTTTTTT	TTTTTACTGA	TAGATGGAAT	TTATTAAGCT	TTTCACATGT	GATAGCACAT	60
AGTTTTTAAT	GCATCCAAAG	TACTAACAAA	AACTCTAGCA	ATCAAGAATG	GCAGCATGTT	120
ATTTTATAAC	AATCAACACC	TGTGGCTTTT	AAAATTTGGT	TTTCATAAGA	TAATTTTATAC	180
TGAAGTAAAT	CTAGCCATGC	TTTTAAAAAA	TGCTTTAGGT	CACTCCAAGC	TTGGCAGTTA	240
ACATTTGGCA	TAAACAATAA	TAAAACAATC	ACAATTTAAT	AAATAACAAA	TACAACATTG	300
TAGGCCATAA	TCATATACAG	TATAAGGAAA	AGGTGGTAGT	GTTGAGTAAG	CAGTTATTAG	360
AATAGAATAC	CTTGGCCTCT	ATGCAAATAT	GTCTAGACAC	TTTGATTTCAC	TCAGCCCTGA	420
CATTCAGTTT	TCAAAGTAGG	AGACAGGTTT	TACAGTATCA	TTTTACAGTT	TCCAACACAT	480
TGAAAACAAG	TAGAAAATGA	TGAGTTGATT	TTTATTTAAT	CATTACATCC	TCAAGAGTTA	540
TCACCAACCC	CTCAGTTATA	AAAAATTTTC	AAGTTATATT	AGTCATATAA	CTTGGTGTGC	600
TTATTTTAAA	TTAGTGCTAA	ATGGATTAA	TGAAGACAAC	AATGGTCCCC	TAATGTGATT	660
GATATTGGTC	ATTTTACCA	GCTTCTAAAT	CTNAACTTTC	AGGCTTTTGA	ACTGGAACAT	720
TGNATNACAG	TGTTCCANAG	TTNCAACCTA	CTGGAACATT	ACAGTGTGCT	TGATTCAAAA	780

TGTTATTTTGG TTAAAAATTA AATTTTAACC TGGTGGAAAA ATAATTTGAA ATNA

834

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 818 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TTTTTTTTTTT	TTTTTTTTTTT	AAGACCCTCA	TCAATAGATG	GAGACATACA	GAAATAGTCA	60
AACCACATCT	ACAAAATGCC	AGTATCAGGC	GGCGGCTTCG	AAGCCAAAGT	GATGTTTGGA	120
TGTAAAGTGA	AATATTAGTT	GGCGGATGAA	GCAGATAGTG	AGGAAAGTTG	AGCCAATAAT	180
GACGTGAAGT	CCGTGGAAGC	CTGTGGCTAC	AAAAAATGTT	GAGCCGTAGA	TGCCGTCGGA	240
AATGGTGAAG	GGAGACTCGA	AGTACTCTGA	GGCTTG TAGG	AGGGTAAAAT	AGAGACCCAG	300
TAAAATTGTA	ATAAGCAGTG	CTTGAATTAT	TTGGTTTCGG	TTGTTTTCTA	TTAGACTATG	360
GTGAGCTCAG	GTGATTGATA	CTCCTGATGC	GAGTAATACG	GATGTGTTTA	GGAGTGGGAC	420
TTCTAGGGGA	TTTAGCGGGG	TGATGCCTGT	TGGGGGCCAG	TGCCCTCCTA	GTTGGGGGGT	480
AGGGGCTAGG	CTGGAGTGGT	AAAAGGCTCA	GAAAAATCCT	GCGAAGAAAA	AAACTTCTGA	540
GGTAATAAAT	AGGATTATCC	CGTATCGAAG	GCCTTTTTGG	ACAGGTGGTG	TGTGGTGGCC	600
TTGGTATGTG	CTTTCTCGTG	TTACATCGCG	CCATCATTGG	TATATGGTTA	GTGTGTTGGG	660
TTANTANGGC	CTANTATGAA	GAACTTTTGG	ANTGGAATTA	AATCAATNGC	TTGGCCGGAA	720
GTCATTANGA	NGGCTNAAAA	GGCCCTGTTA	NGGGTCTGGG	CTNGGTTTTA	CCCNACCCAT	780
GGAATNCNCC	CCCCGGACNA	NTGNATCCCT	ATTCTTAA			818

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 817 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TTTTTTTTTTT	TTTTTTTTTTT	TGGCTCTAGA	GGGGGTAGAG	GGGGTGCTAT	AGGGTAAATA	60
CGGGCCCTAT	TTCAAAGATT	TTTAGGGGAA	TTAATTC TAG	GACGATGGGT	ATGAAACTGT	120
GGTTTGCTCC	ACAGATTTC A	GAGCATTGAC	CGTAGTATAC	CCCCGGTCGT	GTAGCGGTGA	180
AAGTG GTTTG	GTTTAGACGT	CCGGGAATTG	CATCTGTTTT	TAAGCCTAAT	GTGGGGACAG	240
CTCATGAGTG	CAAGACGTCT	TGTGATGTAA	TTATTATACN	AATGGGGGCT	TCAATCGGGA	300
GTACTACTCG	ATTGTCAACG	TCAAGGAGTC	GCAGGTCGCC	TGGTTC TAGG	AATAATGGGG	360
GAAGTATGTA	GGAATTGAAG	ATTAATCCGC	CGTAGTCGGT	GTTCCTCCTAG	GTTCAATACC	420
ATTGGTGGCC	AATTGATTTG	ATGGTAAGGG	GAGGGATCGT	TGAAC TCGTC	TGTTATGTAA	480
AGGATNCCTT	NGGGATGGGA	AGGCNATNAA	GGACTANGGA	TNAATGGCGG	GCANGATATT	540
TCAAACNGTC	TCTANTTCCT	GAAACGTCTG	AAATGTTAAT	AANAATT AAN	TTTNGTTATT	600
GAATNTTNNG	GAAAAGGGCT	TACAGGACTA	GAAACCAAAT	ANGAAAAANTA	ATNNTAANGG	660
CNTTATCNTN	AAAGGTNATA	ACCNCTCCTA	TNATCCCACC	CAATNGNATT	CCCCACNCNN	720
ACNATTGGAT	NCCCCANTTC	CANAAANGGC	CNCCCCCGG	TGNANNCCNC	CTTTTGTTCC	780
CTTNANTGAN	GGTTATTCNC	CCCTNGCNTT	ATCANCC			817

(2) INFORMATION FOR SEQ ID NO:8:

- (A) LENGTH: 799 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CATTTCCGGG	TTTACTTTCT	AAGGAAAGCC	GAGCGGAAGC	TGCTAACGTG	GGAATCGGTG	60
CATAAGGAGA	ACTTTCTGCT	GGCACGCGCT	AGGGACAAGC	GGGAGAGCGA	CTCCGAGCGT	120
CTGAAGCGCA	CGTCCCAGAA	GGTGGACTTG	GCACTGAAAC	AGCTGGGACA	CATCCGCGAG	180
TACGAACAGC	GCCTGAAAGT	GCTGGAGCGG	GAGGTCCAGC	AGTGTAGCCG	CGTCCTGGGG	240
TGGGTGGCCG	ANGCCTGANC	CGCTCTGCCT	TGCTGCCCCC	ANGTGGGCCG	CCACCCCCTG	300
ACCTGCCTGG	GTCCAAACAC	TGAGCCCTGC	TGGCGGACTT	CAAGGANAAC	CCCCACANGG	360
GGATTTTGCT	CCTANANTAA	GGCTCATCTG	GGCCTCGGCC	CCCCACCTG	GTTGGCCTTG	420
TCTTTGANGT	GAGCCCCATG	TCCATCTGGG	CCACTGTCTG	GACCACCTTT	NGGGAGTGTT	480
CTCCTTACAA	CCACANNATG	CCCGGCTCCT	CCCGGAAACC	ANTCCCANCC	TGNGAAGGAT	540
CAAGNCCTGN	ATCCACTTNN	NCTANAACCG	GCCNCCNCCG	CNGTGGAACC	CNCCTTNTGT	600
TCCTTTTCNT	TNAGGGTTAA	TNNCGCCCTG	GCCTTNCCAN	NGTCC'TNCNC	NTTTTCCNNT	660
GT'TNAAATTG	TTANGCNCCAT	NCCNNTCCCN	CNNC'NNCNAN	CCC'GACCCNN	ANN'TTNNANN	720
NCC'TGGGGTG	NCCN'NCGAT	TGACCC'NNCC	NCCCTNTANT	TGCNT'TNGGG	NNC'NNTGCCC	780
CTTTCCTCT	TGGGANNCG					799

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 801 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ACGCCCTTGAT	CCTCCCAGGC	TGGGACTGGT	TCTGGGAGGA	GCCGGGCATG	CTGTGGTTTG	60
TAANGATGAC	ACTCCCAAAG	GTGGTCCTGA	CAGTGGCCCA	GATGGACATG	GGGCTCACCT	120
CAAGGACAAG	GCCACCAGGT	GCGGGGGCCG	AAGCCCACAT	GATCCTTACT	CTATGAGCAA	180
AATCCCC'TGT	GGGGGC'TTCT	CC'TTGAAGTC	CGCCANCAGG	GCTCAGTCTT	TGGACCCANG	240
CAGGTCATGG	GGT'TGTNGNC	CAACTGGGGG	CCNCAACGCA	AAANGGCNCA	TGGCCTCNGN	300
CACCCATCCC	ANGACATCGGC	TACACTNCTG	GACCTCCCN	TCCACCACCT	GTTCGCGCTG	360
TTCNTACCCG	CGNATNTGTC	CCANCTGTTT	CNGTGCCNAC	TCCANCTTCT	NGGACGTGCG	420
CTACATACGC	CCGGANTCNC	NCTCCCGCTT	TGTCCCTATC	CACGTNCCAN	CAACAAATTT	480
CNCCNTANTG	CACCNAT'TCC	CACNT'TTNNC	AGNT'TTCCNC	NNCGNGCTTC	CTTNTAAAAG	540
GGTTGANCCC	CGGAAAATNC	CCCAAAGGGG	GGGGGCCNGG	TACCCAAC'TN	CCCCCTNATA	600
GCTGAANTCC	CCATNACCNN	GNCTCNATGG	ANCCNTCCNT	T'TTAANNACN	T'TCTNAACTT	660
GGGAANANCC	CTCGNCCNTN	CCCCCNTTAA	TCCNCCTTGG	CNANGNNCNT	CCCCCNNTCC	720
NCCNNNTNG	GCNTNTNANN	CNAAAAAGGC	CCNNNANCAA	TCTCCTNNCN	CCTCANTTCG	780
CCANCCCTCG	AAATCGGCCN	C				801

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 789 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CAGTCTATNT	GGCCAGTGTG	GCAGCTTTCC	CTGTGGCTGC	CGGTGCCACA	TGCCTGTCCC	60
ACAGTGTGGC	CGTGGTGACA	GCTTCAGCCG	CCCTCACCGG	GTTACACCTT	TCAGCCCTGC	120
AGATCCTGCC	CTACACACTG	GCCTCCCTCT	ACCACCGGGA	GAAGCAGGTG	TTCCTGCCCA	180
AATACCGAGG	GGACACTGGA	GGTGCTAGCA	GTGAGGACAG	CCTGATGACC	AGCTTCCTGC	240
CAGGCCCTAA	GCCTGGAGCT	CCCTTCCCTA	ATGGACACGT	GGGTGCTGGA	GGCAGTGGCC	300
TGCTCCCACC	TCCACCCGCG	CTCTGCGGGG	CCTCTGCCTG	TGATGTCTCC	GTACGTGTGG	360
TGGTGGGTGA	GCCCACCGAN	GCCAGGGTGG	TTCCGGGCCG	GGGCATCTGC	CTGGACCTCG	420
CCATCCTGGA	TAGTGCTTCC	TGCTGTCCCA	NGTGGCCCCA	TCCCTGTTTA	TGGGCTCCAT	480
TGTCCAGCTC	AGCCAGTCTG	TCAGTGCCTA	TATGGTGTCT	GCCGCAGGCC	TGGGTCTGGT	540
CCCATTTACT	TTGCTACACA	GGTANTATTT	GACAAGAACG	ANTTGGCCAA	ATACTCAGCG	600
TTAAAAAATT	CCAGCAACAT	TGGGGGTGGA	AGGCCTGCCT	CACTGGGTCC	AACTCCCCGC	660
TCCTGTTAAC	CCCATGGGGC	TGCCGGCTTG	GCCGCCAATT	TCTGTTGCTG	CCAAANTNAT	720
GTGGCTCTCT	GCTGCCACCT	GTTGCTGGCT	GAAGTGCNTA	CNGCNCANCT	NGGGGGGTNG	780
GGNGTTCCC						789

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 772 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CCCACCCTAC	CCAAATATTA	GACACCAACA	CAGAAAAGCT	AGCAATGGAT	TCCCTTCTAC	60
TTTGTAAAT	AAATAAGTTA	AATATTTAAA	TGCCTGTGTC	TCTGTGATGG	CAACAGAAGG	120
ACCAACAGGC	CACATCCTGA	TAAAAGGTAA	GAGGGGGGTG	GATCAGCAAA	AAGACAGTGC	180
TGTGGGCTGA	GGGGACCTGG	TTCTTGTGTG	TTGCCCCTCA	GGACTCTTCC	CCTACAAATA	240
ACTTTTCATAT	GTTCAAATCC	CATGGAGGAG	TGTTTCATCC	TAGAAACTCC	CATGCAAGAG	300
CTACATTAAA	CGAAGCTGCA	GGTTAAGGGG	CTTANAGATG	GGAAACCAGG	TGACTGAGTT	360
TATTCAGCTC	CCAAAAACCC	TTCTCTAGGT	GTGTCTCAAC	TAGGAGGCTA	GCTGTTAACC	420
CTGAGCCTGG	GTAATCCACC	TGCAGAGTCC	CCGCATTCCA	GTGCATGGAA	CCCTTCTGGC	480
CTCCCTGTAT	AAGTCCAGAC	TGAAACCCCC	TTGGAAGGNC	TCCAGTCAGG	CAGCCCTANA	540
AACTGGGGAA	AAAAGAAAAG	GACGCCCCAN	CCCCCAGCTG	TGCANCTACG	CACCTCAACA	600
GCACAGGGTG	GCAGCAAAAA	AACCACTTTA	CTTTGGCACA	AACAAAAACT	NGGGGGGGCA	660
ACCCCGGCAC	CCCNANGGGG	GTTAACAGGA	ANCNGGGNAA	CNTGGAACCC	AATTNAGGCA	720
GGCCCNCCAC	CCCNAAATNTT	GCTGGGAAAT	TTTTCTCTCC	CTAAATNTTT	TC	772

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 751 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

GCCCCAATTC CAGCTGCCAC ACCACCCACG GTGACTGCAT TAGTTCGGAT GTCATACAAA      60
AGCTGATTGA AGCAACCCTC TACTTTTTTG TCGTGAGCCT TTTGCTTGGT GCAGGTTTCA      120
TTGGCTGTGT TGGTGACGTT GTCATTGCAA CAGAATGGGG GAAAGGCACT GTTCTCTTTG      180
AAGTANGGTG AGTCCTCAAA ATCCGTATAG TTGGTGAAGC CACAGCACTT GAGCCCTTTC      240
ATGGTGGTGT TCCACACTTG AGTGAAGTCT TCCTGGGAAC CATAATCTTT CTTGATGGCA      300
GGCACTACCA GCAACGTCAG GGAAGTGCTC AGCCATTGTG GTGTACACCA AGGCGACCAC      360
AGCAGCTGCN ACCTCAGCAA TGAAGATGAN GAGGANGATG AAGAAGAACG TCNCGAGGGC      420
ACACTTGCTC TCAGTCTTAN CACCATANCA GCCCNTGAAA ACCAANANCA AAGACCACNA      480
CNCCGGCTGC GATGAAGAAA TNACCCCNCG TTGACAAACT TGCATGGCAC TGGGANCCAC      540
AGTGGCCCNNA AAAATCTTCA AAAAGGATGC CCCATCNATT GACCCCCCAA ATGCCCCTG      600
CCAACAGGGG CTGCCCCACN CNCNNAACGA TGANCCNATT GNACAAGATC TNCNTGGTCT      660
TNATNAACNT GAACCCTGCN TNGTGGCTCC TGTTTCAGGNC CNNGGCCTGA CTTCTNAANN      720
AANGAACTCN GAAGNCCCCA CNGGANANNC G                                     751

```

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 729 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

GAGCCAGGCG TCCCTCTGCC TGCCCACTCA GTGGCAACAC CCGGGAGCTG TTTTGTCTTT      60
TGTGGANCCCT CAGCAGTNCC CTCTTTCAGA ACTCANTGCC AAGANCCCTG AACAGGAGCC      120
ACCATGCAGT GCTTCAGCTT CATTAAAGACC ATGATGATCC TCTTCAATTT GCTCATCTTT      180
CTGTGTGGTG CAGCCCTGTT GGCAGTGGGC ATCTGGGTGT CAATCGATGG GGCATCCTTT      240
CTGAAGATCT TCGGGCCACT GTCGTCCAGT GCCATGCAGT TTGTCAACGT GGGCTACTTC      300
CTCATCGCAG CCGGCGTTGT GGTCTTAGCT CTAGGTTTCC TGGGCTGCTA TGGTGCTAAG      360
ACTGAGAGCA AGTGTGCCCT CGTGACGTTT TTCTTCATCC TCCTCCTCAT CTTCATTGCT      420
GAGGTTGCAA TGCTGTGGTC GCCTTGGTGT ACACCACAAT GGCTGAGCAC TTCCTGACGT      480
TGCTGGTAAT GCCTGCCATC AANAAAAGAT TATGGGTTCC CAGGAANACT TCACTCAAGT      540
GTTGGAACAC CACCATGAAA GGGCTCAAGT GCTGTGGCTT CNNCCAATA TACGGATTTT      600
GAAGANTCAC CTACTTCAAA GAAAANAGTG CCTTCCCCC ATTTCTGTTG CAATTGACAA      660
ACGTCCCCAA CACAGCCAAT TGAAAACCTG CACCCAACCC AAANGGGTCC CCAACCANAA      720
ATTNAAGGG                                     729

```

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 816 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```

TGCTCTTCCT CAAAGTTGTT CTTGTTGCCA TAACAACCAC CATAGGTAAA GCGGGCGCAG      60
TGTTTCGCTGA AGGGGTTGTA GTACCAGCGC GGGATGCTCT CCTTGCAGAG TCCTGTGTCT      120
GGCAGGTCCA CGCAGTGCCC TTTGTCACTG GGGAAATGGA TGCGCTGGAG CTCGTCAAAG      180
CCACTCGTGT ATTTTTCACA GGCAGCCTCG TCCGACGCGT CGGGGCAGTT GGGGGTGTCT      240
TCACACTCCA GGAAACTGTC NATGCAGCAG CCATTGCTGC AGCGGAACTG GGTGGGCTGA      300

```


CANGTGCCAG	AGCACACTGG	ATGGCGCCTT	TCCATGNNAN	GGGCCCTGNG	GGAAAGTCCC	360
TGANCCCCAN	ANCTGCCTCT	CAAANGCCCC	ACCTTGACAC	CCCCGACAGG	CTAGAATGGA	420
ATCTTCTTCC	CGAAAGGTAG	TTNTTCTTGT	TGCCCAANCC	ANCCCCNTAA	ACAAACTCTT	480
GCANATCTGC	TCCGNGGGGG	TCNTANTACC	ANCGTGGGAA	AAGAACCCCA	GGCNGCGAAC	540
CAANCTTGTT	TGGATNCGAA	GCNATAATCT	NCTNTTCTGC	TTGGTGGACA	GCACCANTNA	600
CTGTNNANCT	TTAGNCCNTG	GTCTCNTTGG	GTTGNNCTTG	AACCTAATCN	CCNNTCAACT	660
GGGACAAGGT	AANTNGCCNT	CCTTTNAATT	CCCNANCNTN	CCCCCTGGTT	TGGGGTTTTN	720
CNCNCTCCTA	CCCCAGAAAN	NCCGTGTTCC	CCCCAACCTA	GGGGCCNAAA	CCNNTTNTTC	780
CACAACCCTN	CCCCACCCAC	GGGTTGNGNT	GGTTCNG			816

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 783 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CCAAGGCCTG	GGCAGGCATA	NACTTGAAGG	TACAACCCCA	GGAACCCCTG	GTGCTGAAGG	60
ATGTGGA AAA	CACAGATTGG	CGCCTACTGC	GGGGTGACAC	GGATGTCAGG	GTAGAGAGGA	120
AAGACCCAAA	CCAGGTGGAA	CTGTGGGGAC	TCAAGGAANG	CACCTACCTG	TTCCAGCTGA	180
CAGTGACTAG	CTCAGACCAC	CCAGAGGACA	CGGCCAACGT	CACAGTCACT	GTGCTGTCCA	240
CCAAGCAGAC	AGAAGACTAC	TGCCTCGCAT	CCAACAANGT	GGGTCGCTGC	CGGGGCTCTT	300
TCCCACGCTG	GTACTATGAC	CCCACGGAGC	AGATCTGCAA	GAGTTTCGTT	TATGGAGGCT	360
GCTTGGGCAA	CAAGAACAAC	TACCTTCGGG	AAGAAGAGTG	CATTCTANCC	TGTCNGGGTG	420
TGCAAGGTGG	GCCTTTGANA	NGCANCTCTG	GGGCTCANGC	GACTTTCCCC	CAGGGCCCCCT	480
CCATGGAAAG	GCGCCATCCA	NTGTTCTCTG	GCACCTGTCA	GCCCACCCAG	TTCCGCTGCA	540
NCAATGGCTG	CTGCATCNAC	ANTTTCCTNG	AATTGTGACA	ACACCCCCCA	NTGCCCCCAA	600
CCCTCCCAAC	AAAGCTTCCC	TGTTNAAAAA	TACNCCANTT	GGCTTTTNAC	AAACNCCCCG	660
CNCCTCCNTT	TTCCCCNNTN	AACAAAGGGC	NCTNGCNTTT	GAACTGCCCN	AACCCNGGAA	720
TCTNCCNNGG	AAAAANTNCC	CCCCCTGGTT	CCTNNAANCC	CCTCCNCNAA	ANCTNCCCCC	780
CCC						783

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 801 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGCCCAATTC	CAGCTGCCAC	ACCACCCACG	GTGACTGCAT	TAGTTCGGAT	GTCATACAAA	60
AGCTGATTGA	AGCAACCCCTC	TACTTTTTTG	TCGTGAGCCT	TTTGCTTGGT	GCAGGTTTCA	120
TTGGCTGTGT	TGGTGACGTT	GTCAATTGCAA	CAGAATGGGG	GAAAGGCACT	GTTCTCTTTG	180
AAGTAGGGTG	AGTCCTCAAA	ATCCGTATAG	TTGGTGAAGC	CACAGCACTT	GAGCCCTTTC	240
ATGGTGGTGT	TCCACACTTG	AGTGAAGTCT	TCCTGGGAAC	CATAATCTTT	CTTGATGGCA	300
GGCACTACCA	GCAACGTCAG	GAAGTGCTCA	GCCATTGTGG	TGTACACCAA	GGCGACCACA	360
GCAGCTGCAA	CCTCAGCAAT	GAAGATGAGG	AGGAGGATGA	AGAAGAACGT	CNCGAGGGCA	420
CACTTGCTCT	CCGTCTTAGC	ACCATAGCAG	CCCANGAAAC	CAAGAGCAAA	GACCACAACG	480
CCNGCTGCGA	ATGAAAGAAA	NTACCCACGT	TGACAAACTG	CATGGCCACT	GGACGACAGT	540

TGGCCCGAAN	ATCTTCAGAA	AAGGGATGCC	CCATCGATTG	AACACCCANA	TGCCCCTGTC	600
CNACAGGGCT	GCNCCNCNCN	GAAAGAATGA	GCCATTGAAG	AAGGATCNTC	NTGGTCTTAA	660
TGAACTGAAA	CCNTGTCATG	TGGCCCTGT	TCAGGGCTCT	TGGCAGTGAA	TTCTGANAAA	720
AAGGAACNGC	NTNAGCCCCC	CCAAANGANA	AAACACCCCC	GGGTGTTGCC	CTGAATTGGC	780
GGCCAAGGAN	CCCTGCCCCN	G				801

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 740 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GTGAGAGCCA	GGCGTCCCTC	TGCCTGCCCC	CTCAGTGGCA	ACACCCGGGA	GCTGT'TTTGT	60
CCTTTGTGGA	GCCTCAGCAG	TTCCCTCTTT	CAGAACTCAC	TGCCAAGAGC	CCTGAACAGG	120
AGCCACCATG	CAGTGCTTCA	GC'TTCATTAA	GACCATGATG	ATCCTCTTCA	ATTTGCTCAT	180
CTTTCTGTGT	GGTGCAGCCC	TGTTGGCAGT	GGGCATCTGG	GTGTCAATCG	ATGGGGCATC	240
CTTTCTGAAG	ATCTTCGGGC	CACTGTCTGC	CAGTGCCATG	CAGTTTGTCA	ACGTGGGCTA	300
CTTCCTCATC	GCAGCCGGCG	TTGTGGTCTT	TGCTCTTGGT	TTCTGGGCT	GCTATGGTGC	360
TAAGACGGAG	AGCAAGTGTG	CCCTCGTGAC	GTTCTTCTTC	ATCCTCCTCC	TCATCTTCAT	420
TGCTGAAGTT	GCAGCTGCTG	TGGTCGCCCT	GGTGTAACAC	ACAATGGCTG	AACCATTCCCT	480
GACGTTGCTG	GTANTGCCCTG	CCATCAANAA	AGATTATGGG	TTCCCAGGAA	AAATTCACCTC	540
AANTNTGGAA	CACCNCCATG	AAAAGGGCTC	CAATTTCTGN	TGGCTTCCCC	AACATAACCG	600
GAATTTTGAA	AGANTCNCCC	TACTTCCAAA	AAAAAANANT	TGCCTTTNCC	CCCN'TTCTGT	660
TGCAATGAAA	ACNTCCCAAN	ACNGCCAATN	AAAACCTGCC	CNNNCAAAAA	GGNTCNCAAA	720
CAAAAAAANT	NNAAGGG'TTN					740

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 802 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CCGCTGGTTG	CGCTGGTCCA	GNGNAGCCAC	GAAGCACGTC	AGCATACACA	GCCTCAATCA	60
CAAGGTCTTC	CAGCTGCCGC	ACATTACGCA	GGGCAAGAGC	CTCCAGCAAC	ACTGCATATG	120
GGATACACTT	TACTTTAGCA	GCCAGGGTGA	CAACTGAGAG	GTGTCGAAGC	TTATTCTTCT	180
GAGCCTCTGT	TAGTGGAGGA	AGATTCCGGG	CTTCAGCTAA	GTAGTCAGCG	TATGTCCCAT	240
AAGCAAACAC	TGTGAGCAGC	CGGAAGGTAG	AGGCAAAGTC	ACTCTCAGCC	AGCTCTCTAA	300
CATTGGGCAT	GTCCAGCAGT	TCTCCAAACA	CGTAGACACC	AGNGGCCTCC	AGCACCTGAT	360
GGATGAGTGT	GGCCAGCGCT	GCCCCCTTGG	CCGACTTGGC	TAGGAGCAGA	AATTGCTCCT	420
GGTTCTGCCC	TGTCACCTTC	ACTTCCGCAC	TCATCACTGC	ACTGAGTGTG	GGGGACTTGG	480
GCTCAGGATG	TCCAGAGACG	TGGTTCGGCC	CCCTCNCTTA	ATGACACCGN	CCANNCAACC	540
GTCGGCTCCC	GCCGANTGNG	TTCGTCGTNC	CTGGGTCAGG	GTCTGCTGGC	CNCTACTTGC	600
AANCTTCGTC	NGGCCCATGG	AATTCAACCNC	ACCGGAACTN	GTANGATCCA	CTNN'TTCTAT	660
AACCGGNCGC	CACCGCNNNT	GGAAC'TCCAC	TCTTNTTNCC	TTTACTTGAG	GGTTAAGGTC	720
ACCTTNNCG	TTACCTTGGT	CCAAACCN'TN	CCNTGTGTCTG	ANATNGTNAA	TCNGGNCCNA	780
TNCCANCCNC	ATANGAAGCC	NG				802

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 731 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CNAAGCTTCC	AGGTNACGGG	CCGCNAANCC	TGACCCNAGG	TANCANAANG	CAGNCNGCGG	60
GAGCCCACCG	TCACGNGGNG	GNGTCTTTAT	NGGAGGGGGC	GGAGCCACAT	CNCTGGACNT	120
CNTGACCCCA	ACTCCCNCC	NCNCANTGCA	GTGATGAGTG	CAGAAGTGAA	GGTNACGTGG	180
CAGGAACCAA	GANCAAANNC	TGCTCCNNTC	CAAGTCGGCN	NAGGGGGCGG	GGCTGGCCAC	240
GCNCATCCNT	CNAGTGCTGN	AAAGCCCCNN	CCTGTCTACT	TGTTTGAGAG	ACNGCNNNGA	300
CATGCCCAGN	GTTANATAAC	NGGCNGAGAG	TNANTTTGCC	TCTCCCTTCC	GGCTGCGCAN	360
CGNGTINTGCT	TAGNGGACAT	AACCTGACTA	CTTAAGTGAA	CCCNNGAATC	TNCCNCCCCT	420
CCACTAAGCT	CAGAACAAAA	AACTTCGACA	CCACTCANTT	GTCACCTGNC	TGCTCAAGTA	480
AAGTGTACCC	CATNCCCAAT	GTNTGCTNGA	NGCTCTGNCC	TGCNTTANGT	TCGGTCCTGG	540
GAAGACCTAT	CAATTNAAGC	TATGTTTCTG	ACTGCCTCTT	GCTCCCTGNA	ACAANCNACC	600
CNNC NNTCCA	AGGGGGGGNC	GGCCCCCAAT	CCCCCAACC	NTNAATTNAN	TTTANCCCCN	660
CCCCCNGGCC	CGGCCTTTTA	CNANCNTCNN	NNACNGGGNA	AAACCNNGC	TTTNCCCCAAC	720
NNAATCCNCC	T					731

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 754 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

TTTTTTTTTTT	TTTTTTTTTTT	TAAAAACCCC	CTCCATTNAA	TGNAACTTC	CGAAATTGTC	60
CAACCCCCTC	NTCCAAATNN	CCNTTTCGG	GNGGGGGTTC	CAAACCCAAN	TTANNTTGG	120
ANNTTAAATT	AAATNTTNT	TGGNGGNNA	ANCCNAATGT	NANGAAAGTT	NAACCCANTA	180
TNANCTTNAA	TNCCTGAAA	CCNGTNGNTT	CCAAAAATNT	TTAACCTTA	ANTCCCTCCG	240
AAATNGTTNA	NGGAAAACCC	AANTTCTCNT	AAGGTTGTTT	GAAGGNTNAA	TNAAAANCCC	300
NNCCAATTGT	TTTTNGCCAC	GCCTGAATTA	ATTGGNTTCC	GNTGTTTTCC	NTTAAAANAA	360
GGNNANCCCC	GGTTANTNAA	TCCCCCNCC	CCCAATTATA	CCGANTTTTT	TTNGAATTGG	420
GANCCNCGG	GAATTAACGG	GGNNNNTCCC	TNTTGGGGGG	CNGGNNCCCC	CCCCNTCGGG	480
GGTTNGGGNC	AGGNCNNAAT	TGTTTAAGGG	TCCGAAAAAT	CCCTCCNAGA	AAAAANCTC	540
CCAGGNTGAG	NNTINGGGTTT	NCCCCCCCC	CANGGCCCTT	CTCGNANAGT	TGGGGTTTGG	600
GGGGCCTGGG	ATTTTNTTTC	CCCTNTNTCC	TCCCCCCCC	CCNGGGANAG	AGGTTNGNGT	660
TTTGNTCNNC	GGCCCCNCCN	AAGANCTTTN	CCGANTTNAN	TTAAATCCNT	GCCTNGGCGA	720
AGTCCNTTGN	AGGGNTAAAN	GGCCCCCTNN	CGGG			754

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 755 base pairs
 (B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

ATCANCCCAT	GACCCCNAAAC	NNGGGACCNC	TCANCCGGNC	NNNCNACCNC	CGGCCNATCA	60
NNGTNAGNNC	ACTNCNNTTN	NATCACNCCC	CNCCNACTAC	GCCCNCNANC	CNACGCNCTA	120
NNCANATNCC	ACTGANNGCG	CGANGTNGAN	NGAGAAANCT	NATACCANAG	NCACCANACN	180
CCAGCTGTCC	NANAANGCCT	NNNATAACNG	NNNATCCAAT	NTGNANCCCT	CNAAGTATTN	240
NNCNCANAT	GATTTTCCCTN	ANCCGATTAC	CCNTNCCCCC	TANCCCCCTC	CCCCCAACNA	300
CGAAGGCNCT	GGNCCNAAGG	NNGCGNCNCC	CCGCTAGNTC	CCCNCAAGT	CNCNCNCCTA	360
AACTCANCCN	NATTACNCGC	TTCNTGAGTA	TCACTCCCCG	AATCTCACCC	TACTCAACTC	420
AAAAANATCN	GATACAAAAT	AATNCAAGCC	TGNTTATNAC	ACTNTGACTG	GGTCTCTATT	480
TTAGNGGTCC	NTNAANCNTC	CTAATACTTC	CAGTCTNCCCT	TCNCCAATTT	CCNAANGGCT	540
CTTTCNGACA	GCAATNTTTTG	GTTCCCNNTT	GGGTCTTAN	NGAATTGCCC	TTCNTNGAAC	600
GGGCTCNTCT	TTTCCTTCGG	TTANCTGGN	TTCNNCCGGC	CAGTTATTAT	TTCCCNTTTT	660
AAATTCNTNC	CNTTTANTTT	TGGCNTTCNA	AACCCCCGGC	CTTGAAAACG	GCCCCCTGGT	720
AAAAGGTTGT	TTTGANAAAA	TTTTTGTTTT	GTTCC			755

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 849 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TTTTTTTTTT	TTTTTANGTG	TNGTCGTGCA	GGTAGAGGCT	TACTACAANT	GTGAANACGT	60
ACGCTNGGAN	TAANGCGACC	CGANTTCTAG	GANNCNCCCT	AAAATCANAC	TGTGAAGATN	120
ATCCTGNMNA	CGGAANGGTC	ACCGGNNGAT	NNTGCTAGGG	TGNCCNCTCC	CANNNCNNTN	180
CATAACTCNG	NGGCCCTGCC	CACCACCTTC	GGCGGCCCCG	NGNCCGGGCC	CGGGTCATTN	240
GNNTTAACCN	CACTNNGCNA	NCGGTTTCCN	NCCCCNNCNG	ACCCNGGCCA	TCCGGGGTNC	300
TCTGTCTTCC	CCTGNAGNCN	ANAAANTGGG	CCNCGGNCCC	CTTTACCCCT	NNACAAGCCA	360
CNGCCNTCTA	NCCNCNGCCC	CCCCCTCCANT	NNGGGGGACT	GCCNANNGCT	CCGTTNCTNG	420
NNACCCCNNN	GGGTNCCCTCG	GTTGTCTGANT	CNACCGNANG	CCANGGATTTC	CNAAGGAAGG	480
TGCGTTNTTG	GCCCCTACCC	TTCGCTNCGG	NNCACCCCTTC	CCGACNANGA	NCCGCTCCCG	540
CNCNNGNNG	CCTCNCCTCG	CAACACCCGC	NCTCNTCNGT	NCGGNNNCCC	CCCCACCCGC	600
NCCCTCNCNC	NGNCGNANCN	CTCCNCCNCC	GTCTCANNCA	CCACCCCGCC	CCGCCAGGCC	660
NTCANCCACN	GGNNGACNNG	NAGCNCNNTC	GCNCCGCGCN	GCGNCNCCCT	CGCCNCNGAA	720
CTNCNTCNGG	CCANTNNCGC	TCAANCCNNA	CNAAACGCCG	CTGCGCGGCC	CGNAGCGNCC	780
NCCTCCNCGA	GTCTCTCCGN	CTTCCNACCC	ANGNNTTCCN	CGAGGACACN	NNACCCCGCC	840
NNCANGCGG						849

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 872 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GCGCAA	ACTA	TACTTC	GCCTC	GNACTC	GCCTC	TCTTTT	CCTC	CGCAAC	CATG	60
TCTGAC	NANC	CCGATT	NNGC	NGATAT	CNAN	AAGNTC	CGANC	AGTCCA	AACT	120
CACACN	CNAN	AGANAA	ATCC	NCTGCT	TCC	ANAGTA	NACN	ATTGAA	CNNG	180
NGGCGA	ATCG	TAATNA	GGCG	TGCGCC	GCCA	ATNTGT	CNCC	GTTTAT	TNTN	240
CTNCCN	ACCC	TACNTC	TTCN	NAGCTG	TCNN	ACCCCT	NGTN	CGNAC	CCCC	300
TCGGGT	TTNN	NNTGAC	CGNG	CNNCCC	TCC	CCCCNT	TCCAT	NACGAN	CNCN	360
NANNGC	NCGC	NCCCCG	MNCT	CTTCGC	CNCC	CTGTCT	TNTN	CCCCTG	TNGC	420
ACCGCA	TTGA	CCCTCG	CCNN	CTNCNN	GAAA	NCGNAN	ACGT	CCGGGT	TGNN	480
TGGGNN	NGCG	TCTGCN	CCGC	GTTCTT	TCCN	NCNNCT	TCCA	CCATCT	TCNT	540
CCNCGC	CNTC	TCNNNC	CACN	CCTGGG	ACGC	TNTCCT	TNTGC	CCCCCT	TNAC	600
CGNCGT	GNCC	CGNCCCC	ACC	NTCATT	TNCA	NACGNT	CTTC	ACAANN	NCCT	660
CNANCN	GNCN	GTCANCC	NAG	GGAAGG	GNGG	GGNNCC	NNTG	NTTGAC	GTTG	720
CGAANAN	TCC	TCNCCN	TCAN	CNCTAC	CCCT	CGGGCG	MNCT	CTCNGT	TNCC	780
NTCTCCCC	CG	NGNGCN	CNTC	TCAGCT	CNC	CCNCCCC	NCT	CTCTGC	CANTG	840
TNACCN	NTAC	GANTNT	TCGN	CNCCCT	CCTTT	CC				872

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 815 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GCATGCA	AAGC	TTGAGT	ATTTC	TATAGN	GTCA	CCTAAAT	ANC	TTGGC	NTAAT	CATGGT	CNTA	60
NCTGNCT	TCC	TGTGTCA	AAAT	GTATAC	NAAN	TANATAT	GAA	TCTNAT	NTGA	CAAGAN	NGTA	120
TCNTNC	CATTA	GTAACA	ANTG	TNNTGT	TCCAT	CCTGTC	NGAN	CANATT	CCCCA	TNNATT	NCGN	180
CGCATT	CNCN	GCNCANT	ATN	TAATNG	GGGAA	NTCNMT	TNMN	NCACCN	NCAT	CTATCN	TNCC	240
GCNCCCT	GAC	TGGNAG	AGAT	GGATNA	NNTTC	TNNTNT	GACC	NACATG	TTC	TCTTGG	ATTN	300
AAANAN	CCCCC	CGCNGN	CCAC	CGGTTN	GNGG	CNAGCC	NNTC	CCAAGAC	CCTC	CTGTGG	GAGGT	360
AACCTG	CGTC	AGANNCA	TCA	AACNTG	GGGAA	ACCCGC	NNCC	ANGTNNA	AAGT	NGNNNC	CANAN	420
GATCCCC	GTCC	AGGNTT	NACC	ATCCCT	TCNC	AGCGCC	CCCCCT	TTNGTG	CCTT	ANAGNG	NAGC	480
GTGTCC	NANC	CNCTCA	ACAT	GANACG	CGCC	AGNCCAN	CCG	CAATTN	GGCA	CAATGT	CGNC	540
GAACCCC	CCTA	GGGGGA	NTNA	TNCAAAN	CCCC	CAGGAT	TGTC	CNCNCAN	GAA	ATCCCN	CANC	600
CCNCCC	TAC	CCNNCT	TTGG	GACNGT	GACC	AANTCCC	GGA	GTNCCAG	TCC	GGCCNG	NCTC	660
CCCCACC	GGT	NNCCNT	GGGG	GGGTGA	ANCT	CNGNNT	CANC	CNGNCG	GAGN	NTCGNA	AAGGA	720
ACCGGN	CCTN	GGNCGA	ANNG	ANCNNT	CNGA	AGNGCC	CNCNT	CGTATA	AACC	CCCCTC	NC	780
NCCNAC	NGNT	AGNTCCCC	C	CNGGGT	NC	CG	AANG					815

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 775 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CCGAGATGTC	TCGCTCCGTG	GCCTTAGCTG	TGCTCGCGCT	ACTCTCTCTT	TCTGGCCTGG	60
AGGCTATCCA	GCGTACTCCA	AAGATTGAGG	TTTACTCACG	TCATCCAGCA	GAGAATGGAA	120
AGTCAAATTT	CCTGAATTGC	TATGTGTCTG	GGTTTCATCC	ATCCGACATT	GAANTTGACT	180
TACTGAAGAA	TGGANAGAGA	ATTGAAAAAG	TGGAGCATTC	AGACTTGTCT	TTCAGCAAGG	240
ACTGGTCTTT	CTATCTCNTG	TACTACACTG	AATTCACCCC	CACTGAAAAA	GATGAGTATG	300
CCTGCCGTGT	GAACCATGTG	ACTTTGTGAC	AGCCCAAGAT	AGTTAAGTGG	GATCGAGACA	360
TGTAAGCAGN	CNNCATGGAA	GTTTGAAGAT	GCCGCATTTG	GATTGGATGA	ATTCCAAATT	420
CTGCTTGCTT	GCNTTTTAAT	ANTGATATGC	NTATACACCC	TACCCTTTAT	GNCCCCAAAT	480
TGTAGGGGTT	ACATNANTGT	TCNCNTNGGA	CATGATCTTC	CTTTATAANT	CCNCCNTTCG	540
AATTGCCCGT	CNCCCNGTTN	NGAATGTTTC	CNNAACCACG	GTTGGCTCCC	CCAGGTCNCC	600
TCTTACGGAA	GGGCCTGGGC	CNCTTTNCAA	GGTTGGGGGA	ACCNAAAATT	TCNCTTNTGC	660
CCNCCNCCA	CNNTCTTGNG	NNCNCANTTT	GGAACCCTTC	CNATTCCCCT	TGGCCTCNNA	720
NCCTTNNCTA	ANAAAACCTT	AAANCGTNGC	NAAANNTTTN	ACTTCCCCCC	TTACC	775

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 820 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

ANATTANTAC	AGTGTAATCT	TTTCCCAGAG	GTGTGTANAG	GGAACGGGGC	CTAGAGGCAT	60
CCCANAGATA	NCTTATANCA	ACAGTGCTTT	GACCAAGAGC	TGCTGGGCAC	ATTTCTTGCA	120
GAAAAGGTGG	CGGTCCCCAT	CACTCCTCCT	CTCCCATAGC	CATCCCAGAG	GGGTGAGTAG	180
CCATCANGCC	TTCGGTGCGA	GGGAGTCANG	GAAACAACAN	ACCACAGAGC	ANACAGACCA	240
NTGATGACCA	TGGGCGGGAG	CGAGCCTCTT	CCCTGNACCG	GGGTGGCANA	NGANAGCCTA	300
NCTGAGGGGT	CACACTATAA	ACGTTAACGA	CCNAGATNAN	CACCTGCTTC	AAGTGCACCC	360
TTCTTACCTG	ACNACCAGNG	ACCNNNAACT	GCNGCCTGGG	GACAGCNCTG	GGANCAGCTA	420
ACNNAGCACT	CACCTGCCCC	CCCATGGCCG	TNCGCNTCCC	TGGTCCTGNC	AAGGGAAGCT	480
CCCTGTTGGA	ATTNCGGGGA	NACCAAGGGA	NCCCCCTCCT	CCANCTGTGA	AGGAAAAANN	540
GATGGAATTT	TNCCCTTCCG	GCCNNTCCCC	TCTTCTTTTA	CACGCCCCCT	NNTACTCNTC	600
TCCCTCTNTT	NTCCTGNCNC	ACTTTTNACC	CCNNNATTTT	CCTTNATTGA	TCGGANNCTN	660
GANATTCCAC	TNNCGCCTNC	CNTCNATCNG	NAANACNAAA	NACTNTCTNA	CCNCGGGGAT	720
GGGNNCCTCG	NTCATCCTCT	CTTTTTCNCT	ACCNCCNNTT	CTTTGCCTCT	CCTTNGATCA	780
TCCAACCNTC	GNTGGCCNTN	CCCCCCCNNN	TCCTTTNCCC			820

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 818 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

TCTGGGTGAT	GGCCTCTTCC	TCCTCAGGGA	CCTCTGACTG	CTCTGGGCCA	AAGAATCTCT	60
TGTTTCTTCT	CCGAGCCCCA	GGCAGCGGTG	ATTGAGCCCT	GCCCAACCTG	ATTCTGATGA	120
CTGCGGATGC	TGTGACGGAC	CCAAGGGGCA	AATAGGGTCC	CAGGGTCCAG	GGAGGGGCGC	180
CTGCTGAGCA	CTTCCGCCCC	TCACCCTGCC	CAGCCCCTGC	CATGAGCTCT	GGGCTGGGTC	240
TCCGCCTCCA	GGGTTCTGCT	CTTCCANGCA	NGCCANCAAG	TGGCGCTGGG	CCACACTGGC	300

TTCCTCCTGC	CCCNTCCCTG	GCTCTGANTC	TCTGTCTTCC	TGTCCTGTGC	ANGCNCCTTG	360
GATCTCAGTT	TCCCTCNCCTC	ANNGAACTCT	GTTTCTGANN	TCTTCANTTA	ACTNTGANTT	420
TATNACCNAN	TGGNCTGTNC	TGTCNNACTT	TAATGGGCCN	GACCGGCTAA	TCCCTCCCTC	480
NCTCCCTTCC	ANTTCNNNNA	ACCNGCTTNC	CNTCNTCTCC	CCNTANCCCG	CCNGGGAANC	540
CTCCTTTGCC	CTNACCANGG	GCCNNNACCG	CCCNTNNCTN	GGGGGGCNGG	GTNNCTNCNC	600
CTGNTNNCCC	CNCTCNCNNT	TNCCTCGTCC	CNNCNCNCGN	NNGCANNTTC	NCNGTCCCN	660
TNNCTCTTCN	NGTNTCGNAA	NGNTCNCNTN	TNNNNNGNCN	NGNTNNTNCN	TCCCTCTCNC	720
CNNNTGNANG	TNNTTNNNNC	NCNGNNCCCC	NNNNCNNNNN	NGGNNNTNNN	TCTNCNCNGC	780
CCCNCCCCC	NGNATTAAGG	CCTCCNNTCT	CCGGCCNC			818

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 731 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

AGGAAGGGCG	GAGGGATATT	GTANGGGATT	GAGGGATAGG	AGNATAANGG	GGGAGGTGTG	60
TCCCAACATG	ANGGTGNNGT	TCTCTTTTGA	ANGAGGGTTG	NGTTTTTANN	CCNGGTGGGT	120
GATTNAACCC	CATTGTATGG	AGNNAAGGN	TTTNAGGGAT	TTTTCGGCTC	TTATCAGTAT	180
NTANATTCCCT	GTNAATCGGA	AAATNATNTT	TCNNCNGGAA	AATNTTGCTC	CCATCCGNAA	240
ATTNCTCCCG	GGTAGTGCAT	NTTNGGGGGN	CNGCCANGTT	TCCCAGGCTG	CTANAATCGT	300
ACTAAAGNTT	NAAGTGGGAN	TNCAAATGAA	AACCTNNCAC	AGAGNATCCN	TACCCGACTG	360
TNNNTTNCCT	TCGCCCTNTG	ACTCTGCNNG	AGCCCAATAC	CCNNGNGNAT	GTNCCCNGN	420
NNNGCGNCNC	TGAAANNNNC	TCGNGGCTNN	GANCATCANG	GGGTTTCGCA	TCAAAAGCNC	480
CGTTTCNCAT	NAAGGCACTT	TNGCCTCATC	CAACCNTNG	CCCTCNCNCA	TTTNGCCGTC	540
NGGTTCNCCT	ACGCTNNNTG	CNCCTNNTN	GANATTTTNC	CCGCCTNGGG	NAANCCTCCT	600
GNAATGGGTA	GGGNCCTTNT	TTTTNACCNN	GNGGTNTACT	AATCNNCTNC	ACGCNTNCTT	660
TCTCNACCCC	CCCCCTTTT	CAATCCCAN	GGCNAATGGG	GTCTCCCCNN	CGANGGGGGG	720
NNNCCCANNC	C					731

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 822 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

ACTAGTCCAG	TGTGGTGGA	TTCCATTGTG	TTGGGGNCNC	TTCTATGANT	ANTNTTAGAT	60
CGCTCANACC	TCACANCTC	CCNACNANGC	CTATAANGAA	NANNAATAGA	NCTGTNCNNT	120
ATNTNTACNC	TCATANNCCT	CNNNACCCAC	TCCCTCTTAA	CCCTACTGT	GCCTATNGCN	180
TNNCTANTCT	NTGCCGCCTN	CNANCCACCN	GTGGGCCNAC	CNCNNGNATT	CTCNATCTCC	240
TCNCCATNTN	GCCTANANTA	NGTNCATACC	CTATACCTAC	NCCAATGCTA	NNNCTAANCN	300
TCCATNANTT	ANNNTAACTA	CCACTGACNT	NGACTTTTCNC	ATNANCTCCT	AATTTGAATC	360
TACTCTGACT	CCCACNGCCT	ANNNATTAGC	ANCNTCCCCC	NACNATNTCT	CAACCAAATC	420
NTCAACAACC	TATCTANCTG	TTCNCCAACC	NTTNCCCTCCG	ATCCCCCNAC	AACCCCCCTC	480
CCAAATACCC	NCCACCTGAC	NCCTAACCCN	CACCATCCCCG	GCAAGCCNAN	GGNCATTTAN	540
CCACTGGAAT	CACNATNGGA	NAAAAAAAC	CCNAACTCTC	TANCNCNNAT	CTCCCTAANA	600

AATNCTCCTN	NAATTTACTN	NCANTNCCAT	CAANCCCACN	TGAAACNNAA	CCCCTGTTTT	660
TANATCCCTT	CTTTCGAAAA	CCNACCCTTT	ANNNCCCAAC	CTTTNNGGGCC	CCCCCNCTNC	720
CCNAATGAAG	GNCNCCCAAT	CNANGAAACG	NCCNTGAAAA	ANCNAGGCNA	ANANNNTCCG	780
CANATCCTAT	CCCTTANTTN	GGGGNCCCCTT	NCCCNNGGGCC	CC		822

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 787 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

CGGCCGCCTG	CTCTGGCACA	TGCCTCCTGA	ATGGCATCAA	AAGTGATGGA	CTGCCCATTG	60
CTAGAGAAGA	CCTTCTCTCC	TACTGTCTAT	ATGGAGCCCT	GCAGACTGAG	GGCTCCCCTT	120
GTCTGCAGGA	TTTGATGTCT	GAAGTCGTGG	AGTGTGGCTT	GGAGCTCCTC	ATCTACATNA	180
GCTGGAAGCC	CTGGAGGGCC	TCTCTCGCCA	GCCTCCCCCT	TCTCTCCACG	CTCTCCANGG	240
ACACCAGGGG	CTCCAGGCAG	CCCATTATTC	CCAGNANGAC	ATGGTGTTTC	TCCACGCGGA	300
CCCATGGGGC	CTGNAAGGCC	AGGGTCTCCT	TTGACACCAT	CTCTCCCGTC	CTGCCTGGCA	360
GGCCGTGGGA	TCCACTANTT	CTANAACGGN	CGCCACCNCG	GTGGGAGCTC	CAGCTTTTGT	420
TCCCNTTAAT	GAAGGTTAAT	TGCNCGCTTG	GCGTAATCAT	NGGTCANAAC	TNTTTCCTGT	480
GTGAAATTGT	TTNTCCCCTC	NCNATTCCNC	NCNACATACN	AACCCGGAAN	CATAAAGTGT	540
TAAAGCCTGG	GGGTNGCCTN	NNGAATNAAC	TNAACTCAAT	TAATTGCGTT	GGCTCATGGC	600
CCGCTTTCCN	TTCNGGAAAA	CTGTCNTCCC	CTGCNTTNNT	GAATCGGCCA	CCCCCNGGG	660
AAAAGCGGTT	TGCNTTTTNG	GGGGNTCCTT	CCNCTTCCCC	CCTCNCTAAN	CCCTNCGCCT	720
CGGTCGTTNC	NGGTNGCGGG	GAANGGGNAT	NNNCTCCCNC	NAAGGGGGNG	AGNNNGNTAT	780
CCCCAAA						787

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 799 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

TTTTTTTTTT	TTTTTTTGGC	GATGCTACTG	TTTAATTGCA	GGAGGTGGGG	GTGTGTGTAC	60
CATGTACCAG	GGCTATTAGA	AGCAAGAAGG	AAGGAGGGAG	GGCAGAGCGC	CCTGCTGAGC	120
AACAAAGGAC	TCCTGCAGCC	TTCTCTGTCT	GTCTCTTGGC	GCAGGCACAT	GGGGAGGCCT	180
CCCGCAGGGT	GGGGGCCACC	AGTCCAGGGG	TGGGAGCACT	ACANGGGGTG	GGAGTGGGTG	240
GTGGCTGGTN	CNAATGGCCT	GNCACANATC	CCTACGATTG	TTGACACCTG	GATTTACCA	300
GGGGACCTTC	TGTTCTCCCA	NGGNAACTTC	NTNNATCTCN	AAAGAACACA	ACTGTTTCTT	360
CNGCANTTCT	GGCTGTTTCAT	GGAAAGCACA	GGTGTCNAT	TTNGGCTGGG	ACTTGGGTACA	420
TATGGTTCCG	GCCCACCTCT	CCCNCTCNAAN	AAGTAATTCA	CCCCCCCCCN	CCNTCTNNTG	480
CCTGGGCCCT	TAANTACCCA	CACCGGAACT	CANTTANTTA	TTCATCTTNG	GNTGGGCTTG	540
NTNATCNCCN	CCTGAANGCG	CCAAGTTGAA	AGGCCACGCC	GTNCCCNCTC	CCCATAGNAN	600
NTTTTNNCNT	CANCTAATGC	CCCCCNGGC	AACNATCCAA	TCCCCCCCCN	TGGGGGCCCC	660
AGCCCANGGC	CCCCGNCTCG	GGNNNCCNGN	CNCGNANTCC	CCAGGNTCTC	CCANTCNGNC	720
CCNNNGCNCC	CCCGCACGCA	GAACANAAGG	NTNGAGCCNC	CGCANNNNNN	NGGTNNCNAC	780
CTCGCCCCCC	CCNNCGNNG					799

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 789 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	60
TTTNNCCNAG	GGCAGGTTTA	TTGACAACCT	CNCGGGACAC	AANCAGGCTG	GGGACAGGAC	120
GGCAACAGGC	TCCGGCGGCG	GCGGCGGCGG	CCCTACCTGC	GGTACCAAAT	NTGCAGCCTC	180
CGCTCCCGCT	TGATNTTCCT	CTGCAGCTGC	AGGATGCCNT	AAAACAGGGC	CTCGGCCNTN	240
GGTGGGCACC	CTGGGATTNN	AATTTCCACG	GGCACAATGC	GGTCGCANCC	CCTCACCACC	300
NATTAGGAAT	AGTGGTNTTA	CCCNCCNCCG	TTGGGCNCACT	CCCCNTGGAA	ACCAC'TTNTC	360
GCGGCTCCGG	CATCTGGTCT	TAAACCTTGC	AAACNCTGGG	GCCCTCTTTT	TGGTTANTNT	420
NCCNGCCACA	ATCATNACTC	AGACTGGCNC	GGGCTGGCCC	CAAAAAANCN	CCCCAAAACC	480
GGNCCATGTC	TTNNCGGGGT	TGCTGCNATN	TNCATCACCT	CCCGGGCNCA	NCAGGNCAAC	540
CCAAAAGTTC	TTGNNGGCCN	CAAAAAANCT	CCGGGGGGNC	CCAGTTTCAA	CAAAGTCATC	600
CCCCTTGGCC	CCCAAATCCT	CCCCCGNTT	NCTGGGTTTG	GGAACCCACG	CCTCTNNCTT	660
TGGNNGGCAA	GNTGGNTCCC	CCTTCGGGCC	CCCGGTGGGC	CCNNCTCTAA	NGAAAAACNC	720
NTCCTNNNCA	CCATCCCCC	NNGNNACGNC	TANCAANGNA	TCCCTTTTTT	TANAAACGGG	780
CCCCCNCG						789

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 793 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GACAGAACAT	GTTGGATGGT	GGAGCACCTT	TCTATACGAC	TTACAGGACA	GCAGATGGGG	60
AATTCATGGC	TGTTGGAGCA	ATANAACCCC	AGTTCTACGA	GCTGCTGATC	AAAGGACTTG	120
GACTAAAGTC	TGATGAAC'TT	CCCAATCAGA	TGAGCATGGA	TGATTGGCCA	GAAATGAANA	180
AGAAGTTTGC	AGATGTATTT	GCAAAGAAGA	CGAAGGCAGA	GTGGTGTCAA	ATCTTTTGACG	240
GCACAGATGC	CTGTGTGACT	CCGGTTCTGA	CTTTTGAGGA	GGTTGTTCAT	CATGATCACA	300
ACAANGAACG	GGGCTCGTTT	ATCACCANTG	AGGAGCAGGA	CGTGAGCCCC	CGCCCTGCAC	360
CTCTGCTGTT	AAACACCCCA	GCCATCCC'TT	CTTTCAAAAG	GGATCCACTA	CTTCTAGAGC	420
GGNCGCCACC	GCGGTGGAGC	TCCAGCTTTT	GTTCCCTTTA	GTGAGGGTTA	ATTGCGCGCT	480
TGGCGTAATC	ATGGTCATAN	CTGTTTCTCTG	TGTGAAATTG	TTATCCGCTC	ACAATTCCAC	540
ACAACATACG	ANCCGGAAGC	ATNAAATTTT	AAAGCCTGGN	GGTNGCCTAA	TGANTGAACT	600
NACTCACATT	AAT'TGGCTTT	GCGCTCACTG	CCCGCTTTCC	AGTCCGGAAA	ACCTGTCC'TT	660
GCCAGCTGCC	NTTAATGAAT	CNGGCCACCC	CCCGGGGAAA	AGGCNGTTTG	CTTNTTTGGGG	720
CGCNCTTCCC	GCTTTCTCGC	TTCTTGAANT	CCTTCCCCC	GGTCTTTTCGG	CTTGCGGCNA	780
ACGGTATCNA	CCT					793

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 756 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GCCGCGACCG	GCATGTACGA	GCAACTCAAG	GGCGAGTGGA	ACCGTAAAAG	CCCCAATCTT	60
ANCAAGTGCG	GGGAANAGCT	GGGTCGACTC	AAGCTAGTTC	TTCTGGAGCT	CAACTTCTTG	120
CCAACCACAG	GGACCAAGCT	GACCAAACAG	CAGCTAATTC	TGGCCCGTGA	CATACTGGAG	180
ATCGGGGCCC	AATGGAGCAT	CCTACGCAAN	GACATCCCCCT	CCTTCGAGCG	CTACATGGCC	240
CAGCTCAAAT	GCTACTACTT	TGATTACAAN	GAGCAGCTCC	CCGAGTCAGC	CTATATGCAC	300
CAGCTCTTGG	GCCTCAACCT	CCTCTTCCCTG	CTGTCCCAGA	ACCGGGTGGC	TGANTNCCAC	360
ACGGANTTGG	ANCGGCTGCC	TGCCCAANGA	CATACANACC	AATGTCTACA	TCNACCACCA	420
GTGTCCCTGGA	GCAATACTGA	TGGANGGCAG	CTACCNCAAA	GTNTTCCTGG	CCNAGGGTAA	480
CATCCCCCGC	CGAGAGCTAC	ACCTTCTTTCA	TTGACATCCT	GCTCGACACT	ATCAGGGATG	540
AAAATCGCNG	GGTTGCTCCA	GAAAGGCTNC	AANAANATCC	TTTTCNCTGA	AGGCCCCCGG	600
ATNCNCTAGT	NCTAGAATCG	GCCCGCCATC	GCGGTGGANC	CTCCAACCTT	TCGTTNCCCT	660
TTACTGAGGG	TTNATTGCCG	CCCTTGCGCT	TATCATGGTC	ACNCCNGTTN	CCTGTGTTGA	720
AATTNTTAAC	CCCCACAAT	TCCACGCCNA	CATTNG			756

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 834 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

GGGGATCTCT	ANATCNACCT	GNATGCATGG	TTGTTCGGTGT	GGTCGCTGTC	GATGAANATG	60
AACAGGATCT	TGCCCTTGAA	GCTCTCGGCT	GCTGTNTTTA	AGTTGCTCAG	TCTGCCGTCA	120
TAGTCAGACA	CNCTCTTGGG	CAAAAAACAN	CAGGATNTGA	GTCTTGATTT	CACCTCCAAT	180
AATCTTCNGG	GCTGTCTGCT	CGGTGAACTC	GATGACNANG	GGCAGCTGGT	TGTGTNTGAT	240
AAANTCCANC	ANGTTCTCCT	TGGTGACCTC	CCCTTCAAAG	TTGTTCCGGC	CTTCATCAAA	300
CTTCTNNAAN	ANGANNANCC	CANCTTTGTC	GAGCTGGNAT	TTGGANAACA	CGTCACTGTT	360
GGAAACTGAT	CCCAAATGGT	ATGTCATCCA	TCGCCTCTGC	TGCCTGCAAA	AAACTTGCTT	420
GGCNCAAATC	CGACTCCCCN	TCCTTGAAAG	AAGCCNATCA	CACCCCCCTC	CCTGGACTCC	480
NNCAANGACT	CTNCCGCTNC	CCCNTPCCNG	CAGGGTTGGT	GGCANNCCGG	GCCCN TGCGC	540
TTCTTCAGCC	AGTTCACNAT	NTTCATCAGC	CCCTCTGCCA	GCTGTNTAT	TCCTTGGGGG	600
GGAANCCGTC	TCTCCCTTCC	TGAANNAACT	TTGACCGTNG	GAATAGCCGC	GCNTCNCCNT	660
ACNTNCTGGG	CCGGGTTCAA	ANTCCCTCCN	TTGNCNNTCN	CCTCGGGCCA	TTCTGGATTT	720
NCCNAACTTT	TTCCCTTCCCC	CNCCCCNCGG	NGTTTGGNTT	TTTCATNGGG	CCCCAACTCT	780
GCTNTTGGCC	ANTCCCCTGG	GGCNTNTAN	CNCCCCCTNT	GGTCCCNTNG	GGCC	834

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 814 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

CGGNCGCTTT	CCNGCCGCGC	CCCGTTTCCA	TGACNAAGGC	TCCCTTCANG	TTAAATACNN	60
CCTAGNAAAC	ATTAATGGGT	TGCTCTACTA	ATACATCATA	CNAACCAGTA	AGCCTGCCCA	120
NAACGCCAAC	TCAGGCCATT	CCTACCAAAG	GAAGAAAGGC	TGGTCTCTCC	ACCCCTGTGA	180
GGAAAGGCC	GCCTTGTAAG	ACACCACAA	NCGGCTGAAT	CTNAAGTCTT	GTGTTTACT	240
AATGGAAAAA	AAAAATAAAC	AANAGGTTTT	GTTCTCATGG	CTGCCACCG	CAGCCTGGCA	300
CTAAAACANC	CCAGCGCTCA	CTTCTGCTTG	GANAAATATT	CTTTGCTCTT	TTGGACATCA	360
GGCTTGATGG	TATCACTGCC	ACNTTTCAC	CCAGCTGGGC	NCCCTTCCCC	CATNTTGTG	420
ANTGANCTGG	AAGGCCTGAA	NCTTAGTCTC	CAAAAGTCTC	NGCCACAAAG	ACCGGCCACC	480
AGGGGANGTC	NTTINCAGTG	GATCTGCCAA	ANANTACCCN	TATCATCNNT	GAATAAAAAG	540
GCCCCTGAAC	GANATGCTTC	CANCANCTT	TAAGACCCAT	AATCCTNGAA	CCATGGTGCC	600
CTTCCGGTCT	GATCCNAAAG	GAATGTTTCT	GGGTCCCANT	CCCTCCTTTG	TTNCTTACGT	660
TGTNTTGGAC	CCNTGCTNGN	ATNACCCAAN	TGANATCCCC	NGAAGCACCC	TNCCCCTGGC	720
ATTTGANTTT	CNTAAATTCT	CTGCCCTACN	NCTGAAAGCA	CNATTCCCTN	GGCNCNAAN	780
GGNGAACTCA	AGAAGGTCTN	NGAAAAACCA	CNCN			814

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 760 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GCATGCTGCT	CTTCTCATA	GTTGTTCTTG	TTGCCATAAC	AACCACCATA	GGTAAAGCGG	60
GCGCAGTGTT	CGCTGAAGGG	GTTGTAGTAC	CAGCGCGGGA	TGCTCTCCTT	GCAGAGTCCT	120
GTGTCTGGCA	GGTCCACGCA	ATGCCCTTTG	TCACTGGGGA	AATGGATGCG	CTGGAGCTCG	180
TCNAANCCAC	TCGTGTATTT	TTACACANGCA	GCCTCCTCCG	AAGCNTCCGG	GCAGTTGGGG	240
GTGTCTGCAC	ACTCCACTAA	ACTGTGCGATN	CANCAGCCCA	TTGCTGCAGC	GGAAGTGGGT	300
GGGCTGACAG	GTGCCAGAAC	ACACTGGATN	GGCCTTTCCA	TGGAAGGGCC	TGGGGGAAAT	360
CNCCTNANCC	CAAACTGCCT	CTCAAAGGCC	ACCTTGACACA	CCCCGACAGG	CTAGAAATGC	420
ACTCTTCTTC	CCAAAGGTAG	TTGTTCTTGT	TGCCCAAGCA	NCCTCCANCA	AACCAAAAANC	480
TTGCAAAATC	TGCTCCGTGG	GGGTCATNMN	TACCANGGTT	GGGGAAANAA	ACCCGGCNGN	540
GANCCNCCTT	GTTTGAATGC	NAAGGNAATA	ATCCTCCTGT	CTTGCTTGGG	TGGAANAGCA	600
CAATTGAACT	GTTAACNTTG	GGCCGNGTTC	CNCTNGGGTG	GTCTGAAACT	AATCACCGTC	660
ACTGGAAAAA	GGTANGTGCC	TTCCTTGAAT	TCCCAAANTT	CCCCTNGNTT	TGGGTNNNTT	720
CTCCTCTNCC	CTAAAAATCG	TNTTCCCCCC	CCNTANGGCG			760

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 724 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTAAAAA	CCCCCTCCAT	TGAATGAAAA	60
------------	------------	------------	------------	------------	------------	----

CTTCCNAAAT	TGTCCAACCC	CCTCNNCCAA	ATNNCCATTT	CCGGGGGGGG	GTTCCAAACC	120
CAAATTAATT	TTGGANTTTA	AATTAAATNT	TNATTNGGGG	AANAANCCAA	ATGTNAAGAA	180
AATTTAACCC	ATTATNAACT	TAAATNCCTN	GAAACCCNTG	GNTTCCAAAA	ATTTTTAACC	240
CTTAAATCCC	TCCGAAATTG	NTAANGGAAA	ACCAAATTCN	CCTAAGGCTN	TTTGAAGGTT	300
NGATTTAAAC	CCCCTTNANT	TNTTTTNACC	CNNGNCTNAA	NTATTTNGNT	TCCGGTGTTT	360
TCCTNTTAAN	CNTNGGTAAC	TCCCGNTAAT	GAANNNCCCT	AANCCAATTA	AACCGAATTT	420
TTTTTGAATT	GGAAATTCCN	NGGGAATTNA	CCGGGGTTTT	TCCCNITTGG	GGGCCATNCC	480
CCCNCTTTCG	GGGTTTGGGN	NTAGGTTGAA	TTTTTNANG	NCCCCAAAAA	NCCCCAANA	540
AAAAAACTCC	CAAGNNTTAA	TTNGAATNTC	CCCTTCCCA	GGCCTTTTGG	GAAAGGNGGG	600
TTTNTGGGGG	CCNGGGANTT	CNTTCCCCCN	TTNCCNCCCC	CCCCCNGGT	AAANGGTTAT	660
NGNNTTTGGT	TTTGGGCCC	CTTNANGGAC	CTTCCGGATN	GAAATTAAAT	CCCCGGGNCG	720
GCCG						724

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 751 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

TTTTTTTTTT	TTTTCTTTG	CTCACATTTA	ATTTTTATTT	TGATTTTTTT	TAATGCTGCA	60
CAACACAATA	TTTATTTTCAT	TTGTTTCTTT	TATTTTCATTT	TATTTGTTTG	CTGCTGCTGT	120
TTTATTTATT	TTTACTGAAA	GTGAGAGGGA	ACTTTTGTGG	CCTTTTTTCC	TTTTTCTGTA	180
GGCCGCCTTA	AGCTTTCTAA	ATTTGGAACA	TCTAAGCAAG	CTGAANGGAA	AAGGGGGTTT	240
CGCAAAATCA	CTCGGGGGAA	NGGAAAGGTT	GCTTTGTAA	TCATGCCCTA	TGGTGGGTGA	300
TTAACTGCTT	GTACAATTAC	NTTTCACCTT	TAATTAATTG	TGCTNAANGC	TTTAATTANA	360
CTTGGGGGTT	CCCTCCCCAN	ACCAACCCCN	CTGACAAAAA	GTGCCNGCCC	TCAAATNATG	420
TCCCggcnnt	CNTTGAAACA	CACNGCNGAA	NGTTCTCATT	NTCCCCNCNC	CAGGTNAAAA	480
TGAAGGGTTA	CCATNTTTAA	CNCCACCTCC	ACNTGGCNNN	GCCTGAATCC	TCNAAAAANCN	540
CCCTCAANCN	AATTNCTNNG	CCCCGGTCNC	GCNTNNGTCC	CNCCCGGGCT	CCGGAANTN	600
CACCCCCNGA	ANNCNNTMNC	NAACNAAATT	CCGAAAATAT	TCCCNNTCNC	TCAATTCCCC	660
CNNAGACTNT	CCTCNCNCAN	CNCAATTTTC	TTTTNNTCAC	GAACNCGNNC	CNNAAAATGN	720
NNNNCNCCTC	CNCTNGTCCN	NAATCNCCAN	C			751

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 753 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GTGGTATTTT	CTGTAAGATC	AGGTGTTTCT	CCCTCGTAGG	TTTAGAGGAA	ACACCCTCAT	60
AGATGAAAAC	CCCCCGAGA	CAGCAGCACT	GCAACTGCCA	AGCAGCCGGG	GTAGGAGGGG	120
CGCCCTATGC	ACAGCTGGGC	CCTTGAGACA	GCAGGGCTTC	GATGTCAGGC	TCGATGTCAA	180
TGGTCTGGAA	GCGGCGGCTG	TACCTGCGTA	GGGGCACACC	GTCAGGGCCC	ACCAGGAAC	240
TCTCAAAGTT	CCAGGCAACN	TCGTTGCGAC	ACACCGGAGA	CCAGGTGATN	AGCTTGGGGT	300
CGGTCATAAN	CGCGGTGGCG	TCGTCGCTGG	GAGCTGGCAG	GGCCTCCCGC	AGGAAGGCNA	360
ATAAAAGGTG	CGCCCCCGCA	CCGTTCANCT	CGCACTTCTC	NAANACCATG	ANGTTGGGCT	420

CNAACCCACC	ACCANNCCGG	ACTTCCTTGA	NGGAATTCCC	AAATCTCTTC	GNTCTTGGGC	480
TTCTNCTGAT	GCCCTANCTG	GTTGCCCNNG	ATGCCAANCA	NCCCCAANCC	CCGGGGTCCT	540
AAANCACCCN	CCTCCTCNTT	TCATCTGGGT	TNTTNTCCCC	GGACCNTGGT	TCCTCTCAAG	600
GGANCCCATA	TCTCNACCAN	TACTCACCNT	NCCCCCCCNT	GNNACCCANC	CTTCTANNGN	660
TTCCCNCCCC	NCCTCTGGCC	CNTCAAANAN	GCTTNCACNA	CCTGGGTCTG	CCTTCCCCCC	720
TNCCCTATCT	GNACCCCNEN	TTTGTCTCAN	TNT			753

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 341 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

ACTATATCCA	TCACAACAGA	CATGCTTCAT	CCCATAGACT	TCTTGACATA	GCTTCAAATG	60
AGTGAACCCA	TCCTTGATTT	ATATACATAT	ATGTTCTCAG	TATTTTGGGA	GCCTTTCCAC	120
TTCTTTAAAC	CTTGTTTCAT	ATGAACACTG	AAAATAGGAA	TTTGTGAAGA	GTAAAAAAGT	180
TATAGCTTGT	TTACGTAGTA	AGTTTTTTGAA	GTCTACATTC	AATCCAGACA	CTTAGTTGAG	240
TGTTAAACTG	TGATTTTTTAA	AAAATATCAT	TTGAGAATAT	TCTTTCAGAG	GTATTTTCAT	300
TTTACTTTT	TGATTAATTG	TGTTTTATAT	ATTAGGGTAG	T		341

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 101 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

ACTTACTGAA	TTTAGTTCTG	TGCTCTTCCT	TATTTAGTGT	TGTATCATAA	ATACTTTGAT	60
GTTTCAAACA	TTCTAAATAA	ATAATTTTCA	GTGGCTTCAT	A		101

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 305 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Homo sapiens*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

ACATCTTTGT	TACAGTCTAA	GATGTGTTCT	TAAATCACCA	TTCCTTCCTG	GTCCTCACCC	60
TCCAGGGTGG	TCTCACACTG	TAATTAGAGC	TATTGAGGAG	TCTTTACAGC	AAATTAAGAT	120
TCAGATGCCCT	TGCTAAGTCT	AGAGTTCTAG	AGTTATGTTT	CAGAAAGTCT	AAGAAACCCA	180
CCTCTTGAGA	GGTCAGTAAA	GAGGACTTAA	TATTTTCATAT	CTACAAAATG	ACCACAGGAT	240
TGGATACAGA	ACGAGAGTTA	TCCTGGATAA	CTCAGAGCTG	AGTACCTGCC	CGGGGGCCGC	300
TCGAA						305

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 852 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Homo sapiens*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

ACATAAATAT	CAGAGAAAAG	TAGTCTTTGA	AATATTTACG	TCCAGGAGTT	CTTTGTTTCT	60
GATTATTTGG	TGTGTGTTTT	GGTTTGTGTC	CAAAGTATTG	GCAGCTTCAG	TTTTTCATTTT	120
CTCTCCATCC	TCGGGCATTC	TTCCCAAATT	TATATACCAG	TCTTCGTCCA	TCCACACGCT	180
CCAGAATTTT	TCTTTTGTAG	TAATATCTCA	TAGCTCGGCT	GAGCTTTTCA	TAGGTCATGC	240
TGCTGTTGTT	CTTCTTTTTA	CCCCATAGCT	GAGCCACTGC	CTCTGATTTT	AAGAACCTGA	300
AGACGCCCTC	AGATCGGTCT	TCCCATTTTA	TTAATCCTGG	GTTCTTGCTT	GGGTTCAAGA	360
GGATGTCGCG	GATGAATTCC	CATAAGTGAG	TCCCTCTCGG	GTTGTGCTTT	TTGGTGTGGC	420
ACTTGGCAGG	GGGGTCTTGC	TCCTTTTTCA	TATCAGGTGA	CTCTGCAACA	GGAAGGTGAC	480
TGGTGGTTGT	CATGGAGATC	TGAGCCCGGC	AGAAAGTTTT	GCTGTCCAAC	AAATCTACTG	540
TGCTACCATA	GTTGGTGTCA	TATAAATAGT	TCTNGTCTTT	CCAGGTGTTT	ATGATGGAAG	600
GCTCAGTTTG	TTCAGTCTTG	ACAATGACAT	TGTGTGTGGA	CTGGAACAGG	TCACTACTGC	660
ACTGGCCGTT	CCACTTCAGA	TGCTGCAAGT	TGCTGTAGAG	GAGNTGCCCC	GCCGTCCCCTG	720
CCGCCCCGGT	GAACTCCTGC	AAACTCATGC	TGCAAAGGTG	CTCGCCGTTG	ATGTCGAACT	780
CNTGGAAAGG	GATACAATTG	GCATCCAGCT	GGTTGGTGTG	CAGGAGGTGA	TGGAGCCACT	840
CCCACACCTG	GT					852

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 234 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Homo sapiens*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

ACAACAGACC	CTTGCTCGCT	AACGACCTCA	TGCTCATCAA	GTTGGACGAA	TCCGTGTCCG	60
------------	------------	------------	------------	------------	------------	----

AGTCTGACAC	CATCCGGAGC	ATCAGCATTG	CTTCGCAGTG	CCCTACCGCG	GGGAACTCTT	120
GCCTCGTTTC	TGGCTGGGGT	CTGCTGGCGA	ACGGCAGAAT	GCCTACCGTG	CTGCAGTGCG	180
TGAACGTGTC	GGTGGTGTCT	GAGGAGGTCT	GCAGTAAGCT	CTATGACCCG	CTGT	234

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 590 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

ACTTTTTATT	TAAATGTTTA	TAAGGCAGAT	CTATGAGAAT	GATAGAAAAC	ATGGTGTGTA	60
ATTTGATAGC	AATATTTTGG	AGATTACAGA	GTTTTAGTAA	TTACCAATTA	CACAGTTAAA	120
AAGAAGATAA	TATATTCCAA	GCANATACAA	AATATCTAAT	GAAAGATCAA	GGCAGGAAAA	180
TGANTATAAC	TAATTGACAA	TGGAAAATCA	ATTTTAATGT	GAATTGCACA	TTATCCTTTA	240
AAAGCTTTCA	AAANAAANAA	TTATTGCAGT	CTANTTAATT	CAAACAGTGT	TAAATGGTAT	300
CAGGATAAAN	AACTGAAGGG	CANAAAGAAT	TAATTTTCAC	TTCATGTAAC	NCACCCANAT	360
TTACAATGGC	TTAAATGCAN	GGAAAAAGCA	GTGGAAGTAG	GGAAGTANTC	AAGGTCTTTC	420
TGGTCTCTAA	TCTGCCTTAC	TCTTTGGGTG	TGGCTTTGAT	CCTCTGGAGA	CAGCTGCCAG	480
GGCTCCTGTT	ATATCCACAA	TCCCAGCAGC	AAGATGAAGG	GATGAAAAAG	GACACATGCT	540
GCCTTCCTTT	GAGGAGACTT	CATCTCACTG	GCCAACACTC	AGTCACATGT		590

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 774 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

ACAAGGGGGC	ATAATGAAGG	AGTGGGGANA	GATTTTAAAG	AAGGAAAAAA	AACGAGGCCC	60
TGAACAGAAT	TTTCCTGNAC	AACGGGGCTT	CAAAATAATT	TTCTTGGGGA	GGTTCAAGAC	120
GCTTCACTGC	TTGAAACTTA	AATGGATGTG	GGACANAATT	TTCTGTAATG	ACCCTGAGGG	180
CATTACAGAC	GGGACTCTGG	GAGGAAGGAT	AAACAGAAAG	GGGACAAAGG	CTAATCCCAA	240
AACATCAAAG	AAAGGAAGGT	GGCGTCATAC	CTCCCAGCCT	ACACAGTTCT	CCAGGGCTCT	300
CCTCATCCCT	GGAGGACGAC	AGTGGAGGAA	CAACTGACCA	TGTCCCCAGG	CTCCTGTGTG	360
CTGGCTCCTG	GTCTTCAGCC	CCCAGCTCTG	GAAGCCCACC	CTCTGCTGAT	CCTGCGTGGC	420
CCACACTCCT	TGAACACACA	TCCCCAGGTT	ATATTCTCTG	ACATGGCTGA	ACCTCCTATT	480
CCTACTTCCG	AGATGCCTTG	CTCCCTGCAG	CCTGTCAAAA	TCCCCTCAC	CCTCCAAACC	540
ACGGCATGGG	AAGCCTTTCT	GACTTGCCTG	ATTACTCCAG	CATCTTGGAA	CAATCCCTGA	600
TTCCCCACTC	CTTAGAGGCA	AGATAGGGTG	GTTAAGAGTA	GGGCTGGACC	ACTTGGAGCC	660
AGGCTGCTGG	CTTCAAATTN	TGGCTCATTT	ACGAGCTATG	GGACCTTGGG	CAAGTNATCT	720
TCATTCTAT	GGGCNTCATT	TTGTTCTACC	TGCAAAATGG	GGGATAATAA	TAGT	774

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 124 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

CANAAATTGA AATTTTATAA AAAGGCATTT TTCTCTTATA TCCATAAAAT GATATAATTT	60
TTGCAANTAT ANAAATGTGT CATAAATTAT AATGTTCCCTT AATTACAGCT CAACGCAACT	120
TGGT	124

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 147 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

GCCGATGCTA CTATTTTATT GCAGGAGGTG GGGGTGTTTT TATTATTCTC TCAACAGCTT	60
TGTGGCTACA GGTGGTGTCT GACTGCATNA AAAANTTTTT TACGGGTGAT TGCAAAAATT	120
TTAGGGCACC CATATCCCAA GCANTGT	147

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 107 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

ACATTAAATT AATAAAAGGA CTGTTGGGGT TCTGCTAAAA CACATGGCTT GATATATTGC	60
ATGTTTGTAG GTTAGGAGGA GTTAGGCATA TGTTTTGGGA GAGGGGT	107

(2) INFORMATION FOR SEQ ID NO:51:

863220" 9090E050

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

(2) INFORMATION FOR SEQ ID NO:52:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

ACAAAGATAA	CATTTATCTT	ATAACAAAA	TTTGATAGTT	TTAAAGGTTA	GTATTGTGTA	60
GGGTATTTTC	CAAAAGACTA	AAGAGATAAC	TCAGGTAAAA	AGTTAGAAAT	GTATAAAACA	120
CCATCAGACA	GGTTTTTTAA	AAACAACATA	TTACAAAATT	AGACAATCAT	CCTTAAAAAA	180
AAAAC TTCTT	GTATCAATTT	CTTTTGTTCA	AAATGACTGA	CTTAANTATT	TTTAAATATT	240
TCANAAACAC	TTCCTCAAAA	ATTTTCAANA	TGGTAGCTTT	CANATGTNCC	CTCAGTCCCA	300
ATGTTGCTCA	GATAAATAAA	TCTCGTGAGA	ACTTACCACC	CACCACAAGC	TTTCTGGGGC	360
ATGCAACAGT	GTCTTTTCTT	TNCTTTTCT	TTTTTTTTTT	TTACAGGCAC	AGAAACTCAT	420
CAATTTTATT	TGATAACAA	AGGGTCTCCA	AATTATATTG	AAAAATAAAT	CCAAGTTAAT	480
ATCACTCTTG	T					491

(2) INFORMATION FOR SEQ ID NO:53:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

ACATAATTTA	GCAGGGCTAA	TTACCATAAG	ATGCTATTTA	TTAANAGGTN	TATGATCTGA	60
GTATTAACAG	TTGCTGAAGT	TTGGTATTTT	TATGCAGCAT	TTTCTTTTGT	CTTTGATAAC	120
ACTACAGAAC	CCTTAAGGAC	ACTGAAAATT	AGTAAGTAAA	G TTCAGAAAC	ATTAGCTGCT	180
CAATCAAATC	TCTACATAAC	ACTATAGTAA	TTAAAAACGTT	AAAAAAAAAGT	GTTGAAATCT	240
GCACTAGTAT	ANACCGCTCC	TGTCAGGATA	ANACTGCTTT	GGAACAGAAA	GGGAAAAAANC	300
AGCTTTGANT	TTCTTTGTGC	TGATANGAGG	AAAGGCTGAA	TTACCTTGTT	GCCTCTCCCT	360
AATGATTGGC	AGGTCNGGTA	AATNCCAAAA	CATATTCCAA	CTCAACACTT	CTTTTCCNCG	420
TANCTTGANT	CTGTGTATTC	CAGGANCAGG	CGGATGGAAT	GGGCCAGCCC	NCGGATGTTC	480
CANT						484

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 151 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

ACTAAACCTC	GTGCTTGTTGA	ACTCCATACA	GAAAAACGGTG	CCATCCCTGA	ACACGGCTGG	60
CCACTGGGTA	TACTGCTGAC	AACCGCAACA	ACAAAAACAC	AAATCCTTGG	CACTGGCTAG	120
TCTATGTCCT	CTCAAGTGCC	TTTTTGTTTG	T			151

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 91 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

ACCTGGCTTG	TCTCCGGGTG	GTTCCCGGCG	CCCCCACGG	TCCCCAGAAC	GGACACTTTC	60
GCCCTCCAGT	GGATACTCGA	GCCAAAGTGG	T			91

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 133 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

GGCGGATGTG	CGTTGGTTAT	ATACAAATAT	GTCATTTTAT	GTAAGGGACT	TGAGTATACT	60
TGGATTTTGG	GTATCTGTGG	GTTGGGGGGA	CGGTCCAGGA	ACCAATACCC	CATGGGATACC	120
AAGGGACAAC	TGT					133

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 147 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

ACTCTGGAGA	ACCTGAGCCG	CTGCTCCGCC	TCTGGGATGA	GGTGATGCAN	GCNGTGGCGC	60
GACTGGGAGC	TGAGCCCTTC	CCTTTGCGCC	TGCCTCAGAG	GATTGTTGCC	GACNTGCANA	120
TCTCANTGGG	CTGGATNCAT	GCAGGGT				147

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 198 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

ACAGGGATAT	AGGTTTNAAG	TTATTGTNAT	TGTAAAATAC	ATTGAATTTT	CTGTATACTC	60
TGATTACATA	CATTTATCCT	TTAAAAAAGA	TGTAAATCTT	AATTTTTTATG	CCATCTATTA	120
ATTTACCAAT	GAGTTACCTT	GTAAATGAGA	AGTCATGATA	GCACTGAATT	TTAACTAGTT	180
TTGACTTCTA	AGTTTGGT					198

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 330 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20250920 14:50:00

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

ACAACAAATG	GGTTGTGAGG	AAGTCTTATC	AGCAAACTG	GTGATGGCTA	CTGAAAAGAT	60
CCATTGAAAA	TTATCATTA	TGATTTTAAA	TGACAAGTTA	TCAAAACTC	ACTCAATTTT	120
CACCTGTGCT	AGCTTGCTAA	AATGGGAGTT	AACTCTAGAG	CAAATATAGT	ATCTTCTGAA	180
TACAGTCAAT	AAATGACAAA	GCCAGGGCCT	ACAGGTGGTT	TCCAGACTTT	CCAGACCCAG	240
CAGAAGGAAT	CTATTTTATC	ACATGGATCT	CCGTCTGTGC	TCAAATACC	TAATGATAT	300
TTTCGTCTTT	ATTGGACTTC	TTTGAAGAGT				330

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 175 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

ACCGTGGGTG	CCTTCTACAT	TCCTGACGGC	TCCTTCACCA	ACATCTGGTT	CTACTTCGGC	60
GTCGTGGGCT	CCTTCTCTT	CATCCTCATC	CAGCTGGTGC	TGCTCATCGA	CTTTGCGCAC	120
TCCTGGAACC	AGCGGTGGCT	GGGCAAGGCC	GAGGAGTGCG	ATTCCCGTGC	CTGGT	175

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 154 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

ACCCCACTTT	TCCTCCTGTG	AGCAGTCTGG	ACTTCTCACT	GCTACATGAT	GAGGGTGAGT	60
GGTTGTTGCT	CTTCAACAGT	ATCCTCCCCT	TTCCGGATCT	GCTGAGCCGG	ACAGCAGTGC	120
TGGACTGCAC	AGCCCCGGGG	CTCCACATTG	CTGT			154

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

CGCTCGAGCC CTATAGTGAG TCGTATTAGA

30

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 89 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

ACAAGTCATT TCAGCACCCT TTGCTCTTCA AAAC TGACCA TCTTTTATAT TTAATGCTTC
CTGTATGAAT AAAAATGGTT ATGTCAAGT

60

89

(2) INFORMATION FOR SEQ ID NO:64:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 97 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

ACCGGAGTAA CTGAGTCGGG ACGCTGAATC TGAATCCACC AATAAATAAA GGTTCTGCAG
AATCAGTGCA TCCAGGATTG GTCCTTGGAT CTGGGGT

60

97

(2) INFORMATION FOR SEQ ID NO:65:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 377 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

66220 "90900000

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

ACAACAANAA	NTCCCTTCTT	TAGGCCACTG	ATGGAAACCT	GGAACCCCTT	TTTGATGGCA	60
GCATGGCGTC	CTAGGCCTTG	ACACAGCGGC	TGGGGTTTGG	GCTNTCCCAA	ACCGCACACC	120
CCAACCCTGG	TCTACCCACA	NTTCTGGCTA	TGGGCTGTCT	CTGCCACTGA	ACATCAGGGT	180
TCGGTCATAA	NATGAAATCC	CAANGGGGAC	AGAGGTCAGT	AGAGGAAGCT	CAATGAGAAA	240
GGTGCTGTTT	GCTCAGCCAG	AAAACAGCTG	CCTGGCATTG	GCCGCTGAAC	TATGAACCCG	300
TGGGGGTGAA	CTACCCCAN	GAGGAATCAT	GCCTGGGCGA	TGCAANGGTG	CCAACAGGAG	360
GGGCGGGAGG	AGCATGT					377

(2) INFORMATION FOR SEQ ID NO:66:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 305 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

ACGCCTTTCC	CTCAGAATTC	AGGGAAGAGA	CTGTCGCCTG	CCTTCCTCCG	TTGTTGCGTG	60
AGAACCCGTG	TGCCCCCTTC	CACCATATCC	ACCCTCGCTC	CATCTTTGAA	CTCAAACACG	120
AGGAACTAAC	TGCACCCTGG	TCCTCTCCCC	AGTCCCCAGT	TCACCCTCCA	TCCCTCACCT	180
TCCTCCACTC	TAAGGGATAT	CAACACTGCC	CAGCACAGGG	GCCCTGAATT	TATGTGGTTT	240
TTATATATTT	TTTAATAAGA	TGCACTTTAT	GTCATTTTTT	AATAAAGTCT	GAAGAATTAC	300
TGTTT						305

(2) INFORMATION FOR SEQ ID NO:67:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 385 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

ACTACACACA	CTCCACTTGC	CCTTGTGAGA	CACCTTGTCC	CAGCACTTTA	GGAATGCTGA	60
GGTCGGACCA	GCCACATCTC	ATGTGCAAGA	TTGCCCAGCA	GACATCAGGT	CTGAGAGTTC	120
CCCTTTTAAA	AAAGGGGACT	TGCTTAAAAA	AGAAGTCTAG	CCACGATTGT	GTAGAGCAGC	180
TGTGCTGTGC	TGGAGATTCA	CTTTTGTAGAG	AGTTCTCCTC	TGAGACCTGA	TCTTTAGAGG	240
CTGGGCAGTC	TTGCACATGA	GATGGGGCTG	GTCTGATCTC	AGCACTCCTT	AGTCTGCTTG	300
CCTCTCCAG	GGCCCCAGCC	TGGCCACACC	TGCTTACAGG	GCACTCTCAG	ATGCCCATAC	360
CATAGTTTCT	GTGCTAGTGG	ACCGT				385

(2) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

ACTTAACCAG ATATATTTTT ACCCCAGATG GGGATATTCT TTGTAAAAAA TGAAAATAAA	60
GTTTTTTTAA TGG	73

(2) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 536 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

ACTAGTCCAG TGTGGTGGAA TTCCATTGTG TTGGGGGCTC TCACCCTCCT CTCCTGCAGC	60
TCCAGCTTTG TGCTCTGCCT CTGAGGAGAC CATGGCCCAG CATCTGAGTA CCCTGCTGCT	120
CCTGCTGGCC ACCCTAGCTG TGGCCCTGGC CTGGAGCCCC AAGGAGGAGG ATAGGATAAT	180
CCCGGGTGGC ATCTATAACG CAGACCTCAA TGATGAGTGG GTACAGCGTG CCCTTCACTT	240
CGCCATCAGC GAGTATAACA AGGCCACCAA AGATGACTAC TACAGACGTC CGCTGCGGGT	300
ACTAAGAGCC AGGCAACAGA CCGTTGGGGG GGTGAATTAC TTCTTCGACG TAGAGGTGGG	360
CCGAACCATA TGTACCAAGT CCCAGCCCCA CTTGGACACC TGTGCCTTCC ATGAACAGCC	420
AGAAGTGCAG AAGAAACAGT TGTGCTCTTT CGAGATCTAC GAAGTTCCTT GGGGAGAACA	480
GAANGTCCCT GGGTGAAATC CAGGTGTCAA GAAATCCTAN GGATCTGTTG CCAGGC	536

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 477 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

ATGACCCCTA ACAGGGGCCC TCTCAGCCCT CCTAATGACC TCCGGCCTAG CCATGTGATT	60
TCACTTCCAC TCCATAACGC TCCTCATACT AGGCCTACTA ACCAACACAC TAACCATATA	120
CCAATGATGG CGCGATGTAA CACGAGAAAG CACATACCAA GGCCACCACA CACCACCTGT	180

865220 5098E060

CCAAAAAGGC	CTTCGATACG	GGATAATCCT	ATTTATTACC	TCAGAAAGTTT	TTTTCTTCGC	240
AGGGATTTTT	CTGAGCCTTT	TACCACTCCA	GCCTAGCCCC	TACCCCCCAA	CTAGGAGGGC	300
ACTGGCCCCC	AACAGGCATC	ACCCCGCTAA	ATCCCCTAGA	AGTCCCCTC	CTAAACACAT	360
CCGTATTACT	CGCATCAGGA	GTATCAATCA	CCTGAGCTCA	CCATAGTCTA	ATAGAAAAACA	420
ACCGAAACCA	AATTATTCAA	AGCACTGCTT	ATTACAATTT	TACTGGGTCT	CTATTTT	477

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 533 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

AGAGCTATAG	GTACAGTGTG	ATCTCAGCTT	TGCAAACACA	TTTTCTACAT	AGATAGTACT	60
AGGTATTAAT	AGATATGTAA	AGAAAGAAAT	CACACCATTA	ATAATGGTAA	GATTGGTTTTA	120
TGTGATTTTA	GTGGTATTTT	TGGCACCCCTT	ATATATGTTT	TCCAAACTTT	CAGCAGTGAT	180
ATTATTTTCCA	TAACTTAAAA	AGTGAGTTTG	AAAAAGAAAA	TCTCCAGCAA	GCATCTCATT	240
TAAATAAAGG	TTTGTCTATCT	TTAAAAATAC	AGCAATATGT	GACTTTTTTAA	AAAAGCTGTC	300
AAATAGGTGT	GACCCTACTA	ATAATTATTA	GAAATACATT	TAAAAACATC	GAGTACCTCA	360
AGTCAGTTTG	CCTTGAAAAA	TATCAAATAT	AACTCTTAGA	GAAATGTACA	TAAAAGAATG	420
CTTCGTAATT	TTGGAGTANG	AGGTTCCCTC	CTCAATTTTG	TATTTTTTAAA	AAGTACATGG	480
TAAAAAATAA	AATTCACAAC	AGTATATAAG	GCTGTAAAT	GAAGAATTCT	GCC	533

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 511 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

TATTACGGAA	AAACACACCA	CATAATTCAA	CTANCAAAGA	ANACTGCTTC	AGGGCGTGTA	60
AAATGAAAGG	CTTCCAGGCA	GTTATCTGAT	TAAAGAACAC	TAAAAGAGGG	ACAAGGCTAA	120
AAGCCGACGG	ATGCTACAC	TATANCAGGC	GCTATTTGGG	TTGGCTGGAG	GAGCTGTGGA	180
AAACATGGAN	AGATTGGTGC	TGGANATCGC	CGTGGCTATT	CCTCATTGTT	ATTACANAGT	240
GAGGTTCTCT	GTGTGCCCCAC	TGGTTTGAAA	ACCGTTCTNC	AATAATGATA	GAATAGTACA	300
CACATGAGAA	CTGAAATGGC	CCAAACCCAG	AAAGAAAGCC	CAACTAGATC	CTCAGAAANAC	360
GCTTCTAGGG	ACAATAACCG	ATGAAGAAAA	GATGGCCTCC	TTGTGCCCCC	GTCTGTTATG	420
ATTTCTCTCC	ATTGCAGCNA	NAAACCCGTT	CTTCTAAGCA	AACNCAGGTG	ATGATGGCNA	480
AAATACACCC	CCTCTTGAAG	NACCNGGAGG	A			511

(2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 499 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

CAGTGCCAGC	ACTGGTGCCA	GTACCACTAC	CAATAACAGT	GCCAGTGCCA	GTGCCAGCAC	60
CAGTGGTGGC	TTCAGTGCTG	GTGCCAGCCT	GACCGCCACT	CTCACATTTG	GGCTCTTCGC	120
TGGCCTTGGT	GGAGCTGGTG	CCAGCACCAG	TGGCAGCTCT	GGTGCCTGTG	GTTTCTCCTA	180
CAAGTGAGAT	TTTAGATATT	GTTAATCCTG	CCAGTCTTTC	TCTTCAAGCC	AGGGTGCATC	240
CTCAGAAACC	TACTCAACAC	AGCACTCTAG	GCAGCCACTA	TCAATCAATT	GAAGTTGACA	300
CTCTGCATTA	AATCTATTTG	CCATTTCTGA	AAAAAAAAAA	AAAAAAAAGG	CGGCCGCTCG	360
ANTCTAGAGG	GCCCGTTTAA	ACCCGCTGAT	CAGCCTCGAC	TGTGCCTTCT	ANTTGCCAGC	420
CATCTGTGTG	TTGCCCTCC	CCGNTGCCT	TCCTTGACCC	TGGAAGTGC	CACTCCCACT	480
GTCTTTTCCT	AANTAAAAAT					499

(2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 537 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

TTTCATAGGA	GAACACACTG	AGGAGATACT	TGAAGAATTT	GGATTTCAGCC	GCGAAGAGAT	60
TTATCAGCTT	AACTCAGATA	AAATCATTGA	AAGTAATAAG	GTAAAAGCTA	GTCTCTAACT	120
TCCAGGCCCA	CGGCTCAAGT	GAATTTGAAT	ACTGCATTTA	CAGTGTAGAG	TAACACATAA	180
CATTGTATGC	ATGGAAACAT	GGAGGAACAG	TATTACAGTG	TCCTACCACT	CTAATCAAGA	240
AAAGAATTAC	AGACTCTGAT	TCTACAGTGA	TGATTGAATT	CTAAAAATGG	TAATCATTAG	300
GGCTTTTGAT	TTATAANACT	TTGGGTACTT	ATACTAAATT	ATGGTAGTTA	TACTGCCTTC	360
CAGTTTGCTT	GATATATTTG	TTGATATTAA	GATTCTTGAC	TTATATTTTG	AATGGGTTCT	420
ACTGAAAAAN	GAATGATATA	TTCTTGAAGA	CATCGATATA	CATTTATTTA	CACTCTTGAT	480
TCTACAATGT	AGAAAATGAA	GGAAATGCCC	CAAATTGTAT	GGTGATAAAA	GTCCCGT	537

(2) INFORMATION FOR SEQ ID NO:75:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 467 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

255220" 20502050

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

CAAANACAAT TGTTCAAAAG ATGCAAATGA TACACTACTG CTGCAGCTCA CAAACACCTC	60
TGCATATTAC ACGTACCTCC TCCTGCTCCT CAAGTAGTGT GGTCTATTTT GCCATCATCA	120
CCTGCTGTCT GCTTAGAAGA ACGGCTTTCT GCTGCAANGG AGAGAAATCA TAACAGACGG	180
TGGCACAAGG AGGCCATCTT TTCCTCATCG GTTATTGTCC CTAGAAGCGT CTTCTGAGGA	240
TCTAGTTGGG CTTTCTTTCT GGGTTTGGGC CATTTTCANTT CTCATGTGTG TACTATTCTA	300
TCATTATTGT ATAACGGTTT TCAAACCNGT GGGCACNCAG AGAACCTCAC TCTGTAATAA	360
CAATGAGGAA TAGCCACGGT GATCTCCAGC ACCAAATCTC TCCATGTTNT TCCAGAGCTC	420
CTCCAGCCAA CCCAAATAGC CGCTGCTATN GTGTAGAACA TCCCTGN	467

(2) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 400 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

AAGCTGACAG CATTCTGGGCC GAGATGTCTC GCTCCGTGGC CTTAGCTGTG CTCGCGCTAC	60
TCTCTCTTTC TGGCCTGGAG GCTATCCAGC GTACTCCAAA GATTTCAGGT TACTCACGTC	120
ATCCAGCAGA GAATGGAAAG TCAAATTTCC TGAATTGCTA TGTGTCTGGG TTTCATCCAT	180
CCGACATTGA AGTTGACTTA CTGAAGAATG GAGAGAGAAT TGAAAAAGTG GAGCATTTCAG	240
ACTTGTCTTT CAGCAAGGAC TGGTCTTTCT ATCTCTTGTA CTACACTGAA TTCACCCCCA	300
CTGAAAAAGA TGAGTATGCC TGCCGTGTGA ACCATGTGAC TTTGTACAG CCCAAGATNG	360
TTNAGTGGGA TCGANACATG TAAGCAGCAN CATGGGAGGT	400

(2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 248 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

CTGGAGTGCC TTGGTGTTTC AAGCCCCTGC AGGAAGCAGA ATGCACCTTC TGAGGCACCT	60
CCAGCTGCCC CGGCGGGGGA TGCGAGGCTC GGAGCACCTC TGCCCGGCTG TGATTGCTGC	120
CAGGCACTGT TCATCTCAGC TTTTCTGTCC CTTTGCTCCC GGCAAGCGCT TCTGCTGAAA	180
GTTCATATCT GGAGCCTGAT GTCTTAACGA ATAAAGGTCC CATGCTCCAC CCGAAAAAAA	240
AAAAAAAA	248

(2) INFORMATION FOR SEQ ID NO:78:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 201 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

ACTAGTCCAG	TGTGGTGGAA	TTCCATTGTG	TTGGGCCCAA	CACAATGGCT	ACCTTTAACA	60
TCACCCAGAC	CCCGCCCTGC	CCGTGCCCCA	CGCTGCTGCT	AACGACAGTA	TGATGCTTAC	120
TCTGCTACTC	GGAACTATT	TTTATGTAAT	TAATGTATGC	TTTCTTGTTT	ATAAATGCCT	180
GATTTAAAAA	AAAAAAAAAA	A				201

(2) INFORMATION FOR SEQ ID NO:79:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 552 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

TCCTTTTGTT	AGGTTTTTGA	GACAACCCTA	GACCTAAACT	GTGTCACAGA	CTTCTGAATG	60
TTTAGGCAGT	GCTAGTAATT	TCCTCGTAAT	GATTCTGTTA	TTACTTTCCT	ATTCTTTATT	120
CCTCTTTCTT	CTGAAGATTA	ATGAAGTTGA	AAATTGAGGT	GGATAAATAC	AAAAAGGTAG	180
TGTGATAGTA	TAAGTATCTA	AGTGCAGATG	AAAGTGTGTT	ATATATATCC	ATTCAAAATT	240
ATGCAAGTTA	GTAATTACTC	AGGGTTAACT	AAATTACTTT	AATATGCTGT	TGAACCTACT	300
CTGTTCCCTG	GCTAGAAAAA	ATTATAAACA	GGACTTTGTT	AGTTTGGGAA	GCCAAATTGA	360
TAATATTCTA	TGTTCTAAAA	GTTGGGCTAT	ACATAAANTA	TNAAGAAATA	TGGAATTTTA	420
TTCCCAGGAA	TATGGGGTTC	ATTTATGAAT	ANTACCCGGG	ANAGAAAGTT	TGANTNAAAC	480
CNGTTTGGT	TAATACGTTA	ATATGTCCTN	AATNAACAAG	GCNTGACTTA	TTTCCAAAAA	540
AAAAAAAAAA	AA					552

(2) INFORMATION FOR SEQ ID NO:80:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 476 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

ACAGGGATTT	GAGATGCTAA	GGCCCCAGAG	ATCGTTTGAT	CCAACCCCTCT	TATTTTCAGA	60
GGGGAAAATG	GGGCCTAGAA	GTTACAGAGC	ATCTAGCTGG	TGCGCTGGCA	CCCCTGGCCT	120
CACACAGACT	CCCGAGTAGC	TGGGACTACA	GGCACACAGT	CACTGAAGCA	GGCCCTGTTT	180
GCAATTACAG	TTGCCACCTC	CAACTTAAAC	ATTCTTCATA	TGTGATGTCC	TTAGTCACTA	240
AGGTTAAACT	TTCCCACCCA	GAAAAGGCAA	CTTAGATAAA	ATCTTAGAGT	ACTTTCATAC	300
TCTTCTAAGT	CCTCTTCCAG	CCTCACTTTG	AGTCCTCCTT	GGGGGTTGAT	AGGAANTNTC	360
TCTTGGCTTT	CTCAATAAAA	TCTCTATCCA	TCTCATGTTT	AATTTGGTAC	GCNTAAAAAT	420
GCTGAAAAAA	TTAAATGTT	CTGGTTTCNC	TTTAAAAAAA	AAAAAAAAAA	AAAAAA	476

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 232 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

TTTTTTTTTG	TATGCCNTCN	CTGTGGNGTT	ATTGTTGCTG	CCACCCTGGA	GGAGCCCAGT	60
TTCTTCTGTA	TCTTTCTTTT	CTGGGGGATC	TTCTTGCTC	TGCCCCCTCA	TTCCCAGCCT	120
CTCATCCCCA	TCTTGCACTT	TTGCTAGGGT	TGGAGGCGCT	TTCTTGGTAG	CCCCTCAGAG	180
ACTCAGTCAG	CGGGAATAAG	TCCTAGGGGT	GGGGGGTGTT	GCAAGCCGGC	CT	232

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

AGGCGGGAGC	AGAAGCTAAA	GCCAAAGCCC	AAGAAGAGTG	GCAGTGCCAG	CACTGGTGCC	60
AGTACCAGTA	CCAATAACAT	GCCAGTGCCA	GTGCCAGCAC	CAGTGGTGCC	TTCACTGCTG	120
GTGCCAGCCT	GACCGCCACT	CTCACATTTG	GGCTCTTCGC	TGGCCTTGGT	GGAGCTGGTG	180
CCAGCACCAG	TGGCAGCTCT	GGTGCCGTGT	GTTTCTCCTA	CAAGTGAGAT	TTTAGATATT	240
GTTAATCCTG	CCAGTCTTTT	TCTTCAAGCC	AGGGTGCATC	CTCAGAAACC	TACTCAACAC	300
AGCACTCTNG	GCAGCCACTA	TCAATCAATT	GAAGTTGACA	CTCTGCATTA	AATCTATTTG	360
CCATTTCAAA	AAAAAAAAAA	AAA				383

(2) INFORMATION FOR SEQ ID NO:83:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 494 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

ACCGAATTGG	GACCGCTGGC	TTATAAGCGA	TCATGTCCTC	CAGTATTACC	TCAACGAGCA	60
GGGAGATCGA	GTCTATACGC	TGAAGAAATT	TGACCCGATG	GGACAACAGA	CCTGCTCAGC	120
CCATCCTGCT	CGGTTCTCCC	CAGATGACAA	ATACTCTCGA	CACCGAATCA	CCATCAAGAA	180
ACGCTTCAAG	GTGCTCATGA	CCCAGCAACC	GCGCCCTGTC	CTCTGAGGGT	CCTTAAACTG	240
ATGTCTTTTC	TGCCACCTGT	TACCCCTCGG	AGACTCCGTA	ACCAAACTCT	TCGGACTGTG	300
AGCCCTGATG	CCTTTTTTGGC	AGCCATACTC	TTTGGCNTCC	AGTCTCTCGT	GGCGATTGAT	360
TATGCTTGTG	TGAGGCAATC	ATGGTGGCAT	CACCCATNAA	GGGAACACAT	TTGANTTTTT	420
TTTCNCATAT	TTTAAATTAC	NACCAGAATA	NTTCAGAATA	AATGAATTGA	AAAACCTCTTA	480
AAAAAAAAAA	AAAA					494

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 380 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

GCTGGTAGCC	TATGGCGTGG	CCACGGANGG	GCTCCTGAGG	CACGGGACAG	TGACTTCCCA	60
AGTATCCTGC	GCCGCGTCTT	CTACCGTCCC	TACCTGCAGA	TCTTCGGGCA	GATTCCCCAG	120
GAGGACATGG	ACGTGGCCCT	CATGGAGCAC	AGCAACTGCT	CGTCGGAGCC	CGGCTTCTGG	180
GCACACCCTC	CTGGGGCCCA	GCGGGGCACC	TGCGTCTCCC	AGTATGCCAA	CTGGCTGGTG	240
GTGCTGCTCC	TCGTCATCTT	CCTGCTCGTG	GCCAACATCC	TGCTGGTCAC	TTGCTCATTG	300
CCATGTTCAG	TTACACATTC	GGCAAAGTAC	AGGGCAACAG	CNATCTCTAC	TGGGAAGGCC	360
AGCGTTNCCG	CCTCATCCGG					380

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 481 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

GAGTTAGCTC	CTCCACAACC	TTGATGAGGT	CGTCTGCAGT	GGCCTCTCGC	TTCATACCGC	60
TNCCATCGTC	ATACTGTAGG	TTTGCCACCA	CCTCCTGCAT	CTTGGGGCGG	CTAATATCCA	120
GGAAACTCTC	AATCAAGTCA	CCGTCNATNA	AACCTGTGGC	TGGTTCTGTC	TTCCGCTCGG	180
TGTGAAAGGA	TCTCCAGAAG	GAGTGCTCGA	TCTTCCCCAC	ACTTTTGATG	ACTTTATTGA	240
GTCGATTCTG	CATGTCCAGC	AGGAGGTTGT	ACCAGCTCTC	TGACAGTGAG	GTCACCAGCC	300
CTATCATGCC	NTTGAACGTG	CCGAAGAACA	CCGAGCCTTG	TGTGGGGGGT	GNAGTCTCAC	360
CCAGATTCTG	CATTACCAGA	NAGCCGTGGC	AAAAGANATT	GACAACTCGC	CCAGGNNGAA	420
AAAGAACACC	TCCTGGAAGT	GCTNGCCGCT	CCTCGTCCNT	TGGTGGNNGC	GCNTNCCTTT	480
T						481

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 472 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

AACATCTTCC	TGTATAATGC	TGTGTAATAT	CGATCCGATN	TTGTCTGCTG	AGAATTCATT	60
ACTTGGA AAA	GCAACTTNAA	GCCTGGACAC	TGGTATTAAA	ATTACACAATA	TGCAACACTT	120
TAAACAGTGT	GTCAATCTGC	TCCCTTACTT	TGTCATCACC	AGTCTGGGAA	TAAGGGTATG	180
CCCTATTAC	ACCTGT TAA	AGGGCGCTAA	GCATTTTTTGA	TTCAACATCT	TTTTTTTTTGA	240
CACAAGTCCG	AAAAAAGCAA	AAGTAAACAG	TTNTTAATTT	GTTAGCCAAT	TCACTTTCTT	300
CATGGGACAG	AGCCATTTGA	TTTAAAAAAGC	AAATTGCATA	ATATTGAGCT	TTGGGAGCTG	360
ATATNTGAGC	GGAAGANTAG	CCTTTCTACT	TCACCAGACA	CAACTCCTTT	CATATTGGGA	420
TGTTNACNAA	AGTTATGTCT	CTTACAGATG	GGATGCTTTT	GTGGCAATTC	TG	472

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 413 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

AGAAACCAGT	ATCTCTNAAA	ACAACCTCTC	ATACCTTG TG	GACCTAATTT	TGTGTGCGTG	60
TGTGTGTGCG	CGCATATTAT	ATAGACAGGC	ACATCTTTTT	TACTTTTGTA	AAAGCTTATG	120
CCTCTTTGGT	ATCTATATCT	GTGAAAGTTT	TAATGATCTG	CCATAATGTC	TTGGGGACCT	180
TTGTCTTCTG	TG TAAATGGT	ACTAGAGAAA	ACACCTATNT	TATGAGTCAA	TCTAGTTNGT	240
TTTATTTCGAC	ATGAAGGAAA	TTTCCAGATN	ACAACACTNA	CAAACTCTCC	CTTGACTAGG	300
GGGGACAAAAG	AAAAGCANAA	CTGAACATNA	GAAACAATTN	CCTGGTGAGA	AATTNCATAA	360
ACAGAAATTG	GGTNGTATAT	TGAAANANNG	CATCATTNAA	ACGTTTTTTT	TTT	413

(2) INFORMATION FOR SEQ ID NO:88:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 448 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

CGCAGCGGGT	CCTCTCTATC	TAGCTCCAGC	CTCTCGCCTG	CCCCACTCCC	CGCGTCCCGC	60
GTCCTAGCCN	ACCATGGCCG	GGCCCCCTGC	CGCCCCGCTG	CTCCTGCTGG	CCATCCTGGC	120
CGTGGCCCTG	GCCGTGAGCC	CCGCGGCCGG	CTCCAGTCCC	GGCAAGCCGC	CGCGCCTGGT	180
GGGAGGCCCA	TGGACCCCGC	GTGGAAGAAG	AAGGTGTGCG	GCGTGCACTG	GACTTTGCCG	240
TCGGCNANTA	CAACAAACCC	GCAACNACTT	TTACCNAGCN	CGCGCTGCAG	GTTGTGCCGC	300
CCCAANCAAA	TTGTTACTNG	GGGTAANTAA	TTCTTGGAAG	TTGAACCTGG	GCCAAACNNG	360
TTTACCAGAA	CCNAGCCAAT	TNGAACAAAT	NCCCCCTCCAT	AACAGCCCCT	T'TTAAAAAGG	420
GAANCANTCC	TGNTCTTTTC	CAAATTTT				448

(2) INFORMATION FOR SEQ ID NO:89:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 463 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

GAATTTTGTG	CACTGGCCAC	TGTGATGGAA	CCATTGGGCC	AGGATGCTTT	GAGTTTATCA	60
GTAGTGATTC	TGCCAAAGTT	GGTGTGTGTA	CATGAGTATG	TAAAATGTCA	AAAAATTAGC	120
AGAGGTCTAG	GTCTGCATAT	CAGCAGACAG	TTTGTCCGTG	TATTTTGTAG	CCTTGAAGTT	180
CTCAGTGACA	AGTTNNTTCT	GATGCGAAGT	TCTNATTCCA	GTGTTTTAGT	CCTTTGCATC	240
TTTNATGTTN	AGACTTGCCCT	CTNTNAAATT	GCTTTTGTNT	TCTGCAGGTA	CTATCTGTGG	300
TTTAACAAAA	TAGAANNACT	TCTCTGCTTN	GAANATTTGA	ATATCTTACA	TCTNAAAATN	360
AATTCTCTCC	CCATANNAAA	ACCCANGCCC	TTGGGANAAT	TTGAAAAANG	GNTCCTTCNN	420
AATTCNNANA	ANTTCAGNTN	TCATACAACA	NAACNGGANC	CCC		463

(2) INFORMATION FOR SEQ ID NO:90:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 400 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

AGGGATTGAA	GGTCTNTTNT	ACTGTCTGGAC	TGTTTCANCCA	CCAACTCTAC	AAGTTGCTGT	60
CTTCCACTCA	CTGTCTGTAA	GCNTNTTAAAC	CCAGACTGTA	TCTTCATAAA	TAGAACAAAT	120
TCTTCACCAG	TCACATCTTC	TAGGACCTTT	TTGGATTGAG	TTAGTATAAG	CTCTTCCACT	180
TCCTTTGTTA	AGACTTCATC	TGGTAAAGTC	TTAAGTTTTG	TAGAAAGGAA	TTTAATTGCT	240
CGTTCTCTAA	CAATGTCCTC	TCCTTGAAGT	ATTTGGCTGA	ACAACCCACC	TNAAGTCCCT	300
TTGTGCATCC	ATTTTAAATA	TACTTAATAG	GGCATTGGTN	CACTAGGTTA	AATTCTGCAA	360
GAGTCATCTG	TCTGCAAAAG	TTGCGTTAGT	ATATCTGCCA			400

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 480 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

GAGCTCGGAT	CCAATAATCT	TTGTCTGAGG	GCAGCACACA	TATNCAGTGC	CATGGNAACT	60
GGTCTACCCC	ACATGGGAGC	AGCATGCCGT	AGNTATATAA	GGTCATTCCC	TGAGTCAGAC	120
ATGCCTCTTT	GACTACCGTG	TGCCAGTGCT	GGTGATTCTC	ACACACCTCC	NMCCGCTCTT	180
TGTGGAAAAA	CTGGCACTTG	NCTGGAACCTA	GCAAGACATC	ACTTACAAAT	TCACCCACGA	240
GACACTTGAA	AGGTGTAACA	AAGCGACTCT	TGCATTGCTT	TTTGTCCCTC	CGGCACCACT	300
TGTCAATACT	AACCCGCTGG	TTTGCTCTCA	TCACATTTGT	GATCTGTAGC	TCTGGATACA	360
TCTCCTGACA	GTACTGAAGA	ACTTCTTCTT	TTGTTTCAAA	AGCAACTCTT	GGTGCCTGTT	420
NGATCAGGTT	CCCATTTCCT	AGTCCGAATG	TTCACATGGC	ATATNTTACT	TCCCACAAAA	480

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 477 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

ATACAGCCCA	NATCCCACCA	CGAAGATGCG	CTTGTTGACT	GAGAACCCTGA	TGCGGTCACT	60
GGTCCCGCTG	TAGCCCCAGC	GACTCTCCAC	CTGCTGGAAG	CGGTTGATGC	TGCACTCCTT	120
CCCACGCAGG	CAGCAGCGGG	GCCGGTCAAT	GAATCCCACT	CGTGGCTTGG	GGTTGACGGT	180
TAANTGCAGG	AAGAGGCTGA	CCACCTCGCG	GTCCACCAGG	ATGCCCGACT	GTGCGGGACC	240

TGCAGCGAAA	CTCCTCGATG	GTCATGAGCG	GGAAGCGAAT	GANGCCCAGG	GCCTTGCCCA	300
GAACCTTCCG	CCTGTTCTCT	GGCGTCACCT	GCAGCTGCTG	CCGCTNACAC	TCGGCCTCGG	360
ACCAGCGGAC	AAACGGCGTT	GAACAGCCGC	ACCTCACGGA	TGCCCANTGT	GTCGCGCTCC	420
AGGAACGGCN	CCAGCGTGTC	CAGGTCAATG	TCGGTGAANC	CTCCGCGGGT	AATGGCG	477

(2) INFORMATION FOR SEQ ID NO:93:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 377 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

GAACGGCTGG	ACCTTGCTC	GCATTGTGCT	GCTGGCAGGA	ATACCTTGGC	AAGCAGCTCC	60
AGTCCGAGCA	GCCCCAGACC	GCTGCCGCCC	GAAGCTAAGC	CTGCCTCTGG	CCTTCCCCTC	120
CGCCTCAATG	CAGAACCANT	AGTGGGAGCA	CTGTGTTTAG	AGTTAAGAGT	GAACACTGTN	180
TGATTTTACT	TGGGAATTTT	CTCTGTTATA	TAGCTTTTCC	CAATGCTAAT	TTCCAAACAA	240
CAACAACAAA	ATAACATGTT	TGCCTGTTNA	GTTGTATAAA	AGTANGTGAT	TCTGTATNTA	300
AAGAAAATAT	TACTGTTACA	TATACTGCTT	GCAANTTCTG	TATTTATTGG	TNCTCTGGAA	360
ATAAATATAT	TATTAAA					377

(2) INFORMATION FOR SEQ ID NO:94:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 495 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

CCCTTTGAGG	GGTTAGGGTC	CAGTTCCAG	TGGAAGAAAC	AGGCCAGGAG	AANTGCGTGC	60
CGAGCTGANG	CAGATTTCCC	ACAGTGACCC	CAGAGCCCTG	GGCTATAGTC	TCTGACCCCT	120
CCAAGGAAAG	ACCACCTTCT	GGGGACATGG	GCTGGAGGGC	AGGACCTAGA	GGCACCAAGG	180
GAAGGCCCCA	TTCCGGGGCT	GTTCCCCGAG	GAGGAAGGGA	AGGGGCTCTG	TGTGCCCCCC	240
ACGAGGAANA	GGCCCTGANT	CCTGGGATCA	NACACCCCTT	CACGTGTATC	CCCACACAAA	300
TGCAAGCTCA	CCAAGGTCCC	CTCTCAGTCC	CTTCCCTACA	CCCTGAACGG	NCACTGGCCC	360
ACACCCACCC	AGANCANCCA	CCCGCATGG	GGAATGTNCT	CAAGGAATCG	CNGGGCAACG	420
TGGACTCTNG	TCCCNNAAGG	GGGCAGAAATC	TCCAATAGAN	GGANNGAACC	CTTGCTNANA	480
AAAAAAAAANA	AAAAA					495

(2) INFORMATION FOR SEQ ID NO:95:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 472 base pairs
 - (B) TYPE: nucleic acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

GGTTACTTGG	TTTCATTGCC	ACCACTTAGT	GGATGTCATT	TAGAACCATT	TTGTCTGCTC	60
CCTCTGGAAG	CCTTGCGCAG	AGCGGACTTT	GTAATTGTTG	GAGAATAACT	GCTGAATTTT	120
TAGCTGTTTT	GAGTTGATTC	GCACCACTGC	ACCACAACCTC	AATATGAAAA	CTATTTNACT	180
TATTTATTAT	CTTGTGAAAA	GTATACAATG	AAAATTTTGT	TCATACTGTA	TTTATCAAGT	240
ATGATGAAAA	GCAATAGATA	TATATTCTTT	TATTATGTTN	AATTATGATT	GCCATTATTA	300
ATCGGCAAAA	TGTGGAGTGT	ATGTTCTTTT	CACAGTAATA	TATGCCTTTT	GTAACCTTAC	360
TTGGTTATTT	TATTGTAAAT	GAATTACAAA	ATTCTTAATT	TAAGAAAATG	GTANGTTATA	420
TTTANTTCAN	TAATTTCTTT	CCTTGTTTAC	GTTAATTTTG	AAAAGAATGC	AT	472

(2) INFORMATION FOR SEQ ID NO:96:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 476 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

CTGAAGCATT	TCTTCAAAC	TNTCTACTTT	TGTCATTGAT	ACCTGTAGTA	AGTTGACAAT	60
GTGGTGAAAT	TTCAAAATTA	TATGTAACCT	CTACTAGTTT	TACTTTCTCC	CCCAAGTCTT	120
TTTTAACTCA	TGATTTTAC	ACACACAATC	CAGAACTTAT	TATATAGCCT	CTAAGTCTTT	180
ATTCTTCACA	GATAGATG	AAAGAGTCCT	CCAGTGTCTT	GNGCANAATG	TTCTAGNTAT	240
AGCTGGATAC	ATACNGTGGG	AGTTCTATAA	ACTCATACCT	CAGTGGGACT	NAACCAAAAT	300
TGTGTTAGTC	TCAATTCCTA	CCACACTGAG	GGAGCCTCCC	AAATCACTAT	ATTCTTATCT	360
GCAGGTACTC	CTCCAGAAAA	ACNGACAGGG	CAGGCTTGCA	TGAAAAAGTN	ACATCTGCGT	420
TACAAAGTCT	ATCTTCTCTA	NANGTCTGTN	AAGGAACAAT	TTAATCTTCT	AGCTTT	476

(2) INFORMATION FOR SEQ ID NO:97:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 479 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

ACTCTTTCTA	ATGCTGATAT	GATCTTGAGT	ATAAGAATGC	ATATGTCACT	AGAATGGATA	60
AAATAATGCT	GCAAACTTAA	TGTTCTTATG	CAAAATGGAA	CGCTAATGAA	ACACAGCTTA	120
CAATCGCAAA	TCAAAACTCA	CAAGTGCTCA	TCTGTTGTAG	ATTTAGTGTA	ATAAGACTTA	180
GATTGTGCTC	CTTCGGATAT	GATTGTTTCT	CANATCTTGG	GCAATNTTCC	TTAGTCAAAT	240
CAGGCTACTA	GAATTCTGTT	ATTGGATATN	TGAGAGCATG	AAATTTTTTAA	NAATACACTT	300
GTGATTATNA	AATTAATCAC	AAATTTCACT	TATACCTGCT	ATCAGCAGCT	AGAAAAACAT	360
NTNNTTTTTA	NATCAAAGTA	TTTTGTGTTT	GGAANTGTNN	AAATGAAATC	TGAATGTGGG	420
TTCNATCTTA	TTTTTTCCCN	GACNACTANT	TNCTTTTTTA	GNGNCTATTG	TGANCCATC	479

(2) INFORMATION FOR SEQ ID NO:98:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 461 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

AGTGACTTGT	CCTCCAACAA	AACCCCTTGA	TCAAGTTTGT	GGCACTGACA	ATCAGACCTA	60
TGCTAGTTCC	TGTCATCTAT	TCGCTACTAA	ATGCAGACTG	GAGGGGACCA	AAAAGGGGCA	120
TCAACTCCAG	CTGGATTATT	TTGGAGCCTG	CAAATCTATT	CCTACTTGTA	CGGACTTTGA	180
AGTGATTTCAG	TTTCCTCTAC	GGATGAGAGA	CTGGCTCAAG	AATATCCTCA	TGCAGCTTTA	240
TGAAGCCACT	CTGAACACGC	TGGTTATCTA	GATGAGAACA	GAGAAATAAA	GTCAGAAAAT	300
TTACCTGGAG	AAAAGAGGCT	TTGGCTGGGG	ACCATCCCAT	TGAACCTTCT	CTTAAGGACT	360
TTAAGAAAAA	CTACCACATG	TTGTGTATCC	TGGTGCCGGC	CGTTTATGAA	CTGACCACCC	420
TTTGGAATAA	TCTTGACGCT	CCTGAACCTG	CTCCTCTGCG	A		461

(2) INFORMATION FOR SEQ ID NO:99:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 171 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

GTGGCCGCGC	GCAGGTGTTT	CCTCGTACCG	CAGGGCCCCC	TCCCTTCCCC	AGGCGTCCCT	60
CGGCGCCTCT	GCGGGCCCGA	GGAGGAGCGG	CTGGCGGGTG	GGGGGAGTGT	GACCCACCCT	120
CGGTGAGAAA	AGCCTTCTCT	AGCGATCTGA	GAGGCGTGCC	TTGGGGGTAC	C	171

(2) INFORMATION FOR SEQ ID NO:100:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 269 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

20250520 10:50:50

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

CGGCCGCAAG TGCAACTCCA GCTGGGGCCG TGC GGACGAA GATTCTGCCA GCAGTTGGTC	60
CGACTGCGAC GACGGCGGCG GCGACAGTCG CAGGTGCAGC GCGGGCGCCT GGGGTCTTGC	120
AAGGCTGAGC TGACGCCGCA GAGGTCGTGT CACGTCCCAC GACCTTGACG CCGTCGGGGA	180
CAGCCGGAAC AGAGCCCGGT GAAGCGGGAG GCCTCGGGGA GCCCTCGGG AAGGGCGGCC	240
CGAGAGATAC GCAGGTGCAG GTGGCCGCC	269

(2) INFORMATION FOR SEQ ID NO:101:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 405 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

TTTTTTTTTT TTTTGGAATC TACTGCGAGC ACAGCAGGTC AGCAACAAGT TTATTTTGCA	60
GCTAGCAAGG TAACAGGGTA GGGCATGTTT ACATGTTTTC GTCAACTTCC TTTGTCGTGG	120
TTGATTGGTT TGTCTTTATG GGGGCGGGGT GGGGTAGGGG AAACGAAGCA AATAACATGG	180
AGTGGGTGCA CCCTCCCTGT AGAACCTGGT TACAAAGCTT GGGGCAGTTC ACCTGGTCTG	240
TGACCGTCAT TTTCTTGACA TCAATGTTAT TAGAAGTCAG GATATCTTTT AGAGAGTCCA	300
CTGTTCTGGA GGGAGATTAG GGTTCCTTGC CAAATCCAAC AAAATCCACT GAAAAAGTTG	360
GATGATCAGT ACGAATACCG AGGCATATTC TCATATCGGT GGCCA	405

(2) INFORMATION FOR SEQ ID NO:102:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 470 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

TTTTTTTTTT TTTTTTTTTT TTTTTTTTTT TTTTTTTTTT TTTTTTTTTT TTTTTTTTTT	60
GGCACTTAAT CCATTTTAT TTCAAATGT CTACAAATTT AATCCCATTA TACGGTATTT	120
TCAAAATCTA AATTATTCAA ATTAGCCAAA TCCTTACCAA ATAATACCCA AAAATCAAAA	180
ATATACTTCT TTCAGCAAAC TTGTTACATA AATTAATAAA ATATATACGG CTGGTGTTTT	240
CAAAGTACAA TTATCTTAAC ACTGCAAACA TTTTAAGGAA CTAAATAAAA AAAAAACACT	300

CCGCAAAGGT TAAAGGGAAC AACAAATTCT TTTACAACAC CATTATAAAA ATCATATCTC	360
AAATCTTAGG GGAATATATA CTTACACGG GATCTTAACT TTTACTCACT TTGTTTATTT	420
TTTTAAACCA TTGTTTGGGC CCAACACAAT GGAATCCCC CTGGACTAGT	470

(2) INFORMATION FOR SEQ ID NO:103:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 581 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

TTTTTTTTTT TTTTTTTTGA CCCCCCTCTT ATAAAAACA AGTTACCATT TTATTTTACT	60
TACACATATT TATTTTATAA TTGGTATTAG ATATTCAAAA GGCAGCTTTT AAAATCAAAC	120
TAAATGGAAA CTGCCTTAGA TACATAATTC TTAGGAATTA GCTTAAAAATC TGCCATAAAGT	180
GAAAATCTTC TCTAGCTCTT TTGACTGTAA ATTTTGTGACT CTTGTAAAAC ATCCAAATTC	240
ATTTTCTTG TCTTTAAAAAT TATCTAATCT TTCCATTTTT TCCCTATTCC AAGTCAATTT	300
GCTTCTCTAG CCTCATTTC TAGCTCTTAT CTACTATTAG TAAGTGGCTT TTTTCCTAAA	360
AGGGAAAACA GGAAGAGAAA TGGCACACAA AACAAACATT TTATATTTCAT ATTTCTACCT	420
ACGTTAATAA AATAGCATTT TGTGAAGCCA GCTCAAAAGA AGGCTTAGAT CCTTTTATGT	480
CCATTTTAGT CACTAAACGA TATCAAAGTG CCAGAATGCA AAAGGTTTGT GAACATTTAT	540
TCAAAAGCTA ATATAAGATA TTTACATAC TCATCTTTCT G	581

(2) INFORMATION FOR SEQ ID NO:104:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 578 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

TTTTTTTTTT TTTTTTTTTT TTTTCTCTT CTTTTTTTTT GAAATGAGGA TCGAGTTTTT	60
CACTCTCTAG ATAGGGCATG AAGAAAACTC ATCTTTCCAG CTTTAAAAATA ACAATCAAAT	120
CTCTTATGCT ATATCATATT TTAAGTTAAA CTAATGAGTC ACTGGCTTAT CTTCTCCTGA	180
AGGAAATCTG TTCATTCTTC TCATTATAT AGTTATATCA AGTACTACCT TGCATATTGA	240
GAGGTTTTTC TTCTCTATTT ACACATATAT TTCCATGTGA ATTTGTATCA AACCTTTATT	300
TTCATGCAAA CTAGAAAATA ATGTTTCTTT TGCATAAGAG AAGAGAACAA TATAGCATT	360
CAAACTGCT CAAATTGTTT GTTAAGTTAT CCATTATAAT TAGTTGGCAG GAGCTAATAC	420
AAATCACATT TACGACAGCA ATAATAAAC TGAAGTACCA GTTAAATATC CAAAATAATT	480
AAAGGAACAT TTTTAGCCTG GGTATAATTA GCTAATTCAC TTTACAAGCA TTTATTAGAA	540
TGAATTCACA TGTTATTATT CCTAGCCCCA CACAATGG	578

(2) INFORMATION FOR SEQ ID NO:105:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 538 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

TTTTTTTTTTT	TTTTTCAGTA	ATAATCAGAA	CAATATTTAT	TTTTATATT	AAAATTCATA	60
GAAAAGTGCC	TTACATTTAA	TAAAAGTTTG	TTTCTCAAAG	TGATCAGAGG	AATTAGATAT	120
GTCTTGAACA	CCAATATTAA	TTTGAGGAAA	ATACACCAAA	ATACATTAAG	TAAATTATTT	180
AAGATCATAG	AGCTTGTAAG	TGAAAAGATA	AAATTTGACC	TCAGAAACTC	TGAGCATTAA	240
AAATCCACTA	TTAGCAAATA	AATTACTATG	GACTTCTTGC	TTTAATTTTG	TGATGAATAT	300
GGGGTGTCAC	TGGTAAACCA	ACACATTCTG	AAGGATACAT	TACTTAGTGA	TAGATTCTTA	360
TGTACTTTGC	TAATACGTGG	ATATGAGTTG	ACAAGTTTCT	CTTTCTTCAA	TCTTTTAAGG	420
GGCGAGAAAT	GAGGAAGAAA	AGAAAAGGAT	TACGCATACT	GTTCTTTCTA	TGGAAGGATT	480
AGATATGTTT	CCTTTGCCAA	TATTAAAAAA	ATAATAATGT	TTACTACTAG	TGAAACCC	538

(2) INFORMATION FOR SEQ ID NO:106:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 473 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

TTTTTTTTTTT	TTTTTTAGTC	AAGTTTCTAT	TTTTATTATA	ATTAAAGTCT	TGGTCATTTT	60
ATTTATTAGC	TCTGCAACTT	ACATATTTAA	ATTAAAGAAA	CGTTTTAGAC	AACTGTACAA	120
TTTATAAATG	TAAGGTGCCA	TTATTGAGTA	ATATATTCCT	CCAAGAGTGG	ATGTGTCCCT	180
TCTCCCACCA	ACTAATGAAC	AGCAACATTA	GTTTAATTTT	ATTAGTAGAT	ATACACTGCT	240
GCAAACGCTA	ATTCTCTTCT	CCATCCCCAT	GTGATATTGT	GTATATGTGT	GAGTTGGTAG	300
AATGCATCAC	AATCTACAAT	CAACAGCAAG	ATGAAGCTAG	GCTGGGCTTT	CGGTGAAAAT	360
AGACTGTGTC	TGTCTGAATC	AAATGATCTG	ACCTATCCTC	GGTGGCAAGA	ACTCTTCGAA	420
CCGCTTCCTC	AAAGGCGCTG	CCACATTTGT	GGCTCTTTGC	ACTTGTTTCA	AAA	473

(2) INFORMATION FOR SEQ ID NO:107:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1621 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

CGCCATGGCA CTGCAGGGCA TCTCGGTCAT GGAGCTGTCC GGCCTGGCCC CGGGCCCGTT 60
 CTGTGCTATG GTCCTGGCTG ACTTCGGGGC GCGTGTGGTA CGCGTGGACC GGCCCGGCTC 120
 CCGCTACGAC GTGAGCCGCT TGGGCCGGGG CAAGCGCTCG CTAGTGCTGG ACCTGAAGCA 180
 GCCGCGGGGA GCCGCCGTGC TGCGGCGTCT GTGCAAGCGG TCGGATGTGC TGCTGGAGCC 240
 CTTCCGCCGC GGTGTCATGG AGAAACTCCA ACTGGGCCCA GAGATTCTGC AGCGGGAAAA 300
 TCCAAGGCTT ATTTATGCCA GGCTGAGTGG ATTTGGCCAG TCAGGAAGCT TCTGCCGGTT 360
 AGCTGGCCAC GATATCAACT ATTTGGCTTT GTGAGGTGTT CTCTCAAAAA TTGGCAGAAG 420
 TGGTGAGAAT CCGTATGCCC CGCTGAATCT CCTGGCTGAC TTTGCTGGTG GTGGCCTTAT 480
 GTGTGCACTG GGCATTATAA TGGCTCTTTT TGACCGCACA CGCACTGACA AGGGTCAGGT 540
 CATTGATGCA AATATGGTGG AAGGAACAGC ATATTTAAGT TCTTTTCTGT GGAAACTCA 600
 GAAATCGAGT CTGTGGGAAG CACCTCGAGG ACAGAACATG TTGGATGGTG GAGCACCTTT 660
 CTATACGACT TACAGGACAG CAGATGGGGA ATTCATGGCT GTTGGAGCAA TAGAACCCCA 720
 GTTCTACGAG CTGCTGATCA AAGGACTTGG ACTAAAGTCT GATGAACCTC CCAATCAGAT 780
 GAGCATGGAT GATTGGCCAG AAATGAAGAA GAAGTTTGCA GATGTATTTG CAAAGAAGAC 840
 GAAGGCAGAG TGGTGTCAA TCTTTGACGG CACAGATGCC TGTGTGACTC CGGTTCTGAC 900
 TTTTGAGGAG GTTGTTTCATC ATGATCACAA CAAGGAACGG GGCTCGTTTA TCACCAGTGA 960
 GGAGCAGGAC GTGAGCCCCC GCCCTGCACC TCTGCTGTTA AACACCCAG CCATCCCTTC 1020
 TTTCAAAGG GATCCTTTCA TAGGAGAACA CACTGAGGAG ATACTTGAAG AATTTGGATT 1080
 CAGCCGCGAA GAGATTTATC AGCTTAACTC AGATAAAATC ATTGAAAGTA ATAAGGTAAA 1140
 AGCTAGTCTC TAACTTCCAG GCCCAGCGCT CAAGTGAATT TGAATACTGC ATTTACAGTG 1200
 TAGAGTAACA CATAACATTG TATGCATGGA AACATGGAGG AACAGTATTA CAGTGTCCTA 1260
 CCACTCTAAT CAAGAAAAGA ATTACAGACT CTGATTCTAC AGTGATGATT GAATTCTAAA 1320
 AATGGTTATC ATTAGGGCTT TTGATTTATA AAACTTTGGG TACTTATACT AAATTATGGT 1380
 AGTTATTCTG CCTTCCAGTT TGCTTGATAT ATTTGTTGAT ATTAAGATTC TTGACTTATA 1440
 TTTTGAATGG GTTCTAGTGA AAAAGGAATG ATATATTCTT GAAGACATCG ATATACATTT 1500
 ATTTACACTC TTGATTCTAC AATGTAGAAA ATGAGGAAAT GCCACAAAT GTATGGTGAT 1560
 AAAAGTCACG TGAAACAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 1620
 A 1621

(2) INFORMATION FOR SEQ ID NO:108:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 382 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

Met Ala Leu Gln Gly Ile Ser Val Met Glu Leu Ser Gly Leu Ala Pro
 1 5 10 15
 Gly Pro Phe Cys Ala Met Val Leu Ala Asp Phe Gly Ala Arg Val Val
 20 25 30
 Arg Val Asp Arg Pro Gly Ser Arg Tyr Asp Val Ser Arg Leu Gly Arg
 35 40 45
 Gly Lys Arg Ser Leu Val Leu Asp Leu Lys Gln Pro Arg Gly Ala Ala
 50 55 60
 Val Leu Arg Arg Leu Cys Lys Arg Ser Asp Val Leu Leu Glu Pro Phe
 65 70 75 80

Arg Arg Gly Val Met Glu Lys Leu Gln Leu Gly Pro Glu Ile Leu Gln
 85 90 95
 Arg Glu Asn Pro Arg Leu Ile Tyr Ala Arg Leu Ser Gly Phe Gly Gln
 100 105 110
 Ser Gly Ser Phe Cys Arg Leu Ala Gly His Asp Ile Asn Tyr Leu Ala
 115 120 125
 Leu Ser Gly Val Leu Ser Lys Ile Gly Arg Ser Gly Glu Asn Pro Tyr
 130 135 140
 Ala Pro Leu Asn Leu Leu Ala Asp Phe Ala Gly Gly Gly Leu Met Cys
 145 150 155 160
 Ala Leu Gly Ile Ile Met Ala Leu Phe Asp Arg Thr Arg Thr Asp Lys
 165 170 175
 Gly Gln Val Ile Asp Ala Asn Met Val Glu Gly Thr Ala Tyr Leu Ser
 180 185 190
 Ser Phe Leu Trp Lys Thr Gln Lys Ser Ser Leu Trp Glu Ala Pro Arg
 195 200 205
 Gly Gln Asn Met Leu Asp Gly Gly Ala Pro Phe Tyr Thr Thr Tyr Arg
 210 215 220
 Thr Ala Asp Gly Glu Phe Met Ala Val Gly Ala Ile Glu Pro Gln Phe
 225 230 235 240
 Tyr Glu Leu Leu Ile Lys Gly Leu Gly Leu Lys Ser Asp Glu Leu Pro
 245 250 255
 Asn Gln Met Ser Met Asp Asp Trp Pro Glu Met Lys Lys Lys Phe Ala
 260 265 270
 Asp Val Phe Ala Lys Lys Thr Lys Ala Glu Trp Cys Gln Ile Phe Asp
 275 280 285
 Gly Thr Asp Ala Cys Val Thr Pro Val Leu Thr Phe Glu Glu Val Val
 290 295 300
 His His Asp His Asn Lys Glu Arg Gly Ser Phe Ile Thr Ser Glu Glu
 305 310 315 320
 Gln Asp Val Ser Pro Arg Pro Ala Pro Leu Leu Asn Thr Pro Ala
 325 330 335
 Ile Pro Ser Phe Lys Arg Asp Pro Phe Ile Gly Glu His Thr Glu Glu
 340 345 350
 Ile Leu Glu Glu Phe Gly Phe Ser Arg Glu Glu Ile Tyr Gln Leu Asn
 355 360 365
 Ser Asp Lys Ile Ile Glu Ser Asn Lys Val Lys Ala Ser Leu
 370 375 380

(2) INFORMATION FOR SEQ ID NO:109:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1524 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

GGCACGAGGC	TGCGCCAGGG	CCTGAGCGGA	GGCGGGGGCA	GCCTCGCCAG	CGGGGGCCCC	60
GGGCCTGGCC	ATGCCTCACT	GAGCCAGCGC	CTGCGCCTCT	ACCTCGCCGA	CAGCTGGAAC	120
CAGTGCGACC	TAGTGGCTCT	CACCTGCTTC	CTCCTGGGCG	TGGGCTGCCG	GCTGACCCCG	180
GGTTTGTACC	ACCTGGGCCG	CACCTGTCCTC	TGCATCGACT	TCATGGTTTT	CACGGTGCGG	240

CTGCTTCACA	TCTTCACGGT	CAACAAACAG	CTGGGGCCCA	AGATCGTCAT	CGTGAGCAAG	300
ATGATGAAGG	ACGTGTTCTT	CTTCCTCTTC	TTCCTCGGCG	TGTGGCTGGT	AGCCTATGGC	360
GTGGCCACGG	AGGGGCTCCT	GAGGCCACGG	GACAGTGA	TCCCAAGTAT	CCTGCGCCGC	420
GTCTTCTACC	GTCCCTACCT	GCAGATCTTC	GGGCAGATTC	CCCAGGAGGA	CATGGACGTG	480
GCCCTCATGG	AGCACAGCAA	CTGCTCGTCG	GAGCCCCGGT	TCTGGGCACA	CCCTCCTGGG	540
GCCCAGGCGG	GCACCTGCGT	CTCCCAGTAT	GCCAACTGGC	TGGTGGTGT	GCTCCTCGTC	600
ATCTTCTCTG	TCGTGGCCAA	CATCCTGCTG	GTCAACTTGC	TCATTGCCAT	GTTCAGTTAC	660
ACATTCGGCA	AAGTACAGGG	CAACAGCGAT	CTCTACTGGA	AGGCGCAGCG	TTACCGCCTC	720
ATCCGGGAAT	TCCACTCTCG	GCCCCGCGCT	GCCCCGCCCT	TTATCGTCAT	CTCCCCTTG	780
CGCCTCCTGC	TCAGGCAATT	GTGCAGGCGA	CCCCGGAGCC	CCCAGCCGTC	CTCCCCGGCC	840
CTCGAGCATT	TCCGGGTTTA	CCTTTCTAAG	GAAGCCGAGC	GGAAGCTGCT	AACGTGGGAA	900
TCGGTGCATA	AGGAGAACTT	TCTGCTGGCA	CGCGCTAGGG	ACAAGCGGGA	GAGCGACTCC	960
GAGCGTCTGA	AGCGCACGTC	CCAGAAGGTG	GACTTGGCAC	TGAAACAGCT	GGGACACATC	1020
CGCGAGTACG	AACAGCGCCT	GAAAGTGCTG	GAGCGGGAGG	TCCAGCAGTG	TAGCCGCGTC	1080
CTGGGGTGGG	TGGCCGAGGC	CCTGAGCCGC	TCTGCCTTGG	TGCCCCCAGG	TGGGCCGCCA	1140
CCCCCTGACC	TGCCCTGGTC	CAAAGACTGA	GCCCTGCTGG	CGGACTTCAA	GGAGAAGCCC	1200
CCACAGGGGA	TTTTGTCTCT	AGAGTAAGGC	TCATCTGGGC	CTCGGCCCCC	GCACCTGGTG	1260
GCCTTGTCCT	TGAGGTGAGC	CCCATGTCCA	TCTGGGCCAC	TGTCAGGACC	ACCTTTGGGA	1320
GTGTCATCCT	TACAAACCAC	AGCATGCCCG	GCTCCTCCCA	GAACCAGTCC	CAGCCTGGGA	1380
GGATCAAGGC	CTGGATCCCG	GGCCGTTATC	CATCTGGAGG	CTGCAGGGTC	CTTGGGGTAA	1440
CAGGGACCAC	AGACCCCTCA	CCACTCACAG	ATTCTCTACA	CTGGGGAAAT	AAAGCCATTT	1500
CAGAGGAAAA	AAAAAAAAAA	AAAA				1524

(2) INFORMATION FOR SEQ ID NO:110:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3410 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

GGGAACCAGC	CTGCACGCGC	TGGCTCCGGG	TGACAGCCGC	GCGCCTCGGC	CAGGATCTGA	60
GTGATGAGAC	GTGTCCCCAC	TGAGGTGCCC	CACAGCAGCA	GGTGTGAGC	ATGGGCTGAG	120
AAGCTGGACC	GGCACCAAAG	GGCTGGCAGA	AATGGGCGCC	TGGCTGATTC	CTAGGCAGTT	180
GGCGGCAGCA	AGGAGGAGAG	GCCGCAGCTT	CTGGAGCAGA	GCCGAGACGA	AGCAGTTCTG	240
GAGTGCCCTGA	ACGGCCCCCT	GAGCCCTACC	CGCCTGGCCC	ACTATGGTCC	AGAGGCTGTG	300
GGTGAGCCGC	CTGCTGCGGC	ACCGGAAAGC	CCAGCTCTTG	CTGGTCAACC	TGCTAACCTT	360
TGGCCTGGAG	GTGTGTTTGG	CCGCAGGCAT	CACCTATGTG	CCGCCTCTGC	TGCTGGAAGT	420
GGGGGTAGAG	GAGAAAGTTCA	TGACCATGGT	GCTGGGCATT	GGTCCAGTGC	TGGGCCTGGT	480
CTGTGTCCCG	CTCCTAGGCT	CAGCCAGTGA	CCACTGGCGT	GGACGCTATG	GCCGCCGCCG	540
GCCCTTCATC	TGGGCACTGT	CCTTGGGCAT	CCTGCTGAGC	CTCTTTCTCA	TCCCAAGGGC	600
CGGCTGGCTA	GCAGGGCTGT	TGTGCCCGGA	TCCCAGGCCC	CTGGAGCTGG	CACTGCTCAT	660
CCTGGGCGTG	GGGCTAGCTG	ACTTCTGTGG	CCAGGTGTGC	TTCACTCCAC	TGGAGGCCCT	720
GCTCTCTGAC	CTCTTCCGGG	ACCCGGACCA	CTGTGCGCCAG	GCCTACTCTG	TCTATGCCCTT	780
CATGATCAGT	CTTGGGGGCT	GCCTGGGCTA	CCTCCTGCCT	GCCATTGACT	GGGACACCAG	840
TGCCCTGGCC	CCCTACCTGG	GCACCCAGGA	GGAGTGCCCT	TTTGGCCTGC	TCACCCTCAT	900
CTTCCTCACC	TGCCGTAGCAG	CCACACTGCT	GGTGGCTGAG	GAGGCAGCGC	TGGGCCCCAC	960
CGAGCCAGCA	GAAGGGCTGT	CGGCCCCCTC	CTTGTCGCCC	CACTGCTGTC	CATGCCGGGC	1020
CCGCTTGGCT	TTCCGGAACC	TGGGCGCCCT	GCTTCCCCGG	CTGCACCAGC	TGTGCTGCCG	1080
CATGCCCCGC	ACCCTGCGCC	GGCTCTTCGT	GGCTGAGCTG	TGCAGCTGGA	TGGCACTCAT	1140
GACCTTCACG	CTGTTTACAA	CGGATTTTCGT	GGGCGAGGGG	CTGTACCAGG	GCGTGCCCG	1200

AGCTGAGCCG GGCACCGAGG CCCGGAGACA CTATGATGAA GGCCTTCGGA TGGGCAGCCT 1260
GGGGCTGTTC CTGCAGTGC GGCATCTCCCT GGTCTTCTCT CTGGTCATGG ACCGGCTGGT 1320
GCAGCGATTG GGCAC'CTGAG CAGTCTATTT GGCCAGTGTG GCAGCTTTCC CTGTGGCTGC 1380
CGGTGCCACA TGCCTGTCCC ACAGTGTGGC CGTGGTGACA GCTTCAGCCG CCCTCACC GG 1440
GTTACACCTT TCAGCCCTGC AGATCCTGCC CTACACACTG GCCTCCCTCT ACCACCGGGA 1500
GAAGCAGGTG TTCCTGCCCC AATACCGAGG GGACACTGGA GGTGCTAGCA GTGAGGACAG 1560
CCTGATGACC AGCTTCCTGC CAGGCCCTAA GCCTGGAGCT CCCTTCCCTA ATGGACACGT 1620
GGGTGCTGGA GGCAGTGGCC TGCTCCCACC TCCACCCGCG CTCTGCGGGG CCTCTGCCTG 1680
TGATGTCTCC GTACGTGTGG TGGTGGGTGA GCCCAGGAG GCCAGGGTGG TTCCGGGCCG 1740
GGGCATCTGC CTGGACCTCG CCATCCTGGA TAGTGCTTTC CTGCTGTCCC AGGTGGCCCC 1800
ATCCCTGTTT ATGGGCTCCA TTGTCCAGCT CAGCCAGTCT GTCAGTGCCT ATATGGTGTG 1860
TGCCGACAGC CTGGGTCTGG TCGCCATTTA CTTTGCTACA CAGGTAGTAT TTGACAAGAG 1920
CGACTTGGCC AAATACTCAG CGTAGAAAAC TTCCAGCACA TTGGGGTGGA GGGCTGCCT 1980
CACTGGGTCC CAGCTCCCCG CTCCTGTTAG CCCCATGGGG CTGCCGGGCT GGCCGCCAGT 2040
TTCTGTGTGCT GCCAAAGTAA TGTGGCTCTC TGCTGCCACC CTGTGCTGCT GAGGTGCGTA 2100
GCTGCACAGC TGGGGGCTGG GGCGTCCCTC TCCTCTCTCC CCAGTCTCTA GGGCTGCCTG 2160
ACTGGAGGCC TTCCAAGGGG GTTTCAGTCT GGACTTATAC AGGGAGGCCA GAAGGGCTCC 2220
ATGCACTGGA ATGCGGGGAC TCTGCAGGTG GATTACCCAG GCTCAGGGTT AACAGCTAGC 2280
CTCCTAGTTG AGACACACCT AGAGAAGGGT TTTTGGGAGC TGAATAAACT CAGTCACCTG 2340
GTTTCCCATC TCTAAGCCCC TTAACCTGCA GCTTCGTTTA ATGTAGCTCT TGCATGGGAG 2400
TTTCTAGGAT GAAACACTCC TCCATGGGAT TTGAACATAT GACTTATTTG TAGGGGAAGA 2460
GTCCTGAGGG GCAACACACA AGAACCAGGT CCCCTCAGCC CACAGCACTG TCTTTTGTCT 2520
GATCCACCCC CCTCTTACCT TTTATCAGGA TGTGGCCTGT TGGTCCTTCT GTTGCCATCA 2580
CAGAGACACA GGCATTTAAA TATTTAACTT ATTTATTTAA CAAAGTAGAA GGGAATCCAT 2640
TGCTAGCTTT TCTGTGTTGG TGTCTAATAT TTGGGTAGGG TGGGGGATCC CCAACAATCA 2700
GGTCCCCTGA GATAGCTGGT CATTGGGCTG ATCATTGCCA GAATCTTCTT CTCCTGGGGT 2760
CTGGCCCCCC AAAATGCCTA ACCCAGGACC TTGGAAATTC TACTCATCCC AAATGATAAT 2820
TCCAAATGCT GTTACCCAAG GTTAGGGTGT TGAAGGAAGG TAGAGGGTGG GGCTTCAGGT 2880
CTCAACGGCT TCCCTAACCA CCCCTCTTCT CTTGGCCAG CCTGGTTCCC CCCACTTCCA 2940
CTCCCCTCTA CTCTCTCTAG GACTGGGCTG ATGAAGGCAC TGCCCAAAAT TTCCCCTACC 3000
CCCAACTTTC CCCTACCCCC AACTTTCCCC ACCAGCTCCA CAACCCTGTT TGGAGCTACT 3060
GCAGGACCAG AAGCACAAG TGCGGTTTCC CAAGCCTTTC TCCATCTCAG CCCCAGAGT 3120
ATATCTGTGC TTGGGGAATC TCACACAGAA ACTCAGGAGC ACCCCTGCC TGAGCTAAGG 3180
GAGGTCTTAT CTCTCAGGGG GGGTTTAACT GCGTTTGCA ATAATGTCGT CTTATTTATT 3240
TAGCGGGGTG AATATTTTAT ACTGTAAGTG AGCAATCAGA GTATAATGTT TATGGTGACA 3300
AAATTAAAGG CTTTCTTATA TGTTTAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 3360
AAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAATAA AAAAAAAAAA 3410

(2) INFORMATION FOR SEQ ID NO:111:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1289 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:111:

AGCCAGGCGT CCCTCTGCCT GCCCACTCAG TGGCAACACC CGGGAGCTGT TTTGTCCTTT 60
GTGGAGCCTC AGCAGTTCCT TCTTTCAGAA CTCACTGCCA AGAGCCCTGA ACAGGAGCCA 120
CCATGCAGTG CTTCACTTC ATTAAGACCA TGATGATCCT CTTCAATTTG CTCATCTTTC 180
TGTGTGGTGC AGCCCTGTTG GCAGTGGGCA TCTGGGTGTC AATCGATGGG GCATCCTTTC 240
TGAAGATCTT CGGGCCACTG TCGTCCAGTG CCATGCAGTT TGTCACCGTG GGCTACTTCC 300

```

TCATCGCAGC CGGCGTTGTG GTCTTTGCTC TTGGTTTCCT GGGCTGCTAT GGTGCTAAGA 360
CTGAGAGCAA GTGTGCCCTC GTGACGTTCT TCTTCATCCT CCTCCTCATC TTCATTGCTG 420
AGGTTGCAGC TGCTGTGGTC GCCTTGGTGT ACACCACAAT GGCTGAGCAC TTCCTGACGT 480
TGCTGGTAGT GCCTGCCATC AAGAAAGATT ATGGTTCCCA GGAAGACTTC ACTCAAGTGT 540
GGAACACCAC CATGAAAGGG CTCAAGTGCT GTGGCTTCAC CAACTATACG GATTTTGAGG 600
ACTCACCCTA CTTCAAAGAG AACAGTGCCT TTCCCCCATT CTGTTGCAAT GACAACGTCA 660
CCAACACAGC CAATGAAACC TGCACCAAGC AAAAGGCTCA CGACCAAAAA GTAGAGGGTT 720
GCTTCAATCA GCTTTTGTAT GACATCCGAA CTAATGCAGT CACCGTGGGT GGTGTGGCAG 780
CTGGAATTGG GGGCCTCGAG CTGGCTGCCA TGATTGTGTC CATGTATCTG TACTGCAATC 840
TACAATAAGT CCACTTCTGC CTCTGCCACT ACTGCTGCCA CATGGGAACT GTGAAGAGGC 900
ACCTGGCAA GCAGCAGTGA TTGGGGGAGG GGACAGGATC TAACAATGTC ACTTGGGCCA 960
GAATGGACCT GCCCTTCTG CTCCAGACTT GGGGCTAGAT AGGGACCACT CCTTTTAGCG 1020
ATGCCTGACT TTCCTTCCAT TGGTGGGTGG ATGGGTGGGG GGCATTCCAG AGCCTCTAAG 1080
GTAGCCAGTT CTGTTGCCCA TTCCCCCAGT CTATTAAACC CTTGATATGC CCCCTAGGCC 1140
TAGTGGTGAT CCCAGTGCTC TACTGGGGGA TGAGAGAAAG GCATTTTATA GCCTGGGCAT 1200
AAGTGAATC AGCAGAGCCT CTGGGTGGAT GTGTAGAAGG CACTTCAAAA TGCATAAACC 1260
TGTTACAATG TTAAAAAAA AAAAAAAA 1289

```

(2) INFORMATION FOR SEQ ID NO:112:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 315 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

```

Met Val Phe Thr Val Arg Leu Leu His Ile Phe Thr Val Asn Lys Gln
1           5           10           15

Leu Gly Pro Lys Ile Val Ile Val Ser Lys Met Met Lys Asp Val Phe
20           25           30

Phe Phe Leu Phe Phe Leu Gly Val Trp Leu Val Ala Tyr Gly Val Ala
35           40           45

Thr Glu Gly Leu Leu Arg Pro Arg Asp Ser Asp Phe Pro Ser Ile Leu
50           55           60

Arg Arg Val Phe Tyr Arg Pro Tyr Leu Gln Ile Phe Gly Gln Ile Pro
65           70           75           80

Gln Glu Asp Met Asp Val Ala Leu Met Glu His Ser Asn Cys Ser Ser
85           90           95

Glu Pro Gly Phe Trp Ala His Pro Pro Gly Ala Gln Ala Gly Thr Cys
100          105          110

Val Ser Gln Tyr Ala Asn Trp Leu Val Val Leu Leu Leu Val Ile Phe
115          120          125

Leu Leu Val Ala Asn Ile Leu Leu Val Asn Leu Leu Ile Ala Met Phe

```

130	135	140
Ser Tyr Thr Phe Gly Lys Val Gln Gly Asn Ser Asp Leu Tyr Trp Lys		
145	150	155 160
Ala Gln Arg Tyr Arg Leu Ile Arg Glu Phe His Ser Arg Pro Ala Leu		
	165	170 175
Ala Pro Pro Phe Ile Val Ile Ser His Leu Arg Leu Leu Leu Arg Gln		
	180	185 190
Leu Cys Arg Arg Pro Arg Ser Pro Gln Pro Ser Ser Pro Ala Leu Glu		
	195	200 205
His Phe Arg Val Tyr Leu Ser Lys Glu Ala Glu Arg Lys Leu Leu Thr		
	210	215 220
Trp Glu Ser Val His Lys Glu Asn Phe Leu Leu Ala Arg Ala Arg Asp		
	225	230 235 240
Lys Arg Glu Ser Asp Ser Glu Arg Leu Lys Arg Thr Ser Gln Lys Val		
	245	250 255
Asp Leu Ala Leu Lys Gln Leu Gly His Ile Arg Glu Tyr Glu Gln Arg		
	260	265 270
Leu Lys Val Leu Glu Arg Glu Val Gln Gln Cys Ser Arg Val Leu Gly		
	275	280 285
Trp Val Ala Glu Ala Leu Ser Arg Ser Ala Leu Leu Pro Pro Gly Gly		
	290	295 300 305
Pro Pro Pro Pro Asp Leu Pro Gly Ser Lys Asp		
	305	310 315

(2) INFORMATION FOR SEQ ID NO:113:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 553 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

Met Val Gln Arg Leu Trp Val Ser Arg Leu Leu Arg His Arg Lys Ala
1 5 10 15
Gln Leu Leu Leu Val Asn Leu Leu Thr Phe Gly Leu Glu Val Cys Leu
20 25 30
Ala Ala Gly Ile Thr Tyr Val Pro Pro Leu Leu Leu Glu Val Gly Val
35 40 45

Glu 50	Lys	Phe	Met	Thr	Met	Val	Leu	Gly	Ile	Gly 60	Pro	Val	Leu	Gly	
Leu 65	Val	Cys	Val	Pro	Leu 70	Leu	Gly	Ser	Ala	Ser 75	Asp	His	Trp	Arg 80	
Arg	Tyr	Gly	Arg	Arg 85	Arg	Pro	Phe	Ile	Trp 90	Ala	Leu	Ser	Leu	Gly 95	
Leu	Leu	Ser	Leu 100	Phe	Leu	Ile	Pro	Arg 105	Ala	Gly	Trp	Leu	Ala 110	Gly	Leu
Leu	Cys	Pro 115	Asp	Pro	Arg	Pro	Leu 120	Glu	Leu	Ala	Leu	Leu 125	Ile	Leu	Gly
Val	Gly 130	Leu	Leu	Asp	Phe	Cys 135	Gly	Gln	Val	Cys	Phe 140	Thr	Pro	Leu	Glu
Ala 145	Leu	Leu	Ser	Asp	Leu 150	Phe	Arg	Asp	Pro	Asp 155	His	Cys	Arg	Gln	Ala 160
Tyr	Ser	Val	Tyr	Ala 165	Phe	Met	Ile	Ser	Leu 170	Gly	Gly	Cys	Leu	Gly 175	Tyr
Leu	Leu	Pro 180	Ala	Ile	Asp	Trp	Asp 185	Thr	Ser	Ala	Leu	Ala 190	Pro	Tyr	Leu
Gly	Thr	Gln 195	Glu	Glu	Cys	Leu	Phe 200	Gly	Leu	Leu	Thr	Leu 205	Ile	Phe	Leu
Thr 210	Cys	Val	Ala	Ala	Thr	Leu 215	Leu	Val	Ala	Glu	Glu 220	Ala	Ala	Leu	Gly
Pro 225	Thr	Glu	Pro	Ala	Glu 230	Gly	Leu	Ser	Ala	Pro 235	Ser	Leu	Ser	Pro	His 240
Cys	Cys	Pro	Cys	Arg 245	Ala	Arg	Leu	Ala	Phe 250	Arg	Asn	Leu	Gly	Ala 255	Leu
Leu	Pro	Arg	Leu 260	His	Gln	Leu	Cys 265	Cys	Arg	Met	Pro	Arg	Thr 270	Leu	Arg
Arg	Leu	Phe 275	Val	Ala	Glu	Leu	Cys 280	Ser	Trp	Met	Ala	Leu 285	Met	Thr	Phe
Thr 290	Leu	Phe	Tyr	Thr	Asp	Phe 295	Val	Gly	Glu	Gly	Leu 300	Tyr	Gln	Gly	Val
Pro 305	Arg	Ala	Glu	Pro	Gly 310	Thr	Glu	Ala	Arg	Arg 315	His	Tyr	Asp	Glu	Gly 320
Val	Arg	Met	Gly	Ser 325	Leu	Gly	Leu	Phe	Leu 330	Gln	Cys	Ala	Ile	Ser	Leu
Val	Phe	Ser	Leu 340	Val	Met	Asp	Arg	Leu 345	Val	Gln	Arg	Phe	Gly 350	Thr	Arg

[illegible]

Ala Val Tyr Leu Ala Ser Val Ala Ala Phe Pro Val Ala Ala Gly Ala
 355 360 365

Thr Cys Leu Ser His Ser Val Ala Val Val Thr Ala Ser Ala Ala Leu
 370 375 380

Thr Gly Phe Thr Phe Ser Ala Leu Gln Ile Leu Pro Tyr Thr Leu Ala
 385 390 395 400

Ser Leu Tyr His Arg Glu Lys Gln Val Phe Leu Pro Lys Tyr Arg Gly
 405 410 415

Asp Thr Gly Gly Ala Ser Ser Glu Asp Ser Leu Met Thr Ser Phe Leu
 420 425 430

Pro Gly Pro Lys Pro Gly Ala Pro Phe Pro Asn Gly His Val Gly Ala
 435 440 445

Gly Gly Ser Gly Leu Leu Pro Pro Pro Pro Ala Leu Cys Gly Ala Ser
 450 455 460

Ala Cys Asp Val Ser Val Arg Val Val Val Gly Glu Pro Thr Glu Ala
 465 470 475 480

Arg Val Val Pro Gly Arg Gly Ile Cys Leu Asp Leu Ala Ile Leu Asp
 485 490 495

Ser Ala Phe Leu Leu Ser Gln Val Ala Pro Ser Leu Phe Met Gly Ser
 500 505 510

Ile Val Gln Leu Ser Gln Ser Val Thr Ala Tyr Met Val Ser Ala Ala
 515 520 525

Gly Leu Gly Leu Val Ala Ile Tyr Phe Ala Thr Gln Val Val Phe Asp
 530 535 540

Lys Ser Asp Leu Ala Lys Tyr Ser Ala
 545 550

(2) INFORMATION FOR SEQ ID NO:114:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 241 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

Met Gln Cys Phe Ser Phe Ile Lys Thr Met Met Ile Leu Phe Asn Leu
 1 5 10 15

Leu Ile Phe Leu Cys Gly Ala Ala Leu Leu Ala Val Gly Ile Trp Val

20					25					30					
Ser	Ile	Asp	Gly	Ala	Ser	Phe	Leu	Lys	Ile	Phe	Gly	Pro	Leu	Ser	Ser
		35					40					45			
Ser	Ala	Met	Gln	Phe	Val	Asn	Val	Gly	Tyr	Phe	Leu	Ile	Ala	Ala	Gly
		50					55					60			
Val	Val	Val	Phe	Ala	Leu	Gly	Phe	Leu	Gly	Cys	Tyr	Gly	Ala	Lys	Thr
		65					70					75			
Glu	Ser	Lys	Cys	Ala	Leu	Val	Thr	Phe	Phe	Phe	Ile	Leu	Leu	Leu	Ile
				85					90					95	
Phe	Ile	Ala	Glu	Val	Ala	Ala	Ala	Val	Val	Ala	Leu	Val	Tyr	Thr	Thr
				100					105					110	
Met	Ala	Glu	His	Phe	Leu	Thr	Leu	Leu	Val	Val	Pro	Ala	Ile	Lys	Lys
				115					120					125	
Asp	Tyr	Gly	Ser	Gln	Glu	Asp	Phe	Thr	Gln	Val	Trp	Asn	Thr	Thr	Met
		130							135					140	
Lys	Gly	Leu	Lys	Cys	Cys	Gly	Phe	Thr	Asn	Tyr	Thr	Asp	Phe	Glu	Asp
		145							150					155	
Ser	Pro	Tyr	Phe	Lys	Glu	Asn	Ser	Ala	Phe	Pro	Pro	Phe	Cys	Cys	Asn
				165					170					175	
Asp	Asn	Val	Thr	Asn	Thr	Ala	Asn	Glu	Thr	Cys	Thr	Lys	Gln	Lys	Ala
				180					185					190	
His	Asp	Gln	Lys	Val	Glu	Gly	Cys	Phe	Asn	Gln	Leu	Leu	Tyr	Asp	Ile
				195					200					205	
Arg	Thr	Asn	Ala	Val	Thr	Val	Gly	Gly	Val	Ala	Ala	Gly	Ile	Gly	Gly
				210					215					220	
Leu	Glu	Leu	Ala	Ala	Met	Ile	Val	Ser	Met	Tyr	Leu	Tyr	Cys	Asn	Leu
				225					230					235	
														240	
Gln															

(2) INFORMATION FOR SEQ ID NO:115:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 366 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

GCTCTTTCTC	TCCCCTCCTC	TGAATTTAAT	TCTTTCAACT	TGCAATTTGC	AAGGATTACA	60
CATTTCACTG	TGATGTATAT	TGTGTTGCAA	AAAAAAAAAA	GTGTCTTTGT	TTAAAATTAC	120
TTGGTTTGTG	AATCCATCTT	GCTTTTTCCC	CATTGGAACT	AGTCATTAAC	CCATCTCTGA	180
ACTGGTAGAA	AAACATCTGA	AGAGCTAGTC	TATCAGCATC	TGACAGGTGA	ATTGGATGGT	240
TCTCAGAAC	ATTTACCCA	GACAGCCTGT	TTCTATCCTG	TTAATAAAT	TAGTTTGGGT	300
TCTCTACATG	CATAACAAAC	CCTGCTCCAA	TCTGTACAT	AAAAGTCTGT	GACTTGAAGT	360
TTAGTC						366

(2) INFORMATION FOR SEQ ID NO:116:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 282 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

ACAAAGATGA	ACCATTTCTT	ATATTATAGC	AAAATTAAAA	TCTACCCGTA	TTCTAATATT	60
GAGAAATGAG	ATNAAACACA	ATNTTATAAA	GTCTACTTAG	AGAAGATCAA	GTGACCTCAA	120
AGACTTTACT	ATTTTCATAT	TTTAAGACAC	ATGATTTATC	CTATTTTAGT	AACCTGGTTC	180
ATACGTTAAA	CAAAGGATAA	TGTGAACAGC	AGAGAGGATT	TGTTGGCAGA	AAATCTATGT	240
TCAATCTNGA	ACTATCTANA	TCACAGACAT	TTCTATTCTT	TT		282

(2) INFORMATION FOR SEQ ID NO:117:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 305 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

ACACATGTCG	CTTCACTGCC	TTCTTAGATG	CTTCTGGTCA	ACATANAGGA	ACAGGGACCA	60
TATTTATCCT	CCCTCCTGAA	ACAATTGCAA	AATAANACAA	AATATATGAA	ACAATTGCAA	120
AATAAGGCAA	AATATATGAA	ACAACAGGTC	TCGAGATATT	GGAAATCAGT	CAATGAAGGA	180
TACTGATCCC	TGATCACTGT	CCTAATGCAG	GATGTGGGAA	ACAGATGAGG	TCACCTCTGT	240
GACTGCCCCA	GCTTACTGCC	TGTAGAGAGT	TTCTANGCTG	CAGTTCAGAC	AGGGAGAAAT	300
TGGGT						305

(2) INFORMATION FOR SEQ ID NO:118:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 71 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

ACCAAGGTGT NTGAATCTCT GACGTGGGGA TCTCTGATTC CCGCACAATC TGAGTGGAAA	60
AANTCCTGGG T	71

(2) INFORMATION FOR SEQ ID NO:119:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 212 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:119:

ACTCCGGTTG GTGTCAGCAG CACGTGGCAT TGAACATNGC AATGTGGAGC CCAAACCACA	60
GAAAATGGGG TGAAATTGGC CAACTTTCTA TNAACTTATG TTGGCAANTT TGCCACCAAC	120
AGTAAGCTGG CCCTTCTAAT AAAAGAAAAT TGAAAGGTTT CTCACTAANC GGAATTAANT	180
AATGGANTCA AGANACTCCC AGGCCTCAGC GT	212

(2) INFORMATION FOR SEQ ID NO:120:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 90 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:120:

ACTCGTTGCA NATCAGGGGC CCCCCAGAGT CACCGTTGCA GGAGTCCTTC TGGTCTTGCC	60
CTCCGCCGGC GCAGAACATG CTGGGGTGGT	90

(2) INFORMATION FOR SEQ ID NO:121:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 218 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:121:

TGTANCGTGA	ANACGACAGA	NAGGGTTGTC	AAAAATGGAG	AANCCTTGAA	GTCATTTTGA	60
GAATAAGATT	TGCTAAAAGA	TTTGGGGCTA	AAACATGGTT	ATTGGGAGAC	ATTTCTGAAG	120
ATATNCANGT	AAATTANGGA	ATGAATTCAT	GGTTCTTTTG	GGAATTCCTT	TACGATNGCC	180
AGCATANACT	TCATGTGGGG	ATANCAGCTA	CCCTTGTA			218

(2) INFORMATION FOR SEQ ID NO:122:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 171 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:122:

TAGGGGTGTA	TGCAACTGTA	AGGACAAAAA	TTGAGACTCA	ACTGGCTTAA	CCAATAAAGG	60
CATTTGTTAG	CTCATGGAAC	AGGAAGTCGG	ATGGTGGGGC	ATCTTCAGTG	CTGCATGAGT	120
CACCACCCCG	GCGGGGTCAT	CTGTGCCACA	GGTCCCTGTT	GACAGTGCGG	T	171

(2) INFORMATION FOR SEQ ID NO:123:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 76 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:123:

TGTAGCGTGA	AGACNACAGA	ATGGTGTGTG	CTGTGCTATC	CAGGAACACA	TTTATTATCA	60
TTATCAANTA	TTGTGT					76

(2) INFORMATION FOR SEQ ID NO:124:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 131 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

85220 "9090E050

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:124:

ACCTTTCCCC AAGGCCAATG TCCTGTGTGC TAACTGGCCG GCTGCAGGAC AGCTGCAATT	60
CAATGTGCTG GGTCAATATG AGGGGAGGAG ACTCTAAAAAT AGCCAATTTT ATTCTCTTGG	120
TTAAGATTG T	131

(2) INFORMATION FOR SEQ ID NO:125:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 432 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:125:

ACTTTATCTA CTGGCTATGA AATAGATGGT GGAAAATTGC GTTACCAACT ATACCACTGG	60
CTTGAAAAAG AGGTGATAGC TCTTCAGAGG ACTTGTGACT TTTGCTCAGA TGCTGAAGAA	120
CTACAGTCTG CATTGTGGCAG AAATGAAGAT GAATTTGGAT TAAATGAGGA TGCTGAAGAT	180
TTGCCTCACC AAACAAAAGT GAAACAACCTG AGAGAAAATT TTCAGGAAAA AAGACAGTGG	240
CTCTTGAAGT ATCAGTCACT TTTGAGAATG TTTCTTAGTT ACTGCATACT TCATGGATCC	300
CATGGTGGGG GTCTTGCATC TGTAAGAATG GAATTGATTT TGCTTTTGCA AGAATCTCAG	360
CAGGAAACAT CAGAACCACT ATTTTCTAGC CCTCTGTCAG AGCAAACCTC AGTGCCTCTC	420
CTCTTTGCTT GT	432

(2) INFORMATION FOR SEQ ID NO:126:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 112 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:126:

ACACAACCTTG AATAGTAAAA TAGAACTGA GCTGAAATTT CTAATTCACCT TTCTAACCAT	60
AGTAAGAATG ATATTTCCCC CCAGGGATCA CCAAATATTT ATAAAAATTT GT	112

(2) INFORMATION FOR SEQ ID NO:127:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:127:

ACCACGAAAC CACAAACAAG ATGGAAGCAT CAATCCACTT GCCAAGCACA GCAG

54

(2) INFORMATION FOR SEQ ID NO:128:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 323 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:128:

ACCTCATTAG TAATTGTTTT GTTGTTTCAT TTTTTTCTAA TGTCTCCCCT CTACCAGCTC	60
ACCTGAGATA ACAGAATGAA AATGGAAGGA CAGCCAGATT TCTCCTTTGC TCTCTGCTCA	120
TTCTCTCTGA AGTCTAGGTT ACCCATTTTG GGGACCCATT ATAGGCAATA AACACAGTTC	180
CCAAAGCATT TGGACAGTTT CTTGTTGTGT TTTAGAATGG TTTTCCTTTT TCTTAGCCTT	240
TTCTTGCAAA AGGCTCACTC AGTCCCTTGC TTGCTCAGTG GACTGGGCTC CCCAGGGCCT	300
AGGCTGCCTT CTTTTCATG TCC	323

(2) INFORMATION FOR SEQ ID NO:129:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 192 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:129:

ACATACATGT GTGTATATTT TTAAATATCA CTTTTGTATC ACTCTGACTT TTTAGCATAC	60
TGAAAACACA CTAACATAAT TTNTGTGAAC CATGATCAGA TACAACCCAA ATCATTTCATC	120
TAGCACATTC ATCTGTGATA NAAAGATAGG TGAGTTTCAT TTCCTTCACG TTGGCCAATG	180
GATAAACAAA GT	192

(2) INFORMATION FOR SEQ ID NO:130:

(i) SEQUENCE CHARACTERISTICS:

20250520 09:02:05

- (A) LENGTH: 362 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:130:

CCCTTTT	TTTA	TGGAATGAGT	AGACTGTATG	TTTGAANATT	TANCCACAAC	CTCTTTGACA	60
TATAATGACG	CAACAAAAAG	GTGCTGTTTA	GTCCTATGGT	TCAGTTTATG	CCCCTGACAA		120
GTTTCCATTG	TGTTTTGCCG	ATCTTCTGGC	TAATCGTGGT	ATCCTCCATG	TTATTAGTAA		180
TTCTGTATTC	CATTTTGTTA	ACGCCTGGTA	GATGTAACCT	GCTANGAGGC	TAACTTTATA		240
CTTATTTAAA	AGCTCTTATT	TTGTGGTCAT	TAAAAATGGCA	ATTTATGTGC	AGCACTTTAT		300
TGCAGCAGGA	AGCACGTGTG	GGTTGTTGT	AAAGCTCTTT	GCTAATCTTA	AAAAGTAATG		360
GG							362

(2) INFORMATION FOR SEQ ID NO:131:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 332 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:131:

CTTTT	TGAAA	GATCGTGTCC	ACTCCTGTGG	ACATCTTGTT	TTAATGGAGT	TTCCCATGCA	60
GTANGACTGG	TATGGTTGCA	GCTGTCCAGA	TAAAAACATT	TGAAGAGCTC	CAAAATGAGA		120
GTTCTCCCAG	GTTCCGCTTG	CTGCTCCAAG	TCTCAGCAGC	AGCCTCTTTT	AGGAGGCATC		180
TTCTGAACTA	GATTAAGGCA	GCTTGTAAT	CTGATGTGAT	TTGGTTTATT	ATCCAACTAA		240
CTTCCATCTG	TTATCACTGG	AGAAAGCCCA	GACTCCCCAN	GACNGGTACG	GATTGTGGGC		300
ATANAAGGAT	TGGGTGAAGC	TGGCCTTGTG	GT				332

(2) INFORMATION FOR SEQ ID NO:132:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 322 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:132:

ACTTTTGCCA	TTTTGTATAT	ATAAACAATC	TTGGGACATT	CTCCTGAAAA	CTAGGTGTCC	60
------------	------------	------------	------------	------------	------------	----

```

AGTGGCTAAG AGAACTCGAT TTCAAGCAAT TCTGAAAGGA AAACCAGCAT GACACAGAAT    120
CTCAAATTCC CAAACAGGGG CTCTGTGGGA AAAATGAGGG AGGACCTTTG TATCTCGGGT    180
TTTAGCAAGT TAAAATGAAN ATGACAGGAA AGGCTTATTT ATCAACAAAG AGAAGAGTTG    240
GGATGCTTCT AAAAAAACT TTGGTAGAGA AAATAGGAAT GCTNAATCCT AGGGAAGCCT    300
GTAACAATCT ACAATTGGTC CA                                           322

```

(2) INFORMATION FOR SEQ ID NO:133:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 278 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:133:

```

ACAAGCCTTC ACAAGTTTAA CTAAATTGGG ATTAATCTTT CTGTANTTAT CTGCATAATT    60
CTTGTTTTTC TTTCCATCTG GCTCCTGGGT TGACAATTTG TGGAAACAAC TCTATTGCTA    120
CTATTTAAAA AAAATCACAA ATCTTTCCCT TTAAGCTATG TTNAATTCAA ACTATTCCTG    180
CTATTCCTGT TTTGTCAAAG AAATTATATT TTCAAATA TGTNTATTTG TTTGATGGGT    240
CCCACGAAAC ACTAATAAAA ACCACAGAGA CCAGCCTG                               278

```

(2) INFORMATION FOR SEQ ID NO:134:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 121 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:134:

```

GTTTANAAAA CTTGTTTAGC TCCATAGAGG AAAGAATGTT AAACTTTGTA TTTTAAAAACA    60
TGATTCTCTG AGGTAAACT TGGTTTTCAT ATGTTATTTT TACTTGATT TTGCTTTTGG    120
T                                           121

```

(2) INFORMATION FOR SEQ ID NO:135:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 350 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:135:

ACTTANAACC	ATGCCTAGCA	CATCAGAATC	CCTCAAAGAA	CATCAGTATA	ATCCTATACC	60
ATANCAAGTG	GTGACTGGTT	AAGCGTGCGA	CAAAGGTCAG	CTGGCACATT	ACTTGTGTGC	120
AAACTTGATA	CTTTTGTTC	AAGTAGGAAC	TAGTATACAG	TNCCTAGGAN	TGGTACTCCA	180
GGGTGCCCCC	CAACTCCTGC	AGCCGCTCCT	CTGTGCCAGN	CCCTGNAAGG	AACTTTCGCT	240
CCACCTCAAT	CAAGCCCTGG	GCCATGCTAC	CTGCAATTGG	CTGAACAAAC	GTTTGCTGAG	300
TTCCCAAGGA	TGCAAAGCCT	GGTGCTCAAC	TCCTGGGGCG	TCAACTCAGT		350

(2) INFORMATION FOR SEQ ID NO:136:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 399 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:136:

TGTACCGTGA	AGACGACAGA	AGTTGCATGG	CAGGGACAGG	GCAGGGCCGA	GGCCAGGGTT	60
GCTGTGATTG	TATCCGAATA	NTCCTCGTGA	GAAAAGATAA	TGAGATGACG	TGAGCAGCCT	120
GCAGACTTGT	GTCTGCCTTC	AANAAGCCAG	ACAGGAAGGC	CCTGCCTGCC	TTGGCTCTGA	180
CCTGGCGGCC	AGCCAGCCAG	CCACAGGTGG	GCTTCTTCCT	TTTGTGGTGA	CAACNCCAAG	240
AAAAGTGCAG	AGGCCCAGGG	TCAGGTGTNA	GTGGGTANGT	GACCATAAAA	CACCAGGTGC	300
TCCCAGGAAC	CCGGGCAAAG	GCCATCCCCA	CCTACAGCCA	GCATGCCCAC	TGGCGTGATG	360
GGTGACAGANG	GATGAAGCAG	CCAGNTGTTC	TGCTGTGGT			399

(2) INFORMATION FOR SEQ ID NO:137:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 165 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:137:

ACTGGTGTGG	TNGGGGGTGA	TGCTGGTGGT	ANAAGTTGAN	GTGACTTCAN	GATGGTGTGT	60
GGAGGAAGTG	TGTGAACGTA	GGATGTAGA	NGTTTTGGCC	GTGCTAAATG	AGCTTCGGGA	120
TTGGCTGGTC	CCACTGGTGG	TACTGTTCAT	TGGTGGGGTT	CCTGT		165

(2) INFORMATION FOR SEQ ID NO:138:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 338 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:138:

ACTCACTGGA ATGCCACATT CACAACAGAA TCAGAGGTCT GTGAAAACAT TAATGGCTCC	60
TTAACTTCTC CAGTAAGAAT CAGGGACTTG AAATGGAAAC GTTAACAGCC ACATGCCCAA	120
TGCTGGGCAG TCTCCCATGC CTTCCACAGT GAAAGGGCTT GAGAAAAATC ACATCCAATG	180
TCATGTGTTT CCAGCCACAC CAAAAGGTGC TTGGGGTGGA GGGCTGGGGG CATANANGGT	240
CANGCCTCAG GAAGCCTCAA GTTCCATTCA GCTTTGCCAC TGTACATTCC CCATNTTTAA	300
AAAAACTGAT GCCTTTTTTT TTTTTTTTGT TAAAATTC	338

(2) INFORMATION FOR SEQ ID NO:139:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 382 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:139:

GGGAATCTTG GTTTTTGGCA TCTGGTTTGC CTATAGCCGA GGCCACTTTG ACAGAACAAA	60
GAAAGGGACT TCGAGTAAGA AGGTGATTTA CAGCCAGCCT AGTGCCCGAA GTGAAGGAGA	120
ATTCAAACAG ACCTCGTCAT TCCTGGTGTG AGCCTGGTCG GCTCACCGCC TATCATCTGC	180
ATTTGCCTTA CTCAGGTGCT ACCGGACTCT GGCCCCTGAT GTCTGTAGTT TCACAGGATG	240
CCTTATTTGT CTTCTACACC CCACAGGGCC CCCTACTTCT TCGGATGTGT TTTTAATAAT	300
GTCAGCTATG TGCCCCATCC TCCTTCATGC CCTCCCTCCC TTTCTACCA CTGCTGAGTG	360
GCCTGGAAct TGTTTAAAGT GT	382

(2) INFORMATION FOR SEQ ID NO:140:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 200 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:140:

ACCAAANCTT CTTTCTGTTG TGTTCGATTT TACTATAGGG GTTTCGCTTN TTCTAAANAT	60
ACTTTTCATT TAACANCTTT TGTTAAGTGT CAGGCTGCAC TTTGCTCCAT ANAATTATTG	120
TTTTCACATT TCAACTTGTA TGTGTTTGTG TCTTANAGCA TTGGTGAAAT CACATATTTT	180
ATATTCAGCA TAAAGGAGAA	200

(2) INFORMATION FOR SEQ ID NO:141:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 335 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:141:

ACTTTATTTT	CAAAACACTC	ATATGTTGCA	AAAAACACAT	AGAAAAATAA	AGTTTGGTGG	60
GGGTGCTGAC	TAAACTTCAA	GTCACAGACT	TTTATGTGAC	AGATTGGAGC	AGGGTTTGT	120
ATGCATGTAG	AGAACCCAAA	CTAATTTATT	AAACAGGATA	GAAACAGGCT	GTCTGGGTGA	180
AATGGTTCTG	AGAACCATCC	AATTCACCTG	TCAGATGCTG	ATANACTAGC	TCTTCAGATG	240
TTTTCCTACC	AGTTCAGAGA	TNGGTTAATG	ACTANTTCCA	ATGGGGAAAA	AGCAAGATGG	300
ATTACAAAAC	CAAGTAATTT	TAAACAAAAG	CACTT			335

(2) INFORMATION FOR SEQ ID NO:142:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 459 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:142:

ACCAGGTAA	TATTGCCACA	TATATCCTTT	CCAATTGCGG	GCTAAACAGA	CGTGTATTTA	60
GGGTGTTTA	AAGACAACCC	AGCTTAATAT	CAAGAGAAAT	TGTGACCTTT	CATGGAGTAT	120
CTGATGGAGA	AAACACTGAG	TTTTGACAAA	TCTTATTTTA	TTCAGATAGC	AGTCTGATCA	180
CACATGGTCC	AACAACACTC	AAATAATAAA	TCAAATATNA	TCAGATGTTA	AAGATTGGTC	240
TTCAAACATC	ATAGCCAATG	ATGCCCCGCT	TGCCTATAAT	CTCTCCGACA	TAAAACCACA	300
TCAACACCTC	AGTGCCACC	AAACCATTCA	GCACAGCTTC	CTTAAGTGTG	AGCTGTTTGA	360
AGCTACCAGT	CTGAGCACTA	TTGACTATNT	TTTTTCANGCT	CTGAATAGCT	CTAGGGATCT	420
CAGCANGGGT	GGGAGGAACC	AGCTCAACCT	TGGCGTANT			459

(2) INFORMATION FOR SEQ ID NO:143:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 140 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:143:

ACATTTCTCTT CCACCAAGTC AGGACTCCTG GCTTCTGTGG GAGTTCTTAT CACCTGAGGG	60
AAATCCAAAC AGTCTCTCCT AGAAAGGAAT AGTGTACCA ACCCCACCCA TCTCCCTGAG	120
ACCATCCGAC TTCCCTGTGT	140

(2) INFORMATION FOR SEQ ID NO:144:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 164 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:144:

ACTTCAGTAA CAACATACAA TAACAACATT AAGTGTATAT TGCCATCTTT GTCATTTTCT	60
ATCTATACCA CTCTCCCTTC TGAAAACAAN AATCACTANC CAATCACTTA TACAAATTG	120
AGGCAATTAA TCCATATTG TTTTCAATAA GGAAAAAAG ATGT	164

(2) INFORMATION FOR SEQ ID NO:145:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 303 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:145:

ACGTAGACCA TCCAACTTTG TATTTGTAAT GGCAAACATC CAGNAGCAAT TCCTAAACAA	60
ACTGGAGGGT ATTTATACCC AATTATCCCA TTCATTAACA TGCCCTCCTC CTCAGGCTAT	120
GCAGGACAGC TATCATAAGT CGGCCAGGC ATCCAGATAC TACCATTTGT ATAAACTTCA	180
GTAGGGGAGT CCATCCAAGT GACAGGTCTA ATCAAAGGAG GAAATGGAAC ATAAGCCCAG	240
TAGTAAAATN TTGCTTAGCT GAAACAGCCA CAAAAGACTT ACCGCCGTGG TGATTACCAT	300
CAA	303

(2) INFORMATION FOR SEQ ID NO:146:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 327 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20250909 09:09:09

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:146:

ACTGCAGCTC	AATTAGAAGT	GGTCTCTGAC	TTTCATCANC	TTCTCCCTGG	GCTCCATGAC	60
ACTGGCCTGG	AGTGACTCAT	TGCTCTGGTT	GGTTGAGAGA	GCTCCTTTGC	CAACAGGCCT	120
CCAAGTCAGG	GCTGGGATTT	GTTTCCTTTC	CACATTCTAG	CAACAATATG	CTGGCCACTT	180
CCTGAACAGG	GAGGGTGGGA	GGAGCCAGCA	TGGAACAAGC	TGCCACTTTC	TAAAGTAGCC	240
AGACTTGCCC	CTGGGCCTGT	CACACCTACT	GATGACCTTC	TGTGCCTGCA	GGATGGAATG	300
TAGGGGTGAG	CTGTGTGACT	CTATGGT				327

(2) INFORMATION FOR SEQ ID NO:147:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 173 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:147:

ACATTGTTTT	TTTGAGATAA	AGCATTGANA	GAGCTCTCCT	TAACGTGACA	CAATGGAAGG	60
ACTGGAACAC	ATACCCACAT	CTTTGTTCTG	AGGGATAATT	TTCTGATAAA	GTCTTGCTGT	120
ATATTCAAGC	ACATATGTTA	TATATTATTC	AGTTCCATGT	TTATAGCCTA	GTT	173

(2) INFORMATION FOR SEQ ID NO:148:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 477 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:148:

ACAACCACTT	TATCTCATCG	AATTTTAAAC	CCAAACTCAC	TCACTGTGCC	TTTCTATCCT	60
ATGGGATATA	TTATTTGATG	CTCCATTTC	TCACACATAT	ATGAATAATA	CACTCATACT	120
GCCCTACTAC	CTGCTGCAAT	AATCACATTC	CCTTCCTGTC	CTGACCCTGA	AGCCATTGGG	180
GTGGTCCTAG	TGGCCATCAG	TCCANGCCTG	CACCTTGAGC	CCTTGAGCTC	CATTGCTCAC	240
NCCANCCCC	CTCACCGACC	CCATCCTCTT	ACACAGCTAC	CTCCTTGCTC	TCTAACCCCA	300
TAGATTATNT	CCAAATTCAG	TCAATTAAGT	TACTATTAA	ACTCTACCCG	ACATGTCCAG	360
CACCACTGGT	AAGCCTTCTC	CAGCCAACAC	ACACACACAC	ACACNCACAC	ACACACATAT	420
CCAGGCACAG	GCTACCTCAT	CTTCACAATC	ACCCCTTTAA	TTACCATGCT	ATGGTGG	477

(2) INFORMATION FOR SEQ ID NO:149:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 207 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:149:

ACAGTTGTAT TATAATATCA AGAAATAAAC TTGCAATGAG AGCATTTAAG AGGGAAGAAC	60
TAACGTATTT TAGAGAGCCA AGGAAGGTTT CTGTGGGGAG TGGGATGTAA GGTGGGGCCT	120
GATGATAAAT AAGAGTCAGC CAGGTAAGTG GGTGGTGTGG TATGGGCACA GTGAAGAACA	180
TTTCAGGCAG AGGGAACAGC AGTGAAA	207

(2) INFORMATION FOR SEQ ID NO:150:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 111 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:150:

ACCTTGATTT CATTGCTGCT CTGATGGAAA CCCAACTATC TAATTTAGCT AAAACATGGG	60
CACTTAAATG TGGTCAGTGT TTGGACTTGT TAACTANTGG CATCTTTGGG T	111

(2) INFORMATION FOR SEQ ID NO:151:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 196 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:151:

AGCGCGGCAG GTCATATTGA ACATTCCAGA TACCTATCAT TACTCGATGC TGTGATAAC	60
AGCAAGATGG CTTTGAAGTC AGGGTCACCA CCAGCTATTG GACCTTACTA TGAAAACCAT	120
GGATACCAAC CGGAAAACCC CTATCCCGCA CAGCCCCTG TGGTCCCCAC TGTCTACGAG	180
GTGCATCCGG CTCAGT	196

(2) INFORMATION FOR SEQ ID NO:152:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 132 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:152:

ACAGCACTTT CACATGTAAG AAGGGAGAAA TTCCTAAATG TAGGAGAAAAG ATAACAGAAC	60
CTTCCCCTTT TCATCTAGTG GTGGAAACCT GATGCTTTAT GTTGACAGGA ATAGAACCAG	120
GAGGGAGTTT GT	132

(2) INFORMATION FOR SEQ ID NO:153:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 285 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:153:

ACAAANACCCA NGANAGGCCA CTGGCCGTGG TGTCATGGCC TCCAAACATG AAAGTGTGTCAG	60
CTTCTGCTCT TATGTCCTCA TCTGACAACT CTTTACCATT TTTATCCTCG CTCAGCAGGA	120
GCACATCAAT AAAGTCCAAA GTCTTGGA CTGGCCTTGGC TTGGAGGAAG TCATCAACAC	180
CCTGGCTAGT GAGGGTGCGG CGCCGCTCCT GGATGACGGC ATCTGTGAAG TCGTGCACCA	240
GTCTGCAGGC CCTGTGGAAG CGCCGTCCAC ACGGAGTNAG GAATT	285

(2) INFORMATION FOR SEQ ID NO:154:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 333 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:154:

ACCACAGTCC TGTTGGGCCA GGGCTTCATG ACCCTTTCCTG TGAAAAGCCA TATTATCACC	60
ACCCCAAATT TTTCCTTAAA TATCTTTAAC TGAAGGGGTC AGCCTCTTGA CTGCAAAGAC	120
CCTAAGCCGG TTACACAGCT AACTCCCACT GGCCCTGATT TGTGAAATTG CTGCTGCCTG	180
ATTGGCACAG GAGTCGAAGG TGTTCACTC CCCTCCTCCG TGGAACGAGA CTCTGATTG	240
AGTTTCACAA ATTCTCGGGC CACCTCGTCA TTGCTCCTCT GAAATAAAAT CCGGAGAATG	300

GTCAGGCCTG TCTCATCCAT ATGGATCTTC CGG

333

(2) INFORMATION FOR SEQ ID NO:155:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 308 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:155:

ACTGGAAATA ATAAAACCCA CATCACAGTG TTGTGTCAAA GATCATCAGG GCATGGATGG	60
GAAAGTGCTT TGGGAAGTGT AAAGTGCCTA ACACATGATC GATGATTTTT GTTATAATAT	120
TTGAATCAGG GTGCATACAA ACTCTCCTGC CTGCTCCTCC TGGGCCCCAG CCCCAGCCCC	180
ATCACAGCTC ACTGCTCTGT TCATCCAGGC CCAGCATGTA GTGGCTGATT CTTCTTGCT	240
GCTTTTAGCC TCCANAAGTT TCTCTGAAGC CAACCAAACC TCTANGTGTA AGGCATGCTG	300
GCCCTGGT	308

(2) INFORMATION FOR SEQ ID NO:156:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 295 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:156:

ACCTTGCTCG GTGCTTGGAA CATATTAGGA ACTCAAAATA TGAGATGATA ACAGTGCCTA	60
TTATTGATTA CTGAGAGAAC TGTTAGACAT TTAGTTGAAG ATTTTCTACA CAGGAAGTGA	120
GAATAGGAGA TTATGTTTGG CCCTCATATT CTCTCCTATC CTCCTTGCCT CATTCTATGT	180
CTAATATATT CTCAATCAA TAAGGTTAGC ATAATCAGGA AATCGACCAA ATACCAATAT	240
AAAACCAGAT GTCTATCCTT AAGATTTTCA AATAGAAAAC AAATTAACAG ACTAT	295

(2) INFORMATION FOR SEQ ID NO:157:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 126 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo sapiens

20250909 15:22:20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:157:

ACAAGTTTAA ATAGTGCTGT CACTGTGCAT GTGCTGAAAT GTGAAATCCA CCACATTTCT	60
GAAGAGCAAA ACAAATTCTG TCATGTAATC TCTATCTTGG GTCGTGGGTA TATCTGTCCC	120
CTTAGT	126

(2) INFORMATION FOR SEQ ID NO:158:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 442 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:158:

ACCCACTGGT CTTGGAAACA CCCATCCTTA ATACGATGAT TTTTCTGTCG TGTGAAAATG	60
AANCCAGCAG GCTGCCCCTA GTCAGTCCTT CCTTCCAGAG AAAAAGAGAT TTGAGAAAGT	120
GCCTGGGTAA TTCACCATT AATTCCTCCC CCAAACCTCTC TGAGTCTTCC CTTAATATTT	180
CTGGTGGTTC TGACCAAAGC AGGTCATGGT TTGTTGAGCA TTTGGGATCC CAGTGAAGTA	240
NATGTTTGTA GCCTTGCATA CTTAGCCCTT CCCACGCACA AACGGAGTGG CAGAGTGGTG	300
CCAACCCTGT TTTCCAGTC CACGTAGACA GATTCACAGT GCGGAATTCT GGAAGCTGGA	360
NACAGACGGG CTCTTTGCAG AGCCGGGACT CTGAGANGGA CATGAGGGCC TCTGCCTCTG	420
TGTTCAATTCT CTGATGTCCT GT	442

(2) INFORMATION FOR SEQ ID NO:159:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 498 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:159:

ACTTCCAGGT AACGTTGTTG TTTCCGTTGA GCCTGAACTG ATGGGTGACG TTGTAGGTTT	60
TCCAACAAGA ACTGAGGTTG CAGAGCGGGT AGGGAAGAGT GCTGTTCCAG TTGCACCTGG	120
GCTGCTGTGG ACTGTTGTTG ATTCCCTCACT ACGGCCCAAG GTTGTGGAAC TGGCANAAAG	180
GTGTGTTGTT GGANTTGAGC TCGGGCGGCT GTGGTAGGTT GTGGGCTCTT CAACAGGGGC	240
TGCTGTGGTG CCGGGANGTG AANGTGTTGT GTCACCTGAG CTTGGCCAGC TCTGGAAAGT	300
ANTANATTCT TCCTGAAGGC CAGCGCTTGT GGAGCTGGCA NGGGTCANTG TTGTGTGTAA	360
CGAACCAGTG CTGCTGTGGG TGGGTGTANA TCCTCCACAA AGCCTGAAGT TATGGTGTCTN	420
TCAGGTAANA ATGTGGTTTC AGTGTCCCTG GGCNGCTGTG GAAGGTTGTA NATTGTCAAC	480
AAGGGAATAA GCTGTGGT	498

(2) INFORMATION FOR SEQ ID NO:160:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 380 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:160:

ACCTGCATCC AGCTTCCCTG CCAAACCTCAC AAGGAGACAT CAACCTCTAG ACAGGGAAAC	60
AGCTTCAGGA TACTTCCAGG AGACAGAGCC ACCAGCAGCA AAACAAATAT TCCCATGCCT	120
GGAGCATGGC ATAGAGGAAG CTGANAAATG TGGGGTCTGA GGAAGCCATT TGAGTCTGGC	180
CACTAGACAT CTCATCAGCC ACTTGTGTGA AGAGATGCCC CATGACCCCA GATGCCTCTC	240
CCACCCTTAC CTCCATCTCA CACACTTGAG CTTTCCACTC TGTATAATTC TAACATCCTG	300
GAGAAAAATG GCAGTTTGAC CGAACCTGTT CACAACGGTA GAGGCTGATT TCTAACGAAA	360
CTTGTAGAAT GAAGCCTGGA	380

(2) INFORMATION FOR SEQ ID NO:161:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 114 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:161:

ACTCCACATC CCCTCTGAGC AGGCGGTTGT CGTTCAAGGT GTATTTGGCC TTGCCTGTCA	60
CACTGTCCAC TGGCCCCTTA TCCACTTGGT GCTTAATCCC TCGAAAGAGC ATGT	114

(2) INFORMATION FOR SEQ ID NO:162:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 177 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:162:

ACTTTCTGAA TCGAATCAAA TGATACTTAG TGTAGTTTAA ATATCCTCAT ATATATCAAA	60
GTTTTACTAC TCTGATAATT TTGTAAACCA GGTAACCAGA ACATCCAGTC ATACAGCTTT	120
TGGTGATATA TAACTTGGCA ATAACCCAGT CTGGTGATAC ATAAAACTAC TCACTGT	177

(2) INFORMATION FOR SEQ ID NO:163:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 137 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:163:

CATTTATACA GACAGGCGTG AAGACATTCA CGACAAAAAC GCGAAATTCT ATCCCGTGAC	60
CANAGAAGGC AGCTACGGCT ACTCCTACAT CCTGGCGTGG GTGGCCTTCG CCTGCACCTT	120
CATCAGCGGC ATGATGT	137

(2) INFORMATION FOR SEQ ID NO:164:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 469 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:164:

CTTATCACAA TGAATGTTCT CCTGGGCAGC GTTGTGATCT TTGCCACCTT CGTGACTTTA	60
TGCAATGCAT CATGCTATTT CATACTAAT GAGGGAGTTC CAGGAGATTC AACCAGGAAA	120
TGCATGGATC TCAAAGGAAA CAAACACCCA ATAAACTCGG AGTGGCAGAC TGACAACTGT	180
GAGACATGCA CTTGCTACGA AACAGAAATT TCATGTTGCA CCCTTGTTTC TACACCTGTG	240
GGTTATGACA AAGACAACCTG CCAAAGAATC TTCAAGAAGG AGGACTGCAA GTATATCGTG	300
GTGGAGAAGA AGGACCCAAA AAAGACCTGT TCTGTCACTG AATGGATAAT CTAATGTGCT	360
TCTAGTAGGC ACAGGGCTCC CAGGCCAGGC CTCATTCTCC TCTGGCCTCT AATAGTCAAT	420
GATTGTGTAG CCATGCCTAT CAGTAAAAAG ATNTTTGAGC AAACACTTT	469

(2) INFORMATION FOR SEQ ID NO:165:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 195 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:165:

ACAGTTTTTT ATANATATCG ACATTGCCGG CACTTGTGTT CAGTTTCATA AAGCTGGTGG	60
---	----

ATCCGCTGTC ATCCACTATT CCTTGGCTAG AGTAAAAATT ATTCTTATAG CCCATGTCCC	120
TGCAGGCCGC CCGCCCGTAG TTCTCGTTCC AGTCGTCTTG GCACACAGGG TGCCAGGACT	180
TCCTCTGAGA TGAGT	195

(2) INFORMATION FOR SEQ ID NO:166:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 383 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:166:

ACATCTTAGT AGTGTGGCAC ATCAGGGGGC CATCAGGGTC ACAGTCACTC ATAGCCTCGC	60
CGAGGTCGGA GTCCACACCA CCGGTGTAGG TGTGCTCAAT CTTGGGCTTG GCGCCACCT	120
TTGGAGAAGG GATATGCTGC ACACACATGT CCACAAAGCC TGTGAACTCG CCAAAGAATT	180
TTTGCAGACC AGCCTGAGCA AGGGGCGGAT GTTCAGCTTC AGCTCCTCCT TCGTCAGGTG	240
GATGCCAACC TCGTCTANGG TCCGTGGGAA GCTGGTGTCC ACNTCACCTA CAACCTGGGC	300
GANGATCTTA TAAAGAGGCT CCNAGATAAA CTCCACGAAA CTTCTCTGGG AGCTGCTAGT	360
NGGGGCCTTT TTGGTGAAC TTC	383

(2) INFORMATION FOR SEQ ID NO:167:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 247 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:167:

ACAGAGCCAG ACCTTGGCCA TAAATGAANC AGAGATTAAG ACTAAACCCC AAGTCGANAT	60
TGGAGCAGAA ACTGGAGCAA GAAGTGGGCC TGGGGCTGAA GTAGAGACCA AGGCCACTGC	120
TATANCCATA CACAGAGCCA ACTCTCAGGC CAAGGCNATG GTTGGGGCAG ANCCAGAGAC	180
TCAATCTGAN TCCAAAGTGG TGGCTGGAAC ACTGGTCATG ACANAGGCAG TGA CTCTGAC	240
TGANGTC	247

(2) INFORMATION FOR SEQ ID NO:168:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 273 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:168:

ACTTCTAAGT TTTCTAGAAG TGGAAGGATT GTANTCATCC TGAAAATGGG TTTACTTCAA	60
AATCCCTCAN CCTTGTTCTT CACNACTGTC TATACTGANA GTGTCATGTT TCCACAAAGG	120
GCTGACACCT GAGCCTGNAT TTTCACTCAT CCCTGAGAAG CCCTTTCAG TAGGGTGGGC	180
AATTCCCAAC TTCCTTGCCA CAAGCTTCCC AGGCTTTCTC CCCTGGAAAA CTCCAGCTTG	240
AGTCCCAGAT ACACTCATGG GCTGCCCTGG GCA	273

(2) INFORMATION FOR SEQ ID NO:169:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 431 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:169:

ACAGCCTTGG CTTCCCCAAA CTCCACAGTC TCAGTGCAGA AAGATCATCT TCCAGCAGTC	60
AGCTCAGACC AGGGTCAAAG GATGTGACAT CAACAGTTTC TGGTTTCAGA ACAGGTTCTA	120
CTACTGTCAA ATGACCCCCC AACTTCCCTC AAAGGCTGTG GTAAGTTTTG CACAGGTGAG	180
GGCAGCAGAA AGGGGGTANT TACTGATGGA CACCATCTTC TCTGTATACT CCACACTGAC	240
CTTGCCATGG GCAAAGGCC CTACCACAAA AACAATAGGA TCACTGCTGG GCACCAGCTC	300
ACGCACATCA CTGACAACCG GGATGGAAAA AGAANTGCCA ACTTTCATAC ATCCAAGTGG	360
AAAGTGATCT GATACTGGAT TCTTAATTAC CTTCAAAAGC TTCTGGGGGC CATCAGCTGC	420
TCGAACACTG A	431

(2) INFORMATION FOR SEQ ID NO:170:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 266 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:170:

ACCTGTGGGC TGGGCTGTTA TGCCTGTGCC GGCTGCTGAA AGGGAGTTCA GAGGTGGAGC	60
TCAAGGAGCT CTGCAGGCAT TTTGCCAANC CTCTCCANAG CANAGGGAGC AACCTACACT	120
CCCCGCTAGA AAGACACCAG ATTGGAGTCC TGGGAGGGGG AGTTGGGGTG GGCATTTGAT	180
GTATACTTGT CACCTGAATG AANGAGCCAG AGAGGAANGA GACGAANATG ANATTGGCCT	240
TCAAAGCTAG GGGTCTGGCA GGTGGA	266

(2) INFORMATION FOR SEQ ID NO:171:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1248 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:171:

GGCAGCCAAA	TCATAAACGG	CGAGGACTGC	AGCCCGCACT	CGCAGCCCTG	GCAGGCGGCA	60
CTGGTCATGG	AAAACGAATT	GTTCTGCTCG	GGCGTCCTGG	TGCATCCGCA	GTGGGTGCTG	120
TCAGCCGCAC	ACTGTTTCCA	GAAGTGAGTG	CAGAGCTCCT	ACACCATCGG	GCTGGGCGTG	180
CACAGTCTTG	AGGCCGACCA	AGAGCCAGGG	AGCCAGATGG	TGGAGGCCAG	CCTCTCCGTA	240
CGGCACCCAG	AGTACAACAG	ACCCTTGCTC	GCTAACGACC	TCATGCTCAT	CAAGTTGGAC	300
GAATCCGTGT	CCGAGTCTGA	CACCATCCGG	AGCATCAGCA	TTGCTTCGCA	GTGCCCTACC	360
GCGGGGAACT	CTTGCTTCGT	TTCTGGCTGG	GGTCTGCTGG	CGAACGGCAG	AATGCCTACC	420
GTGCTGCAGT	GCGTGAACGT	GTCGGTGGTG	TCTGAGGAGG	TCTGCAGTAA	GCTCTATGAC	480
CCGCTGTACC	ACCCACGAT	GTTCTGCGCC	GGCGGAGGGC	AAGACCAGAA	GGACTCCTGC	540
AACGGTGACT	CTGGGGGGCC	CCTGATCTGC	AACGGGTACT	TGCAGGGCCT	TGTGTCTTTC	600
GGAAAAGCCC	CGTGTGGCCA	AGTTGGCGTG	CCAGGTGTCT	ACACCAACCT	CTGCAAATTC	660
ACTGAGTGGA	TAGAGAAAAC	CGTCCAGGCC	AGTTAACTCT	GGGGACTGGG	AACCCATGAA	720
ATTGACCCCC	AAATACATCC	TGCGGAAGGA	ATTCAGGAAT	ATCTGTTCCC	AGCCCCCTCT	780
CCCTCAGGCC	CAGGAGTCCA	GGCCCCAGC	CCCTCCTCCC	TCAAACCAAG	GGTACAGATC	840
CCCAGCCCCCT	CCTCCCTCAG	ACCCAGGAGT	CCAGACCCCC	CAGCCCCCTCC	TCCCTCAGAC	900
CCAGGAGTCC	AGCCCCCTCT	CCCTCAGACC	CAGGAGTCCA	GACCCCCCAG	CCCCCTCCTCC	960
CTCAGACCCA	GGGGTCCAGG	CCCCCAACCC	CTCCTCCCTC	AGACTCAGAG	GTCCAAGCCC	1020
CCAACCCNTC	ATTCCCCAGA	CCCAGAGGTC	CAGGTCCCAG	CCCCTCNTCC	CTCAGACCCA	1080
GCGGTCCAAT	GCCACCTAGA	CTNTCCCTGT	ACACAGTGCC	CCCTTGTTGG	ACGTTGACCC	1140
AACCTTACCA	GTTGGTTTTT	CATTTTTNGT	CCCTTTCCCC	TAGATCCAGA	AATAAAGTTT	1200
AAGAGAAGNG	CAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA		1248

(2) INFORMATION FOR SEQ ID NO:172:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 159 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:172:

Met	Val	Glu	Ala	Ser	Leu	Ser	Val	Arg	His	Pro	Glu	Tyr	Asn	Arg	Pro
1					5				10					15	
Leu	Leu	Ala	Asn	Asp	Leu	Met	Leu	Ile	Lys	Leu	Asp	Glu	Ser	Val	Ser
			20					25					30		
Glu	Ser	Asp	Thr	Ile	Arg	Ser	Ile	Ser	Ile	Ala	Ser	Gln	Cys	Pro	Thr

35	40	45
Ala Gly Asn Ser Cys Leu Val Ser Gly Trp Gly Leu Leu Ala Asn Gly		
50	55	60
Arg Met Pro Thr Val Leu Gln Cys Val Asn Val Ser Val Val Ser Glu		
65	70	75
Glu Val Cys Ser Lys Leu Tyr Asp Pro Leu Tyr His Pro Ser Met Phe		
85	90	95
Cys Ala Gly Gly Gly Gln Xaa Gln Xaa Asp Ser Cys Asn Gly Asp Ser		
100	105	110
Gly Gly Pro Leu Ile Cys Asn Gly Tyr Leu Gln Gly Leu Val Ser Phe		
115	120	125
Gly Lys Ala Pro Cys Gly Gln Val Gly Val Pro Gly Val Tyr Thr Asn		
130	135	140
Leu Cys Lys Phe Thr Glu Trp Ile Glu Lys Thr Val Gln Ala Ser		
145	150	155

(2) INFORMATION FOR SEQ ID NO:173:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1265 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:173:

GGCAGCCCCG	ACTCGCAGCC	CTGGCAGGCG	GCACTGGTCA	TGGAAAACGA	ATTGTTCTGC	60
TCGGGCGTCC	TGGTGTCATCC	GCAGTGGGTG	CTGTTCAGCCG	CACACTGTTT	CCAGAACTCC	120
TACACCATCG	GGCTGGGCGT	GCACAGTCTT	GAGGCCGACC	AAGAGCCAGG	GAGCCAGATG	180
GTGGAGGCCA	GCCTCTCCGT	ACGGCACCCA	GAGTACAACA	GACCCCTTGCT	CGCTAACGAC	240
CTCATGCTCA	TCAAGTTGGA	CGAATCCGTG	TCCGAGTCTG	ACACCATCCG	GAGCATCAGC	300
ATTGCTTCGC	AGTGCCCTAC	CGCGGGGAAC	TCTTGCCCTCG	TTTCTGGCTG	GGGTCTGCTG	360
GCGAACGGTG	AGCTCACGGG	TGTGTGTCTG	CCCTCTTCAA	GGAGGTCCTC	TGCCCAGTCG	420
CGGGGGCTGA	CCCAGAGCTC	TGCGTCCCAG	GCAGAATGCC	TACCGTGCTG	CAGTGCGTGA	480
ACGTGTCCGT	GGTGTCTGAG	GAGGTCTGCA	GTAAGCTCTA	TGACCCGCTG	TACCACCCCA	540
GCACTGTTCTG	CGCCGGCGGA	GGGCAAGACC	AGAAGGACTC	CTGCAACGGT	GACTCTGGGG	600
GGCCCCTGAT	CTGCAACGGG	TACTTGCAGG	GCCTTGTGTC	TTTCGGAAAA	GCCCCGTGTG	660
GCCAAGTTGG	CGTGCCAGGT	GTCTACACCA	ACCTCTGCAA	ATTCACTGAG	TGGATAGAGA	720
AAACCGTCCA	GGCCAGTTAA	CTCTGGGGAC	TGGGAACCCA	TGAAATTGAC	CCCCAAATAC	780
ATCCTGCGGA	AGGAATTTCAG	GAATATCTGT	TCCCAGCCCC	TCCTCCCTCA	GGCCCAGGAG	840
TCCAGGCCCC	CAGCCCCTCC	TCCCTCAAAC	CAAGGGTACA	GATCCCCAGC	CCCTCCTCCC	900
TCAGACCCAG	GAGTCCAGAC	CCCCCAGCCC	CTCCTCCCTC	AGACCCAGGA	GTCCAGCCCC	960
TCCTCCNTCA	GACCCAGGAG	TCCAGACCCC	CCAGCCCCTC	CTCCCTCAGA	CCCAGGGGTT	1020
GAGGCCCCCA	ACCCCTCCTC	CTTCAGAGTC	AGAGGTCCAA	GCCCCCAACC	CCTCGTTCCC	1080
CAGACCCAGA	GGTNNAGGTC	CCAGCCCCTC	TTCCNTCAGA	CCCAGNGGTC	CAATGCCACC	1140
TAGATTTTCC	CTGNACACAG	TGCCCCCTTG	TGGNANGTTG	ACCCAACCTT	ACCAGTTGGT	1200

TTTTTCATTTT TNGTCCCTTT CCCCTAGATC CAGAAATAAA GTTTAAGAGA NGNGCAAAAA 1260
AAAAA 1265

(2) INFORMATION FOR SEQ ID NO:174:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1459 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:174:

GGTCAGCCGC AACTGTTTC CAGAAGTGAG TGCAGAGCTC CTACACCATC GGGCTGGGCC 60
 TGCACAGTCT TGAGGCCGAC CAAGAGCCAG GGAGCCAGAT GGTGGAGGCC AGCCTCTCCG 120
 TACGGCACCC AGAGTACAAC AGACCCTTGC TCGCTAACGA CCTCATGCTC ATCAAGTTGG 180
 ACGAATCCGT GTCCGAGTCT GACACCATCC GGAGCATCAG CATTGCTTCG CAGTGCCTTA 240
 CCGCGGGGAA CTCTTGCTC GTTTCTGGCT GGGGTCTGCT GGCGAACGGT GAGCTCACGG 300
 GTGTGTGTCT GCCCTCTTCA AGGAGGTCCT CTGCCCAGTC GCGGGGGCTG ACCCAGAGCT 360
 CTGCGTCCCA GGCAGAATGC CTACCGTGCT GCAGTGCGTG AACGTGTCGG TGGTGTCTGA 420
 NGAGGTCTGC ANTAAGCTCT ATGACCCGCT GTACCACCCC ANCATGTTCT GCGCCGGCGG 480
 AGGGCAAGAC CAGAAGGACT CCTGCAACGT GAGAGAGGGG AAAGGGGAGG GCAGGCGACT 540
 CAGGGAAGGG TGGAGAAGGG GGAGACAGAG ACACACAGGG CCGCATGGCG AGATGCAGAG 600
 ATGGAGAGAC ACACAGGGAG ACAGTGACAA CTAGAGAGAG AAAGTGAGAG AAACAGAGAA 660
 ATAAACACAG GAATAAAGAG AAGCAAAGGA AGAGAGAAAC AGAAACAGAC ATGGGGAGGC 720
 AGAAACACAC ACACATAGAA ATGCAGTTGA CCTTCCAACA GCATGGGGCC TGAGGGCGGT 780
 GACCTCCACC CAATAGAAAA TCCTCTTATA ACTTTTGACT CCCCCAAAAC CTGACTAGAA 840
 ATAGCCTACT GTTGACGGGG AGCCTTACCA ATAACATAAA TAGTCGATTT ATGCATACGT 900
 TTTATGCATT CATGATATAC CTTTGTGTTGA ATTTTGTGAT ATTTCTAAGC TACACAGTTC 960
 GTCTGTGAAT TTTTAAAT TGTTGCAACT CTCCTAAAT TTTTCTGATG TGTTTATTGA 1020
 AAAAATCCAA GTATAAGTGG ACTTGTGCAT TCAAACCAGG GTTGTTCAGG GGTCAACTGT 1080
 GTACCCAGAG GGAAACAGTG ACACAGATTC ATAGAGGTGA AACACGAAGA GAAACAGGAA 1140
 AAATCAAGAC TCTACAAAGA GGCTGGGCAG GGTGGCTCAT GCCTGTAATC CCAGCACTTT 1200
 GGGAGGCGAG GCAGGCAGAT CACTTGAGGT AAGGAGTTCA AGACCAGCCT GGCCAAAATG 1260
 GTGAAATCCT GTCTGTACTA AAAATACAAA AGTTAGCTGG ATATGGTGGC AGGCGCCTGT 1320
 AATCCCAGCT ACTTGGGAGG CTGAGGCAGG AGAATTGCTT GAATATGGGA GGCAGAGGTT 1380
 GAAGTGAGTT GAGATCACAC CACTATACTC CAGCTGGGGC AACAGAGTAA GACTCTGTCT 1440
 CAAAAA AAAAAA 1459

(2) INFORMATION FOR SEQ ID NO:175:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1167 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:175:

(2) INFORMATION FOR SEQ ID NO:176:

(A) LENGTH: 205 amino acids

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

Met Glu Asn Glu Leu Phe Cys Ser Gly Val Leu Val His Pro Gln Trp
1 5 10 15

Val Leu Ser Ala Ala His Cys Phe Gln Asn Ser Tyr Thr Ile Gly Leu
20 25 30

Gly Leu His Ser Leu Glu Ala Asp Gln Glu Pro Gly Ser Gln Met Val
35 40 45

Glu Ala Ser Leu Ser Val Arg His Pro Glu Tyr Asn Arg Leu Leu Leu
50 55 60

Ala Asn Asp Leu Met Leu Ile Lys Leu Asp Glu Ser Val Ser Glu Ser
65 70 75 80

Asp Thr Ile Arg Ser Ile Ser Ile Ala Ser Gln Cys Pro Thr Ala Gly
85 90 95

Asn Ser Cys Leu Val Ser Gly Trp Gly Leu Leu Ala Asn Gly Arg Met
100 105 110

```

Pro Thr Val Leu His Cys Val Asn Val Ser Val Val Ser Glu Xaa Val
    115                      120                      125

Cys Ser Lys Leu Tyr Asp Pro Leu Tyr His Pro Ser Met Phe Cys Ala
    130                      135                      140

Gly Gly Gly Gln Asp Gln Lys Asp Ser Cys Asn Gly Asp Ser Gly Gly
    145                      150                      155                      160

Pro Leu Ile Cys Asn Gly Tyr Leu Gln Gly Leu Val Ser Phe Gly Lys
    165                      170                      175

Ala Pro Cys Gly Gln Leu Gly Val Pro Gly Val Tyr Thr Asn Leu Cys
    180                      185                      190

Lys Phe Thr Glu Trp Ile Glu Lys Thr Val Gln Xaa Ser
    195                      200                      205

```

(2) INFORMATION FOR SEQ ID NO:177:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1119 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:177:

```

GCGCACTCGC AGCCCTGGCA GCGGCACTG GTCATGGAAA ACGAATTGTT CTGCTCGGGC      60
GTCCTGGTGC ATCCGCAGTG GGTGCTGTCA GCCGCACACT GTTTCAGAA CTCCTACACC      120
ATCGGGCTGG GCCTGCACAG TCTTGAGGCC GACCAAGAGC CAGGGAGCCA GATGGTGGAG      180
GCCAGCCTCT CCGTACGGCA CCCAGAGTAC AACAGACCCT TGCTCGCTAA CGACCTCATG      240
CTCATCAAGT TGGACGAATC CGTGTCGAG TCTGACACCA TCCGGAGCAT CAGCATTGCT      300
TCGCAGTGCC CTACCGCGGG GAACTCTTGC CTCGTTTCTG GCTGGGGTCT GCTGGCGAAC      360
GATGCTGTGA TTGCCATCCA GTCCCAGACT GTGGGAGGCT GGGAGTGTGA GAAGCTTTCC      420
CAACCTGGC AGGGTTGTAC CATTTTCGGA ACTTCCAGTG CAAGGACGTC CTGCTGCATC      480
CTCACTGGGT GCTCACTACT GCTCACTGCA TCACCCGGAA CACTGTGATC AACTAGCCAG      540
CACCATAGTT CTCCGAAGTC AGACTATCAT GATTACTGTG TTGACTGTGC TGTCTATTGT      600
ACTAACCATG CCGATGTTTA GGTGAAATTA GCGTCACTTG GCCTCAACCA TCTTGGTATC      660
CAGTTATCCT CACTGAATTG AGATTTCTCTG CTTCAAGGAT GCTGGTACTC CCCTCACAAA      720
TGACCTACAG AGGTGAGGGA TCATATAGCT CTTCAAGGAT GCTGGTACTC CCCTCACAAA      780
TTCATTTCTC CTGTTGTAGT GAAAGGTGCG CCCTCTGGAG CCTCCCAGGG TGGGTGTGCA      840
GGTCACAATG ATGAATGTAT GATCGTGTTC CCATTACCCA AAGCCTTTAA ATCCCTCATG      900
CTCAGTACAC CAGGGCAGGT CTAGCATTTT TGCATTTAGT GTATGCTGTC CATTATGCA      960
ACCACCTCAG GACTCCTGGA TTCTCTGCCT AGTTGAGCTC CTGCATGCTG CCTCCTTGGG      1020
GAGGTGAGGG AGAGGGCCCA TGGTTCAATG GGATCTGTGC AGTTGTAACA CATTAGGTGC      1080
TTAATAAACA GAAGCTGTGA TGTAAAAAAA AAAAAAAAAA      1119

```

(2) INFORMATION FOR SEQ ID NO:178:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 164 amino acids
 - (B) TYPE: amino acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:178:

```

Met Glu Asn Glu Leu Phe Cys Ser Gly Val Leu Val His Pro Gln Trp
1          5          10          15

Val Leu Ser Ala Ala His Cys Phe Gln Asn Ser Tyr Thr Ile Gly Leu
          20          25          30

Gly Leu His Ser Leu Glu Ala Asp Gln Glu Pro Gly Ser Gln Met Val
          35          40          45

Glu Ala Ser Leu Ser Val Arg His Pro Glu Tyr Asn Arg Pro Leu Leu
          50          55          60

Ala Asn Asp Leu Met Leu Ile Lys Leu Asp Glu Ser Val Ser Glu Ser
65          70          75          80

Asp Thr Ile Arg Ser Ile Ser Ile Ala Ser Gln Cys Pro Thr Ala Gly
          85          90          95

Asn Ser Cys Leu Val Ser Gly Trp Gly Leu Leu Ala Asn Asp Ala Val
          100          105          110

Ile Ala Ile Gln Ser Xaa Thr Val Gly Gly Trp Glu Cys Glu Lys Leu
          115          120          125

Ser Gln Pro Trp Gln Gly Cys Thr Ile Ser Ala Thr Ser Ser Ala Arg
          130          135          140

Thr Ser Cys Cys Ile Leu Thr Gly Cys Ser Leu Leu Leu Thr Ala Ser
145          150          155          160

Pro Gly Thr Leu

```

(2) INFORMATION FOR SEQ ID NO:179:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 250 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:179:

```

CTGGAGTGCC TTGGTGTTTC AAGCCCCTGC AGGAAGCAGA ATGCACCTTC TGAGGCACCT      60
CCAGCTGCCC CCGGCCGGGG GATGCGAGGC TCGGAGCACC CTTGCCCGGC TGTGATTGCT      120
GCCAGGCACT GTTCATCTCA GCTTTTCTGT CCCTTTGCTC CCGGCAAGCG CTTCTGCTGA      180
AAGTTCATAT CTGGAGCCTG ATGTCTTAAC GAATAAAGGT CCCATGCTCC ACCCGAAAAA      240
AAAAAAAAAA                                     250

```

(2) INFORMATION FOR SEQ ID NO:180:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 202 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:180:

ACTAGTCCAG	TGTGGTGGAA	TTCCATTGTG	TTGGGCCCAA	CACAATGGCT	ACCTTTAACA	60
TCACCCAGAC	CCCGCCCCTG	CCCGTGCCCC	ACGCTGCTGC	TAACGACAGT	ATGATGCTTA	120
CTCTGCTACT	CGGAAACTAT	TTTTATGTAA	TTAATGTATG	CTTTCTTGTT	TATAAATGCC	180
TGATTTAAAA	AAAAAAAAAA	AA				202

(2) INFORMATION FOR SEQ ID NO:181:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 558 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:181:

TCCYTTTGKT	NAGGTTTKKG	AGACAMCCCK	AGACCTWAAN	CTGTGTCACA	GACTTCYNGG	60
AATGTTTAGG	CAGTGCTAGT	AATTTCTCG	TAATGATTCT	GTTATTACTT	TCCTNATTCT	120
TTATTCCTCT	TTCTTCTGAA	GATTAATGAA	GTTGAAAATT	GAGGTGGATA	AATACAAAAA	180
GGTAGTGTGA	TAGTATAAGT	ATCTAAGTGC	AGATGAAAGT	GTGTTATATA	TATCCATTCA	240
AAATTATGCA	AGTTAGTAAT	TACTCAGGGT	TAACTAAATT	ACTTTAATAT	GCTGTTGAAC	300
CTACTCTGTT	CCTTGGCTAG	AAAAAATTAT	AAACAGGACT	TTGTTAGTTT	GGGAAGCCAA	360
ATTGATAATA	TTCTATGTTT	TAAAAGTTGG	GCTATACATA	AATTATTAAG	AAATATGGAW	420
TTTTATTCCC	AGGAATATGG	KGTTTCATTTT	ATGAATATTA	CSCRGGATAG	AWGTWTGAGT	480
AAAAYCAGTT	TTGGTWAATA	YGTWAATATG	TCMTAAATAA	ACAAGGCTTT	GACTTATTTC	540
CAAAAAAAA	AAAAAAA					558

(2) INFORMATION FOR SEQ ID NO:182:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 479 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:182:

ACAGGGWTTK	GRGGATGCTA	AGSCCCCRGA	RWTYGTTTGA	TCCAACCCTG	GCTTWTTTTTC	60
AGAGGGGAAA	ATGGGGCCTA	GAAGTTACAG	MSCATYTAGY	TGGTGCGMTG	GCACCCCTGG	120
CSTCACACAG	ASTCCCAGT	AGCTGGGACT	ACAGGCACAC	AGTCACTGAA	GCAGGCCCTG	180
TTWGCAATTC	ACGTTGCCAC	CTCCAACCTA	AACATTCTTC	ATATGTGATG	TCCTTAGTCA	240
CTAAGGTAA	ACTTTCCCAC	CCAGAAAAGG	CAACTTAGAT	AAAATCTTAG	AGTACTTTCA	300
TACTMTTCTA	AGTCCTCTTC	CAGCCTCACT	KKGAGTCCTM	CYTGGGGGTT	GATAGGAANT	360
NTCTCTTGGC	TTTCTCAATA	AARTCTCTAT	YCATCTCATG	TTTAATTTGG	TACGCATARA	420
AWTGSTGARA	AAATTAAAA	GTTCTGGTTY	MACTTTAAAA	AAAAAAAAAA	AAAAAAAAAA	479

(2) INFORMATION FOR SEQ ID NO:183:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:183:

AGGCGGGAGC	AGAAGCTAAA	GCCAAAGCCC	AAGAAGAGTG	GCAGTGCCAG	CACTGGTGCC	60
AGTACCAGTA	CCAATAACAG	TGCCAGTGCC	AGTGCCAGCA	CCAGTGGTGG	CTTCAGTGCT	120
GGTGCCAGCC	TGACCGCCAC	TCTCACATTT	GGGCTCTTCG	CTGGCCTTGG	TGGAGCTGGT	180
GCCAGCACCA	GTGGCAGCTC	TGGTGCCTGT	GGTTTCTCCT	ACAAGTGAGA	TTTGTAGATAT	240
TGTTAATCCT	GCCAGTCTTT	CTCTTCAAGC	CAGGGTGCAT	CCTCAGAAAC	CTACTCAACA	300
CAGCACTCTA	GGCAGCCACT	ATCAATCAAT	TGAAGTTGAC	ACTCTGCATT	ARATCTATTT	360
GCCATTTCAA	AAAAAAAAAA	AAAA				384

(2) INFORMATION FOR SEQ ID NO:184:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:184:

ACCGAATTGG	GACCGCTGGC	TTATAAGCGA	TCATGTYTNT	CCRGATATKAC	CTCAACGAGC	60
AGGGAGATCG	AGTCTATACG	CTGAAGAAAT	TTGACCCGAT	GGGACAACAG	ACCTGCTCAG	120
CCCATCCTGC	TCGGTTCTCC	CCAGATGACA	AATACTCTSG	ACACCGAATC	ACCATCAAGA	180
AACGCTTCAA	GGTGCTCATG	ACCCAGCAAC	CGCGCCCTGT	CCTCTGAGGG	TCCCTTAAAC	240
TGATGTCTTT	TCTGCCACCT	GTTACCCCTC	GGAGACTCCG	TAACCAAAC	CTTCGGACTG	300
TGAGCCCTGA	TGCCTTTTTC	CCAGCCATAC	TCTTTGGCAT	CCAGTCTCTC	GTGGCGATTG	360
ATTATGCTTG	TGTGAGGCAA	TCATGGTGGC	ATCACCCATA	AAGGGAACAC	ATTTGACTTT	420
TTTTTCTCAT	ATTTTAAATT	ACTACMAGAW	TATTWMAGAW	WAAATGAWTT	GAAAAACTST	480
TAAAAAAAAA	AAAAAA					496

(2) INFORMATION FOR SEQ ID NO:185:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:185:

GCTGGTAGCC	TATGGCGKGG	CCCACGGAGG	GGCTCCTGAG	GCCACGGRAC	AGTGAATTCC	60
CAAGTATCYT	GCGSGCGTC	TTCTACCGTC	CCTACCTGCA	GATCTTCGGG	CAGATTCCCC	120
AGGAGGACAT	GGACGTGGCC	CTCATGGAGC	ACAGCAACTG	YTCGTCGGAG	CCCGGCTTCT	180
GGGCACACCC	TCCTGGGGCC	CAGGCGGGCA	CCTGCGTCTC	CCAGTATGCC	AACTGGCTGG	240
TGGTGCTGCT	CCTCGTCATC	TTCTGCTCG	TGGCCAACAT	CCTGCTGGTC	AACTTGCTCA	300
TTGCCATGTT	CAGTTACACA	TTGGGCAAAG	TACAGGGCAA	CAGCGATCTC	TACTGGGAAG	360

GCGCAGCGTT ACCGCCTCAT CCGG

384

(2) INFORMATION FOR SEQ ID NO:186:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 577 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:186:

GAGTTAGCTC	CTCCACAACC	TTGATGAGGT	CGTCTGCAGT	GGCCTCTCGC	TTCATACCGC	60
TNCCATCGTC	ATACTGTAGG	TTTGCCACCA	CYTCTGGCA	TCTTGGGGCG	GCNTAATATT	120
CCAGGAAACT	CTCAATCAAG	TCACCGTCGA	TGAAACCTGT	GGGCTGGTTC	TGTCTTCCGC	180
TCGGTGTGAA	AGGATCTCCC	AGAAGGAGTG	CTCGATCTTC	CCCACACTTT	TGATGACTTT	240
ATTGAGTCGA	TTCTGCATGT	CCAGCAGGAG	GTTGTACCAG	CTCTCTGACA	GTGAGGTCAC	300
CAGCCCTATC	ATGCCGTTGA	MCGTGCCGAA	GARCACCGAG	CCTTGTGTGG	GGGKKGAAGT	360
CTCACCAGAG	TTCTGCATTA	CCAGAGAGCC	GTGGCAAAAG	ACATTGACAA	ACTCGCCCAG	420
GTGGAAAAAG	AMCAMCTCCT	GGARGTGCTN	GCCGCTCCTC	GTCMGTGGT	GGCAGCGCTW	480
TCCTTTTGAC	ACACAAACAA	GTTAAAGGCA	TTTTTCAGCCC	CCAGAAANTT	GTCATCATCC	540
AAGATNTCGC	ACAGCACTNA	TCCAGTTGGG	ATTAAAT			577

(2) INFORMATION FOR SEQ ID NO:187:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 534 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:187:

AACATCTTCC	TGTATAATGC	TGTGTAATAT	CGATCCGATN	TTGTCTGSTG	AGAATYCATW	60
ACTKGGAAAA	GMAACATTAA	AGCCTGGACA	CTGGTATTAA	AATTCACAAT	ATGCAACACT	120
TTAAACAGTG	TGTCAATCTG	CTCCCYYNAC	TTTGTTCATCA	CCAGTCTGGG	AACAAGGGTA	180
TGCCCTATTG	ACACCTGTGA	AAAGGGCGCT	AAGCATTTTT	GATTCAACAT	CTTTTTTTTT	240
GACACAAGTC	CGAAAAAAGC	AAAAGTAAAC	AGTTATYAAT	TTGTTAGCCA	ATTCACTTTC	300
TTTCATGGGAC	AGAGCCATYT	GATTTAAAAA	GCAAATTGCA	TAATATTGAG	CTTYGGGAGC	360
TGATATTTGA	GCGGAAGAGT	AGCCTTTCTA	CTTCACCAGA	CACAACTCCC	TTTCATATTG	420
GGATGTTNAC	NAAAGTWATG	TCTCTWACAG	ATGGGATGCT	TTTGTGGCAA	TTCTGTTCTG	480
AGGATCTCCC	AGTTTATTTA	CACTTGCAC	AAGAAGGCGT	TTTCTTCCTC	AGGC	534

(2) INFORMATION FOR SEQ ID NO:188:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 761 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:188:

AGAAACCAGT	ATCTCTNAAA	ACAACCTCTC	ATACCTTGTG	GACCTAATTT	TGTGTGCGTG	60
TGTGTGTGCG	CGCATATTAT	ATAGACAGGC	ACATCTTTTT	TACTTTTGTA	AAAGCTTATG	120

CCTCTTTGGT	ATCTATATCT	GTGAAAAGTTT	TAATGATCTG	CCATAATGTC	TTGGGGACCT	180
TTGTCTTCTG	TGTAAATGGT	ACTAGAGAAA	ACACCTATNT	TATGAGTCAA	TCTAGTTNGT	240
TTTATTTCGAC	ATGAAGGAAA	TTTCCAGATN	ACAACACTNA	CAAACCTCTCC	CTKGACKARG	300
GGGGACAAAG	AAAAGCAAAA	CTGAMCATAA	RAAACAATWA	CCTGGTGAGA	ARTTGCATAA	360
ACAGAAATWR	GGTAGTATAT	TGAARNACAG	CATCATTAATA	RMGTTWTKTT	WTTCTCCCTT	420
GCAAAAAACA	TGTACNGACT	TCCCCTTGAG	TAATGCCAAG	TTGTTTTTTT	TATNATAAAA	480
CTTGCCCTTC	ATTACATGTT	TNAAAGTGGT	GTGGTGGGCC	AAAATATTGA	AATGATGGAA	540
CTGACTGATA	AAGCTGTACA	AATAAGCAGT	GTGCCTAACA	AGCAACACAG	TAATGTTGAC	600
ATGCTTAATT	CACAAATGCT	AATTTTATTA	TAAATGTTTG	CTAAAATACA	CTTTGAACTA	660
TTTTTCTGTN	TTCCCAGAGC	TGAGATNTTA	GATTTTATGT	AGTATNAAGT	GAAAAANTAC	720
GAAAATAATA	ACATTGAAGA	AAAANANAAA	AAANAAAAAA	A		761

(2) INFORMATION FOR SEQ ID NO:189:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 482 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:189:

TTTTTTTTTT	TTTGCCGATN	CTACTATTTT	ATTGCAGGAN	GTGGGGGTGT	ATGCACCGCA	60
CACCGGGGCT	ATNAGAAGCA	AGAAGGAAGG	AGGGAGGGCA	CAGCCCCTTG	CTGAGCAACA	120
AAGCCGCTTG	CTGCCTTCTC	TGTCTGTCTC	CTGGTGCAGG	CACATGGGGA	GACCTTCCCC	180
AAGGCAGGGG	CCACCAGTCC	AGGGGTGGGA	ATACAGGGGG	TGGGANGTGT	GCATAAGAAG	240
TGATAGGCAC	AGGCCACCCG	GTACAGACCC	CTCGGCTCCT	GACAGGTNGA	TTTCGACCAG	300
GTCATTGTGC	CCTGCCCAGG	CACAGCGTAN	ATCTGGAATA	GACAGAATGC	TTTCCTTTTC	360
AAATTTGGCT	NGTCATNGAA	NGGGCANTTT	TCCAANTTNG	GCTNGGTCTT	GGTACNCTTG	420
GTTCGGCCCA	GCTCCNCGTC	CAAAAANTAT	TCACCCNNCT	CCNAATTGCT	TGCNNGNCCC	480
CC						482

(2) INFORMATION FOR SEQ ID NO:190:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 471 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:190:

TTTTTTTTTT	TTTTAAAACA	GTTTTTCACA	ACAAAATTTA	TTAGAAGAAT	AGTGGTTTTG	60
AAAACCTCTG	CATCCAGTGA	GAACCTACAT	ACACCACATT	ACAGCTNGGA	ATGTNCTCCA	120
AATGTCTGGT	CAAATGATAC	AATGGAACCA	TTCAATCTTA	CACATGCACG	AAAGAACAAG	180
CGCTTTTGAC	ATACAATGCA	CAAAAAAAAA	AGGGGGGGGG	GACCACATGG	ATTAAAATTT	240
TAAGTACTCA	TCACATACAT	TAAGACACAG	TTCTAGTCCA	GTCNAAAATC	AGAACTGCNT	300
TGAAAAATTT	CATGTATGCA	ATCCAACCAA	AGAACTTNAT	TGGTGATCAT	GANTNCTCTA	360
CTACATCNAC	CTTGATCATT	GCCAGGAACN	AAAAGTTNAA	ANCACNCNGT	ACAAAAANAA	420
TCTGTAATTN	ANTTCAACCT	CCGTACNGAA	AAATNTTNNT	TATACACTCC	C	471

(2) INFORMATION FOR SEQ ID NO:191:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 402 base pairs
- (B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:191:

GAGGGATTGA	AGGTCTGTTC	TASTGTCTGGM	CTGTTTCAGCC	ACCAACTCTA	ACAAGTTGCT	60
GTCTTCCACT	CACTGTCTGT	AAGCTTTTTA	ACCCAGACWG	TATCTTCATA	AATAGAACAA	120
ATTCTTCACC	AGTCACATCT	TCTAGGACCT	TTTTGGATTG	AGTTAGTATA	AGCTCTTCCA	180
CTTCCTTTGT	TAAGACTTCA	TCTGGTAAAG	TCTTAAGTTT	TGTAGAAAGG	AATTYAATTG	240
CTCGTTCTCT	AACAATGTCC	TCTCCTTGAA	GTATTTGGCT	GAACAACCCA	CCTAAAGTCC	300
CTTTGTGCAT	CCATTTTAAA	TATACTTAAT	AGGGCATTGK	TNCACTAGGT	TAAATTCTGC	360
AAGAGTCATC	TGTCTGCAA	AGTTGCGTTA	GTATATCTGC	CA		402

(2) INFORMATION FOR SEQ ID NO:192:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 601 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:192:

GAGCTCGGAT	CCAATAATCT	TTGTCTGAGG	GCAGCACACA	TATNCAGTGC	CATGGNAACT	60
GGTCTACCCC	ACATGGGAGC	AGCATGCCGT	AGNTATATAA	GGTCATTCCC	TGAGTCAGAC	120
ATGCYTYTTT	GAYTACCGTG	TGCCAAGTGC	TGGTGATTCT	YAACACACYT	CCATCCCGYT	180
CTTTTGTGGA	AAAAGTGGCA	CTTKTCTGGA	ACTAGCARGA	CATCACTTAC	AAATTCACCC	240
ACGAGACACT	TGAAAGGTGT	AACAAAGCGA	YTCTTGCATT	GCTTTTGTGTC	CCTCCGGCAC	300
CAGTTGTCAA	TACTAACCCG	CTGGTTTGCC	TCCATCACAT	TTGTGATCTG	TAGCTCTGGA	360
TACATCTCCT	GACAGTACTG	AAGAACTTCT	TCTTTTGTTT	CAAAAGCARC	TCTTGGTGCC	420
TGTTGGATCA	GGTTCCCATT	TCCCAGTCYG	AATGTTTACA	TGGCATATTT	WACTTCCCAC	480
AAAACATTGC	GATTTGAGGC	TCAGCAACAG	CAAATCCTGT	TCCGGCATTG	GCTGCAAGAG	540
CCTCGATGTA	GCCGGCCAGC	GCCAAGGCAG	GCGCCGTGAG	CCCCACCAGC	AGCAGAAGCA	600
G						601

(2) INFORMATION FOR SEQ ID NO:193:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 608 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:193:

ATACAGCCCCA	NATCCCACCA	CGAAGATGCG	CTTGTTGACT	GAGAACCTGA	TGCGGTCACT	60
GGTCCCGCTG	TAGCCCCAGC	GACTCTCCAC	CTGCTGGAAG	CGGTTGATGC	TGCACTCYTT	120
CCCAACGCAG	GCAGMAGCGG	GSCCGGTCAA	TGAACTCCAY	TCGTGGCTTG	GGGTKGACGG	180
TKAAGTGCA	GAAGAGGCTG	ACCACCTCGC	GGTCCACCAG	GATGCCCCGAC	TGTGCGGGAC	240
CTGCAGCGAA	ACTCCTCGAT	GGTCATGAGC	GGGAAGCGAA	TGAGGCCCGAG	GGCCTTGCCC	300
AGAACCTTCC	GCCTGTTCCTC	TGGCGTCACC	TGCAGCTGCT	GCCGCTGACA	CTCGGCCTCG	360
GACCAGCGGA	CAAACGGCRT	TGAACAGCCG	CACCTCACGG	ATGCCCAAGT	TGTCGCGCTC	420
CAGGAMMGSC	ACCAGCGTGT	CCAGGTCAAT	GTCGGTGAAG	CCCTCCGCGG	GTRATGGCGT	480
CTGCAGTGTT	TTTGTCGATG	TTCTCCAGGC	ACAGGCTGGC	CAGCTGCGGT	TCATCGAAGA	540
GTCGCGCCTG	CGTGAGCAGC	ATGAAGGCGT	TGTCGGCTCG	CAGTTCTTCT	TCAGGAACTC	600

CACGCAAT

608

(2) INFORMATION FOR SEQ ID NO:194:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 392 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:194:

GAACGGCTGG	ACCTTGCCCTC	GCATTGTGCT	TGCTGGCAGG	GAATACCTTG	GCAAGCAGYT	60
CCAGTCCGAG	CAGCCCCAGA	CCGCTGCCGC	CCGAAGCTAA	GCCTGCCCTCT	GGCCTTCCCC	120
TCCGCCTCAA	TGCAGAACCA	GTAGTGGGAG	CACCTGTGTTT	AGAGTTAAGA	GTGAACACTG	180
TTTGATTTTA	CTTGGAATT	TCCTCTGTTA	TATAGCTTTT	CCCAATGCTA	ATTTCCAAAC	240
AACAACAACA	AAATAACATG	TTTGCCTGTT	AAGTTGTATA	AAAGTAGGTG	ATTCTGTATT	300
TAAAGAAAAT	ATTACTGTTA	CATATACTGC	TTGCAATTC	TGTATTTATT	GKTNCTSTGG	360
AAATAAATAT	AGTTATTAAA	GGTTGTCANT	CC			392

(2) INFORMATION FOR SEQ ID NO:195:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 502 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:195:

CCSTTKGAGG	GGTKAGGKYC	CAGTTYCCGA	GTGGAAGAAA	CAGGCCAGGA	GAAGTGCCTG	60
CCGAGCTGAG	GCAGATGTTC	CCACAGTGAC	CCCCAGAGCC	STGGGSTATA	GTYTCTGACC	120
CCTCNCAAGG	AAAGACCACS	TTCTGGGGAC	ATGGGCTGGA	GGGCAGGACC	TAGAGGCACC	180
AAGGGAAGGC	CCCATTCCGG	GGSTGTTCCC	CGAGGAGGAA	GGGAAGGGGC	TCTGTGTGCC	240
CCCCASGAGG	AAGAGGCCCT	GAGTCCTGGG	ATCAGACACC	CCTTCACGTG	TATCCCCACA	300
CAAATGCAAG	CTCACCAAGG	TCCCCTCTCA	GTCCCCTTCC	STACACCCTG	AMCGGCCACT	360
GSCSCACACC	CACCCAGAGC	ACGCCACCCG	CCATGGGGAR	TGTGCTCAAG	GARTCGCNGG	420
GCARCGTGGA	CATCTNGTCC	CAGAAGGGGG	CAGAATCTCC	AATAGANGGA	CTGARCMSTT	480
GCTNANAAAA	AAAAANAAAA	AA				502

(2) INFORMATION FOR SEQ ID NO:196:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 665 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:196:

GGTTACTTGG	TTTCATTGCC	ACCACTTAGT	GGATGTCATT	TAGAACCATT	TTGTCTGCTC	60
CCTCTGGAAG	CCTTGCGCAG	AGCGGACTTT	GTAATTGTTG	GAGAATAACT	GCTGAATTTT	120
WAGCTGTTTK	GAGTTGATTS	GCACCACTGC	ACCCACAAC	TCAATATGAA	AACYAWTTGA	180
ACTWATTTAT	TATCTTGTTGA	AAAGTATAAC	AATGAAAATT	TTGTTTCATAC	TGTATTKATC	240
AAGTATGATG	AAAAGCAAWA	GATATATATT	CTTTTATTAT	GTAAATTAT	GATTGCCATT	300

ATTAATCGGC	AAAATGTGGA	GTGTATGTTC	TTTTACACAGT	AATATATGCC	TTTTGTAACT	360
TCACCTGGTT	ATTTTATTGT	AAATGARTTA	CAAAATTCTT	AATTTAAGAR	AATGGTATGT	420
WATATTTATT	TCATTAATTT	CTTTCCTKGT	TTACGTWAAT	TTTGAAAAGA	WTGCATGATT	480
TCTTGACAGA	AATCGATCTT	GATGCTGTGG	AAGTAGTTTG	ACCCACATCC	CTATGAGTTT	540
TTCTTAGAAT	GTATAAAGGT	TGTAGCCCAT	CNAACTTCAA	AGAAAAAAT	GACCACATAC	600
TTTGCAATCA	GGCTGAAATG	TGGCATGCTN	TTCTAATTC	AACTTTATAA	ACTAGCAAAN	660
AAGTG						665

(2) INFORMATION FOR SEQ ID NO:197:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 492 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:197:

TTTTNTTTTT	TTTTTTTTGC	AGGAAGGATT	CCATTTATTG	TGGATGCATT	TTCACAATAT	60
ATGTTTATTG	GAGCGATCCA	TTATCAGTGA	AAAGTATCAA	GTGTTTATAA	NATTTTTAGG	120
AAGGCAGATT	CACAGAACAT	GCTNGTCNGC	TTGCAGTTTT	ACCTCGTANA	GATNACAGAG	180
AATTTATAGTC	NAACCAGTAA	ACNAGGAATT	TACTTTTCAA	AAGATTAAAT	CCAAACTGAA	240
CAAAATTCTA	CCCTGAAACT	TACTCCATCC	AAATATTGGA	ATAANAGTCA	GCAGTGATAC	300
ATTCTCTTCT	GAACTTTAGA	TTTTCTAGAA	AAATATGTAA	TAGTGATCAG	GAAGAGCTCT	360
TGTTCAAAAG	TACAACNAAG	CAATGTTCCC	TTACCATAGG	CCTTAATTCA	AACTTTGATC	420
CATTTCACTC	CCATCACGGG	AGTCAATGCT	ACCTGGGACA	CTTGTATTTT	GTTTCATNCTG	480
ANCNTGGCTT	AA					492

(2) INFORMATION FOR SEQ ID NO:198:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 478 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:198:

TTTNTTTTGN	ATTTCAANTCT	GTANNAANTA	TTTTCATTTAT	GTTTATTANA	AAAATATNAA	60
TGTNTCCACN	ACAAATCATN	TTACNTNAGT	AAGAGGCCAN	CTACATTGTA	CAACATACAC	120
TGAGTATATT	TTGAAAAGGA	CAAGTTTAAA	GTANACNCAT	ATTGCCGANC	ATANCACATT	180
TATACATGGC	TTGATTGATA	TTTAGCACAG	CANAACTGA	GTGAGTTACC	AGAAAANAAT	240
NATATATGTC	AATCNGATTT	AAGATACAAA	ACAGATCCTA	TGGTACATAN	CATCNTGTAG	300
GAGTTGTGGC	TTTATGTTTA	CTGAAAGTCA	ATGCAGTTCC	TGTACAAAAGA	GATGGCCGTA	360
AGCATTCCTAG	TACCTCTACT	CCATGGTTAA	GAATCGTACA	CTTATGTTTA	CATATGTNCA	420
GGGTAAGAAT	TGTGTTAAGT	NAANTTATGG	AGAGGTCCAN	GAGAAAAATT	TGATNCAA	478

(2) INFORMATION FOR SEQ ID NO:199:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 482 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:199:

AGTGA	CTTGT	CCTCCA	ACAA	AACCCCT	TGA	TCAAGT	TTTGT	GGCACT	GACA	ATCAGAC	CCTA	60
TGCTAG	TTCC	TGTCAT	CTAT	TCGCTAC	TAA	ATGCAG	ACTG	GAGGGG	ACCA	AAAAGGG	GCA	120
TCAACT	CCAG	CTGGAT	TATT	TTGGAG	CCTG	CAAATC	TATT	CCTACT	TGTA	CGGACT	TTGA	180
AGTGAT	TCAG	TTTCCT	CTAC	GGATG	GAGA	CTGGCT	CAAG	AATATC	CCTCA	TGCAGC	TTTA	240
TGAAGC	CNAC	TCTGA	ACAC	GCTGGT	TATCT	NAGATG	GAGAA	NCAGAG	AAAT	AAAGTC	NAGA	300
AAATTT	ACCT	GGANGA	AAAAG	AGGCTT	TNNG	CTGGGG	ACCA	TCCCATT	GAA	CCTTCT	CTTA	360
ANGGAC	TTTA	AGAANA	AACT	ACCACAT	GTN	TGTNGT	TATCC	TGGTGC	CNGG	CCGTTT	ANTG	420
AACNTN	GACN	NCACCCT	TNT	GGAATAN	ANT	CTTGAC	NGCN	TCCTGA	ACTT	GCTCCT	CTGC	480
GA												482

(2) INFORMATION FOR SEQ ID NO:200:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 270 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:200:

CGGCCG	CAAG	TGCAAC	TCCA	GCTGGG	GCCG	TGCGGA	CGAA	GATTCT	GGCA	GCAGTT	GGTC	60
CGACTG	CGAC	GACGGC	GGCG	GCGAC	AGTC	GAGGTG	CAGC	GCGGGC	GCCT	GGGGT	CTTG	120
AAGGCT	GAGC	TGACGC	CGCA	GAGGTC	GTGT	CACGTC	CCAC	GACCTT	GACG	CCGTC	GGGA	180
CAGCCG	GAAC	AGAGCC	CGGT	GAANGC	GGGA	GGCCTC	GGGG	AGCCCC	CTCG	GAAGGG	CGGC	240
CCGAG	AGATA	CGCAGG	TGCA	GGTGGC	CGCC							270

(2) INFORMATION FOR SEQ ID NO:201:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 419 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:201:

TTTTTT	TTTT	TGGA	ATC	TACTGC	GAGC	ACAGCA	GAGT	AGCAAC	AAAGT	TTATTT	TGCA	60
GCTAGC	AAAG	TAACAG	GGTA	GGGCAT	TGGT	ACATGT	TCAG	GTCAAC	TTCC	TTTGT	CGTG	120
TTGATT	TGGT	TGTCTT	TATG	GGGGC	GGGGT	GGGGT	AGGG	AAANCG	AAGC	ANAANTA	AACA	180
TGGAGT	TGGG	GCACCC	TCCC	TGTAGA	ACCT	GGTTAC	NAAA	GCTTGG	GGCA	GTTTAC	CCTG	240
TCTGTG	ACCG	TCATTT	TCTT	GACATC	CAATG	TTATTAG	AAG	TCAGGA	TATC	TTTTAG	AGAG	300
TCCACT	GTNT	CTGGAG	GGGAG	ATTAGG	GTTT	CTTGCC	AAANA	TCCAAN	CAAA	ATCCAC	NTGA	360
AAAAGT	TGGA	TGATNC	CANGT	ACNGA	AATACC	GANGGC	CATAN	TTCTCA	TANT	CGGTGG	CCA	419

(2) INFORMATION FOR SEQ ID NO:202:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 509 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:202:

TTTNTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	60
TGGCACTTAA	TCCATTTTTA	TTTCAAAATG	TCTACAAANT	TTNAATNCNC	CATTATACNG	120
GTNATTTTNC	AAAATCTAAA	NNTTATTCAA	ATNTNAGCCA	AANTCCTTAC	NCAAATNNAA	180
TACNCNCAAA	AATCAAAAAT	ATACNTNTCT	TTCAGCAAAC	TTNGTTACAT	AAATTAAGAA	240
AATATATACG	GCTGGTGTTC	TCAAAGTACA	ATTATCTTAA	CACTGCAAAC	ATNTTTNNAA	300
GGAAGTAAAA	TAAAAAATAA	CACTNCCGCA	AAGGTTAAAG	GGAACAACAA	ATTCNTTTTA	360
CAACANCNNC	NATTATAAAA	ATCATATCTC	AAATCCTTAGG	GGAATATATA	CTTCACACNG	420
GGATCTTAAC	TTTTACTNCA	CTTTGTTTAT	TTTTTTANAA	CCATTGTNTT	GGGCCCAACA	480
CAATGGNAAT	NCCNCCNCNC	TGGACTAGT				509

(2) INFORMATION FOR SEQ ID NO:203:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 583 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:203:

TTTTTTTTTT	TTTTTTTTGA	CCCCCTCTT	ATAAAAACA	AGTTACCATT	TTATTTTACT	60
TACACATATT	TATTTTATAA	TTGGTATTAG	ATATTCAAAA	GGCAGCTTTT	AAAATCAAAC	120
TAAATGGAAA	CTGCCCTAGA	TACATAATTC	TTAGGAATTA	GCTTAAAATC	TGCCATAAGT	180
GAAAATCTTC	TCTAGCTCTT	TTGACTGTAA	ATTTTGTGACT	CTTGTAAGAA	ATCCAAATTC	240
ATTTTCTCTG	TCTTTAAAAA	TATCTAATCT	TTCCATTTT	TCCCTATTCC	AAGTCAATTT	300
GCTTCTCTAG	CCTCATTTCC	TAGCTCTTAT	CTACTATTAG	TAAGTGGCTT	TTTTCTTAAA	360
AGGGAAAAACA	GGAAGAGANA	ATGGCACACA	AAACAAACAT	TTTATATTCA	TATTTCTACC	420
TACGTTAATA	AAATAGCATT	TTGTGAAGCC	AGCTCAAAAG	AAGGCTTAGA	TCCTTTTATG	480
TCCATTTTAG	TCACTAAACG	ATATCNAAAG	TGCCAGAATG	CAAAAGGTTT	GTGAACATTT	540
ATTCAAAAGC	TAATATAAGA	TATTTACAT	ACTCATCTTT	CTG		583

(2) INFORMATION FOR SEQ ID NO:204:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 589 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:204:

TTTTTTTTNT	TTTTTTTTTT	TTTTTTNCTC	TTCTTTTTTT	TTGANAATGA	GGATCGAGTT	60
TTTCACTCTC	TAGATAGGGC	ATGAAGAAAA	CTCATCTTTC	CAGCTTTTAA	ATAACAATCA	120
AATCTCTTAT	GCTATATCAT	ATTTTAAAGT	AAACTAATGA	GTCAGTGGCT	TATCTTCTCC	180
TGAAGGAAAT	CTGTTTATT	TTCTCATTTA	TATAGTTATA	TCAAGTACTA	CCTTGCATAT	240
TGAGAGGTTT	TTCTTCTCTA	TTTACACATA	TATTTCCATG	TGAATTTGTA	TCAAACCTTT	300
ATTTTCATGC	AAACTAGAAA	ATAATGINTT	CTTTTGCATA	AGAGAAGAGA	ACAATATNAG	360
CATTACAAAA	CTGCTCAAAAT	TGTTTGTTAA	GNTTATCCAT	TATAATTAGT	TNGGCAGGAG	420
CTAATACAAA	TCACATTTAC	NGACNAGCAA	TAATAAAACT	GAAGTACCAG	TTAAATATCC	480
AAAATAATTA	AAGGAACATT	TTTAGCCTGG	GTATAATTAG	CTAATTCAT	TTACAAGCAT	540
TTATTNAGAA	TGAATTCACA	TGTTATTATT	CCNTAGCCCA	ACACAATGG		589

(2) INFORMATION FOR SEQ ID NO:205:

(i) SEQUENCE CHARACTERISTICS:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:205:

(2) INFORMATION FOR SEQ ID NO:206:

(A) LENGTH: 487 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:206:

(2) INFORMATION FOR SEQ ID NO:207:

(A) LENGTH: 332 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:207:

TGAATTGGCT	AAAAGACTGC	ATTTTTANAA	CTAGCAACTC	TTATTTCTTT	CCTTTAAAAA	60
TACATAGCAT	TAAATCCCAA	ATCCTATTTA	AAGACCTGAC	AGCTTGAGAA	GGTCACTACT	120
GCATTTATAG	GACCTTCTGG	TGGTTCCTGCT	GTTACNTTTG	AANTCTGACA	ATCCTTGANA	180
ATCTTTGCAT	GCAGAGGAGG	TAAAAGGTAT	TGGATTTTCA	CAGAGGAANA	ACACAGCGCA	240
GAATGAAGG	GGCCAGGCTT	ACTGAGCTTG	TCCACTGGAG	GGCTCATGGG	TGGGACATGG	300

AAAAGAAGGC AGCCTAGGCC CTGGGGAGCC CA

332

(2) INFORMATION FOR SEQ ID NO:208:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 524 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:208:

AGGGCGTGGT	GCGGAGGGCG	TTACTGTTTT	GTCTCAGTAA	CAATAAATAC	AAAAAGACTG	60
GTTGTGTTCC	GGCCCCATCC	AACCACGAAG	TTGATTTCTC	TTGTGTGCAG	AGTGACTGAT	120
TTTAAAGGAC	ATGGAGCTTG	TCACAATGTC	ACAATGTCAC	AGTGTGAAGG	GCACACTCAC	180
TCCCGCGTGA	TTCACATTTA	GCAACCAACA	ATAGCTCATG	AGTCCATACT	TGTAAATACT	240
TTTGGCAGAA	TACTTNTTGA	AACTTGCAGA	TGATAACTAA	GATCCAAGAT	ATTTCCCAAA	300
GTAAATAGAA	GTGGGTCATA	ATATTAATTA	CCTGTTCACA	TCAGCTTCCA	TTTACAAGTC	360
ATGAGCCAG	ACACTGACAT	CAAACCTAAGC	CCACTTAGAC	TCCTCACCAC	CAGTCTGTCC	420
TGTCATCAGA	CAGGAGGCTG	TCACCTTGAC	CAAATCTCTCA	CCAGTCAATC	ATCTATCCAA	480
AAACCATTAC	CTGATCCACT	TCCGGTAATG	CACCACCTTG	GTGA		524

(2) INFORMATION FOR SEQ ID NO:209:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 159 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:209:

GGGTGAGGAA	ATCCAGAGTT	GCCATGGAGA	AAATTCCAGT	GTCAGCATTC	TTGCTCCTTG	60
TGGCCCTCTC	CTACACTCTG	GCCAGAGATA	CCACAGTCAA	ACCTGGAGCC	AAAAAGGACA	120
CAAAGGACTC	TCGACCCAAA	CTGCCCCAGA	CCCTCTCCA			159

(2) INFORMATION FOR SEQ ID NO:210:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 256 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:210:

ACTCCCTGGC	AGACAAAGGC	AGAGGAGAGA	GCTCTGTTAG	TTCTGTGTTG	TTGAACTGCC	60
ACTGAATTTT	TTTCCACTTG	GACTATTACA	TGCCANTTGA	GGGACTAATG	GAAAAACGTA	120
TGGGGAGATT	TTANCCAATT	TANGTNTGTA	AATGGGGAGA	CTGGGGCAGG	CGGGAGAGAT	180
TTGCAGGGTG	NAAATGGGAN	GGCTGGTTTG	TTANATGAAC	AGGGACATAG	GAGGTAGGCA	240
CCAGGATGCT	AAATCA					256

(2) INFORMATION FOR SEQ ID NO:211:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:211:

ACATTGTTTT	TTTGAGATAA	AGCATTGAGA	GAGCTCTCCT	TAACGTGACA	CAATGGAAGG	60
ACTGGAACAC	ATACCCACAT	CTTTGTTCTG	AGGGATAATT	TTCTGATAAA	GTCTTGCTGT	120
ATATTCAAGC	ACATATGTTA	TATATTATTC	AGTTCCATGT	TTATAGCCTA	GTTAAGGAGA	180
GGGGAGATAC	ATTCNGAAAG	AGGACTGAAA	GAAATACTCA	AGTNGGAAAA	CAGAAAAAGA	240
AAAAAAGGAG	CAAATGAGAA	GCCT				264

(2) INFORMATION FOR SEQ ID NO:212:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 328 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:212:

ACCCAAAAAT	CCAATGCTGA	ATATTTGGCT	TCATTATTCC	CANATTCTTT	GATTGTCAAA	60
GGATTTAATG	TTGTCTCAGC	TTGGGCACTT	CAGTTAGGAC	CTAAGGATGC	CAGCCGGCAG	120
GTTTATATAT	GCAGCAACAA	TATTCAAGCG	CGACAACAGG	TTATTGAACT	TGCCCCGCCAG	180
TTNAATTTCA	TTCCCATTTGA	CTTGGGATCC	TTATCATCAG	CCAGAGAGAT	TGAAAATTTA	240
CCCCTACNAC	TCTTTACTCT	CTGGANAGGG	CCAGTGGTGG	TAGCTATAAG	CTTGGCCACA	300
TTTTTTTTTC	CTTTATTCCT	TTGTCAGA				328

(2) INFORMATION FOR SEQ ID NO:213:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 250 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:213:

ACTTATGAGC	AGAGCGACAT	ATCCNAGTGT	AGACTGAATA	AAACTGAATT	CTCTCCAGTT	60
TAAAGCATTG	CTCACTGAAG	GGATAGAAGT	GACTGCCAGG	AGGGAAAGTA	AGCCAAGGCT	120
CATTATGCCA	AAGGANATAT	ACATTTCAAT	TCTCCAAACT	TCTTCCTCAT	TCCAAGAGTT	180
TTCAATATTT	GCATGAACCT	GCTGATAANC	CATGTTAANA	AACAAATATC	TCTCTNACCT	240
TCTCATCGGT						250

(2) INFORMATION FOR SEQ ID NO:214:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 444 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:214:

ACCCAGAATC	CAATGCTGAA	TATTTGGCTT	CATTATTCCC	AGATTCTTTG	ATTGTCAAAG	60
GATTTAATGT	TGTCTCAGCT	TGGGCACTTC	AGTTAGGACC	TAAGGATGCC	AGCCGGCAGG	120
TTTATATATG	CAGCAACAAT	ATTCAAGCGC	GACAACAGGT	TATTGAACTT	GCCCGCCAGT	180
TGAATTTTCAT	TCCCATTGAC	TTGGGATCCT	TATCATCAGC	CANAGAGATT	GAAAATTTAC	240
CCCTACGACT	CTTTACTCTC	TGGAGAGGGC	CAGTGGTGGT	AGCTATAAGC	TTGGCCACAT	300
TTTTTTTTTCC	TTTATTCCTT	TGTCAGAGAT	GCGATTTCATC	CATATGCTAN	AAACCAACAG	360
AGTGACTTTT	ACAAAATTC	TATAGANATT	GTGAATAAAA	CCTTACCTAT	AGTTGCCATT	420
ACTTTGCTCT	CCCTAATATA	CCTC				444

(2) INFORMATION FOR SEQ ID NO:215:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 366 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:215:

ACTTATGAGC	AGAGCGACAT	ATCCAAGTGT	ANACTGAATA	AAACTGAATT	CTCTCCAGTT	60
TAAAGCATTG	CTCACTGAAG	GGATAGAAGT	GACTGCCAGG	AGGGAAAAGTA	AGCCAAGGCT	120
CATTATGCCA	AAGGANATAT	ACATTTCAAT	TCTCCAAACT	TCTTCCTCAT	TCCAAGAGTT	180
TTCAATATTT	GCATGAACCT	GCTGATAAGC	CATGTTGAGA	AACAAATATC	TCTCTGACCT	240
TCTCATCGGT	AAGCAGAGGC	TGTAGGCAAC	ATGGACCATA	GCGAANAAAA	AACTTAGTAA	300
TCCAAGCTGT	TTTCTACACT	GTAACCAGGT	TTCCAACCAA	GGTGGAATC	TCCTATACTT	360
GGTGCC						366

(2) INFORMATION FOR SEQ ID NO:216:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 260 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:216:

CTGTATAAAC	AGAACTCCAC	TGCANGAGGG	AGGGCCGGGC	CAGGAGAATC	TCCGCTTGTC	60
CAAGACAGGG	GCCTAAGGAG	GGTCTCCACA	CTGCTNNTAA	GGGCTNTTNC	ATTTTTTTAT	120
TAATAAAAAG	TNNAAGGC	CTCTTCTCAA	CTTTTTTCCC	TTNGGCTGGA	AAATTTAAAA	180
ATCAAAAATT	TCCTNAAGTT	NTCAAGCTAT	CATATATACT	NTATCCTGAA	AAAGCAACAT	240
AATTCTTCT	TCCCTCCTTT					260

(2) INFORMATION FOR SEQ ID NO:217:

(A) LENGTH: 262 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:217:

ACCTACGTGG	GTAAGTTTAN	AAATGTTATA	ATTTTCAGGAA	NAGGAACGCA	TATAATTGTA	60
TCCTTGCCAT	AATTTTCTAT	TTTAATAAGG	AAATAGCAAA	TTGGGGTGGG	GGGAATGTAG	120
GGCATTCTAC	AGTTTGAGCA	AAATGCAATT	AAATGTGGAA	GGACAGCACT	GAAAAATTTT	180
ATGAATAATC	TGTATGATTA	TATGTCCTTA	GAGTAGATTT	ATAATTAGCC	ACTTACCCTA	240
ATATCCTTCA	TGCTTGTA	GT				262

(A) LENGTH: 205 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:218:

ACCAAGGTGG	TGCATTACCG	GAANTGGATC	AANGACACCA	TCGTGGCCAA	CCCCTGAGCA	60
CCCCTATCAA	CTCCCTTTTG	TAGTAAACTT	GGAACCTTGG	AAATGACCAG	GCCAAGACTC	120
AGGCCTCCCC	AGTTCTACTG	ACCTTTGTCC	TTANGTNTNA	NGTCCAGGGT	TGCTAGGAAA	180
ANAAATCAGC	AGACACAGGT	GTAAA				205

(A) LENGTH: 114 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:219:

TACTGTTTTG TCTCAGTAAC AATAAATACA AAAAGACTGG TTGTGTTCCG GCCCATCCA 60
ACCACGAAGT TGATTTCTCT TGTGTGCAGA GTGACTGATT TTAAAGGACA TGGA 114

(A) LENGTH: 93 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:220:

ACTAGCCAGC	ACAAAAGGCA	GGGTAGCCTG	AATTGCTTTC	TGCTCTTTAC	ATTTCTTTTA	60
AAATAAGCAT	TTAGTGCTCA	GTCCCTACTG	AGT			93

(2) INFORMATION FOR SEQ ID NO:221:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 167 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:221:

ACTANGTGCA	GGTGCGCACA	AATATTTGTC	GATATTCCCT	TCATCTTGGA	TTCCATGAGG	60
TCTTTTGCCC	AGCCTGTGGC	TCTACTGTAG	TAAGTTTCTG	CTGATGAGGA	GCCAGNATGC	120
CCCCCACTAC	CTTCCCTGAC	GCTCCCCANA	AATCACCCAA	CCTCTGT		167

(2) INFORMATION FOR SEQ ID NO:222:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 351 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:222:

AGGGCGTGTT	GCGGAGGGCG	GTACTGACCT	CATTAGTAGG	AGGATGCATT	CTGGCACCCC	60
GTTCTTCACC	TGTCCCCCAA	TCCTTAAAG	GCCATACTGC	ATAAAGTCAA	CAACAGATAA	120
ATGTTTGCTG	AATTAAAGGA	TGGATGAAAA	AAATTAATAA	TGAATTTTGT	CATAATCCAA	180
TTTTCTCTTT	TATATTCTA	GAAGAAGTTT	CTTTGAGCCT	ATTAGATCCC	GGGAATCTTT	240
TAGGTGAGCA	TGATTAGAGA	GCTTGTAGGT	TGCTTTTACA	TATATCTGGC	ATATTTGAGT	300
CTCGTATCAA	AACAATAGAT	TGGTAAAGGT	GGTATTATTG	TATTGATAAG	T	351

(2) INFORMATION FOR SEQ ID NO:223

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:223:

AAAACAAACA	AACAAAAAAA	ACAATTCTTC	ATTCAGAAAA	ATTATCTTAG	GGAATGATAT	60
TGGTAATTAT	GGTCAATTTA	ATWRTRTTKT	GGGGCATTTC	CTTACATTGT	CTTGACAAGA	120
TTAAAATGTC	TGTGCCAAAA	TTTTGTATTT	TATTTGGAGA	CTTCTTATCA	AAAGTAATGC	180
TGCCAAAGGA	AGTCTAAGGA	ATTAGTAGTG	TTCCCMTCAC	TTGTTTGGAG	TGTGCTATTC	240
TAAAAGATTT	TGATTTCTCTG	GAATGACAAT	TATATTTTAA	CTTTGGTGGG	GGAAANAGTT	300
ATAGGACCAC	AGTCTTCACT	TCTGATACTT	GTAAATTAAT	CTTTTATTGC	ACTTGTTTTG	360
ACCATTAAGC	TATATGTTTA	AAA				383

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 320 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

CCCCTGAAGG	CTTCTTGTTA	GAAAATAGTA	CAGTTACAAC	CAATAGGAAC	AACAAAAAGA	60
AAAAGT'TTGT	GACATTGTAG	TAGGGAGTGT	GTACCCCTTA	CTCCCATCA	AAAAAAAAT	120
GGATACATGG	TTAAAGGATA	RAAGGGCAAT	ATTTTATCAT	ATGTTCTAA	AGAGAAGGAA	180
GAGAAAATAC	TACT'TTCTCR	AAATGGAAGC	CCTTAAAGGT	GC'TTTGATAC	TGAAGGACAC	240
AAATGTGGCC	GTCCATCCTC	CTTTARAGTT	GCATGACTTG	GACACGGTAA	CTGTTGCAGT	300
TTTARACTCM	GCATTGTGAC					320

CLAIMS

1. A method for detecting prostate cancer in a patient, comprising:
- (a) contacting a biological sample obtained from the patient with a binding agent which is capable of binding to a polypeptide, the polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, wherein said protein comprises an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences and variants of said nucleotide sequences; and
 - (b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting prostate cancer in the patient.
2. The method of claim 1 wherein the binding agent is a monoclonal antibody.
3. The method of claim 2 wherein the binding agent is a polyclonal antibody.
4. A method for monitoring the progression of prostate cancer in a patient, comprising:
- (a) contacting a biological sample obtained from the patient with a binding agent that is capable of binding to a polypeptide, said polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, wherein said protein comprises an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences and variants of said nucleotide sequences;
 - (b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent;
 - (c) repeating steps (a) and (b); and

(d) comparing the amount of polypeptide detected in steps (b) and (c) to monitor the progression of prostate cancer in the patient.

5. A monoclonal antibody that binds to a polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, wherein said protein comprises an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 8-29, 41-45, 47-52, 54-65, 70, 73, 74, 79, 81, 87, 90, 92, 93, 97, 103, 104, 107, 109-111, 115-160, 171, 173-175, 177, 181, 188, 191, 193, 194, 198, 203, 204, 207-224, the complements of said nucleotide sequences variants of said nucleotide sequences.

6. A method for inhibiting the development of prostate cancer in a patient, comprising administering to the patient a therapeutically effective amount of a monoclonal antibody according to claim 5.

7. The method of claim 6 wherein the monoclonal antibody is conjugated to a therapeutic agent.

8. A method for detecting prostate cancer in a patient comprising:

- (a) obtaining a biological sample from the patient;
- (b) contacting the sample with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotides is specific for a DNA molecule encoding a polypeptide comprising an immunogenic portion of a prostate protein or of a variant thereof, said protein comprising an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences variants of said nucleotide sequences; and
- (c) detecting in the sample a DNA sequence that amplifies in the presence of the oligonucleotide primers, thereby detecting prostate cancer.

9. The method of claim 8, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule having a

sequence selected from SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224.

10. A diagnostic kit comprising:

- (a) one or more monoclonal antibodies of claim 5; and
- (b) a detection reagent.

~~11.~~ A diagnostic kit comprising:

- ~~(a)~~ one or more monoclonal antibodies that bind to a polypeptide encoded by a DNA molecule having a nucleotide sequence selected from the group consisting of SEQ ID Nos: 5-7, 30-40, 46, 53, 66-69, 71, 72, 75-78, 80, 82-86, 88, 89, 91, 94-96, 98-102, 105, 106, 161-170, 179, 180, 182-187, 189, 190, 192, 195-197, 199-202, 205, 206, the complements of said sequences and variants of said nucleotide sequences; and
- (b) a detection reagent.

12. The kit of claims 10 or 11 wherein the monoclonal antibodies are immobilized on a solid support.

13. The kit of claim 12 wherein the solid support comprises nitrocellulose, latex or a plastic material.

14. The kit of claims 10 or 11 wherein the detection reagent comprises a reporter group conjugated to a binding agent.

15. The kit of claim 14 wherein the binding agent is selected from the group consisting of anti-immunoglobulins, Protein G, Protein A and lectins.

16. The kit of claim 14 wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.

~~17.~~ A diagnostic kit comprising at least two oligonucleotide primers, at least one of the oligonucleotide primers being specific for a DNA molecule encoding a

polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, said protein comprising an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences and variants of said nucleotide sequences.

18. A diagnostic kit of claim 17 wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule having a sequence selected from SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224.

19. A method for detecting prostate cancer in a patient, comprising:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide probe specific for a DNA molecule encoding a polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, said protein comprising an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences and variants of said nucleotide sequences; and
- (c) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe, thereby detecting prostate cancer in the patient.

20. The method of claim 19 wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule having a sequence selected from the group consisting of SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224.

21. A diagnostic kit comprising an oligonucleotide probe specific for a DNA molecule encoding a polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, said protein comprising an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences

recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences variants of said nucleotide sequences.

22. The diagnostic kit of claim 21 wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule having a sequence selected from the group consisting of SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224.

ABSTRACT OF THE DISCLOSURE

WPN/DJM/210121/428C3-AP/1

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Jiangchun Xu and Davin C. Dillon
Filed : February 25, 1998
For : COMPOUNDS FOR IMMUNODIAGNOSIS OF PROSTATE
CANCER AND METHODS FOR THEIR USE

Docket No. : 210121.428C3

Date : February 25, 1998



Box Patent Application
Assistant Commissioner for Patents
2011 Jefferson Davis Highway
Washington, D.C. 20231


DECLARATION

Sir:

I, Lawrence Teague, in accordance with 37 C.F.R. § 1.821(f) do hereby declare that, to the best of my knowledge, the content of the paper entitled "Sequence Listing" and the computer readable copy contained within the floppy disk are the same.

I declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Dated this 25th day of February, 1998.



Lawrence Teague
Legal Assistant

6300 Columbia Center
Seattle, WA 98104-7092
(206) 622-4900
FAX (206) 682-6031